



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 17, 2023 – 05:16 am GMT

PDB ID : 4A6V
Title : X-ray structures of oxazole hydroxamate EcMetAp-Mn complexes
Authors : Huguet, F.; Melet, A.; AlvesdeSousa, R.; Lieutaud, A.; Chevalier, J.; Deschamps, P.; Tomas, A.; Leulliot, N.; Pages, J.M.; Artaud, I.
Deposited on : 2011-11-09
Resolution : 1.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	265	 2% 88% 10% ..
1	B	265	 2% 89% 9% ..

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4979 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called METHIONINE AMINOPEPTIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	264	2055	1297	355	388	15	0	2	0
1	B	262	2034	1281	351	387	15	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	265	ALA	-	expression tag	UNP C6EAB7
B	265	ALA	-	expression tag	UNP C6EAB7

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mn	0	0
			2	2		
2	B	2	Total	Mn	0	0
			2	2		

- Molecule 3 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	1	3		

- Molecule 4 is N-HYDROXY-2-[2-(TRIFLUOROMETHYL)PHENYL]-1,3-OXAZOLE-4-CARBOXAMIDE (three-letter code: IKY) (formula: C₁₁H₇F₃N₂O₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	0	0
			19	11	3	2	3		
4	B	1	Total	C	F	N	O	0	0
			19	11	3	2	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	441	Total 441	O 441	0	0
5	B	403	Total 403	O 403	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: METHIONINE AMINOPEPTIDASE



- Molecule 1: METHIONINE AMINOPEPTIDASE



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	53.23Å 61.25Å 77.76Å 90.00° 107.62° 90.00°	Depositor
Resolution (Å)	74.11 – 1.46 19.96 – 1.46	Depositor EDS
% Data completeness (in resolution range)	98.1 (74.11-1.46) 98.1 (19.96-1.46)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 1.46Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.159 , 0.223 0.157 , 0.218	Depositor DCC
R_{free} test set	4030 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	14.7	Xtrriage
Anisotropy	0.038	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4979	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 77.09 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.1210e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CO3, MN, IKY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.10	3/2100 (0.1%)	1.18	13/2838 (0.5%)
1	B	1.08	1/2069 (0.0%)	1.08	7/2796 (0.3%)
All	All	1.09	4/4169 (0.1%)	1.13	20/5634 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	2	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	265	ALA	CA-CB	-7.74	1.36	1.52
1	B	168	TYR	CD1-CE1	5.74	1.48	1.39
1	A	167	GLU	CD-OE1	5.57	1.31	1.25
1	A	265	ALA	CA-C	-5.49	1.38	1.52

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	124	ARG	NE-CZ-NH2	-19.87	110.37	120.30
1	A	124	ARG	NE-CZ-NH1	15.31	127.96	120.30
1	B	127	ARG	NE-CZ-NH2	-12.15	114.22	120.30
1	B	26	GLU	OE1-CD-OE2	-7.38	114.44	123.30
1	A	242	ASP	CB-CG-OD1	7.02	124.62	118.30
1	A	264	GLU	N-CA-CB	6.86	122.94	110.60
1	A	228	ARG	NE-CZ-NH2	-6.31	117.14	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	265	ALA	N-CA-CB	6.08	118.61	110.10
1	B	127	ARG	NE-CZ-NH1	5.97	123.29	120.30
1	B	124	ARG	NE-CZ-NH1	5.94	123.27	120.30
1	A	119	THR	N-CA-CB	-5.93	99.03	110.30
1	B	124	ARG	NE-CZ-NH2	-5.88	117.36	120.30
1	A	124	ARG	CD-NE-CZ	5.64	131.50	123.60
1	A	7	THR	OG1-CB-CG2	5.52	122.70	110.00
1	B	189	ARG	CG-CD-NE	-5.31	100.64	111.80
1	A	177	PHE	CB-CG-CD1	5.28	124.50	120.80
1	A	43	ARG	NE-CZ-NH1	5.17	122.88	120.30
1	A	166	ARG	NE-CZ-NH1	-5.04	117.78	120.30
1	B	189	ARG	NE-CZ-NH1	-5.03	117.78	120.30
1	A	124	ARG	CG-CD-NE	-5.01	101.27	111.80

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	7	THR	CB
1	A	264	GLU	CA

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	264	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2055	0	2073	42	0
1	B	2034	0	2053	36	2
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	4	0	0	0	0
4	A	19	0	6	0	0
4	B	19	0	6	0	0
5	A	441	0	0	21	3

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	403	0	0	17	1
All	All	4979	0	4138	75	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:THR:HB	5:A:2263:HOH:O	1.26	1.26
1:B:26:GLU:CD	5:B:2066:HOH:O	1.75	1.25
1:B:26:GLU:HG2	5:B:2066:HOH:O	1.38	1.18
5:A:2321:HOH:O	1:B:218:LYS:HD3	1.42	1.17
1:B:26:GLU:CG	5:B:2066:HOH:O	1.83	1.15
1:A:265:ALA:O	1:B:177:PHE:HB3	1.56	1.05
1:B:37:SER:OG	1:B:40:GLU:HG3	1.63	0.98
1:B:26:GLU:OE2	5:B:2066:HOH:O	1.73	0.97
1:A:265:ALA:HB1	1:B:62:TYR:OH	1.74	0.88
1:B:121:MET:HE1	1:B:162:PHE:HE1	1.38	0.88
1:A:265:ALA:HB2	5:A:2438:HOH:O	1.73	0.87
1:A:120:ILE:HD12	5:A:2268:HOH:O	1.73	0.86
1:A:119:THR:HG23	1:A:122:GLY:HA3	1.59	0.84
1:A:119:THR:CG2	1:A:122:GLY:HA3	2.09	0.83
1:A:2:ALA:N	5:A:2002:HOH:O	2.13	0.81
1:B:121:MET:CE	1:B:162:PHE:HE1	1.92	0.81
1:A:120:ILE:HB	5:A:2268:HOH:O	1.80	0.80
1:A:123:GLU:OE1	1:A:127:ARG:NH1	2.13	0.80
1:A:86:LYS:HE3	5:A:2215:HOH:O	1.83	0.78
1:B:263:ASP:OD2	5:B:2400:HOH:O	2.01	0.78
1:A:192:ASN:HB3	5:A:2300:HOH:O	1.82	0.77
1:B:86:LYS:HE3	5:B:2203:HOH:O	1.85	0.77
1:A:123:GLU:OE2	1:A:127:ARG:NH1	2.18	0.76
1:A:52:GLU:HG3	5:A:2139:HOH:O	1.85	0.75
1:A:252:LYS:HG2	5:A:2425:HOH:O	1.84	0.75
1:B:40:GLU:HB3	5:B:2103:HOH:O	1.86	0.75
1:B:37:SER:OG	1:B:40:GLU:CG	2.34	0.75
1:B:121:MET:CE	1:B:162:PHE:CE1	2.71	0.73
1:A:184:LEU:HD13	5:A:2353:HOH:O	1.90	0.71
1:A:62:TYR:CE2	1:A:63[B]:HIS:HD2	2.09	0.71
1:A:123:GLU:CD	1:A:127:ARG:NH1	2.47	0.68
1:A:7:THR:HG22	1:A:10:ASP:CG	2.14	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:26:GLU:OE2	5:B:2065:HOH:O	2.14	0.65
1:B:121:MET:HE1	1:B:162:PHE:CE1	2.25	0.64
1:A:192:ASN:CB	5:A:2300:HOH:O	2.41	0.63
1:A:246:GLU:HG3	1:A:259:ILE:CD1	2.29	0.62
1:B:121:MET:HE3	1:B:162:PHE:CE1	2.34	0.62
1:A:119:THR:CA	5:A:2263:HOH:O	2.45	0.61
1:A:7:THR:CG2	1:A:10:ASP:H	2.15	0.60
1:A:120:ILE:CG1	5:A:2268:HOH:O	2.49	0.60
1:B:215:ARG:NH2	1:B:226:LYS:HE2	2.16	0.59
1:A:119:THR:HG23	1:A:122:GLY:CA	2.30	0.58
1:B:2:ALA:HB3	5:B:2001:HOH:O	2.02	0.58
1:A:246:GLU:HG3	1:A:259:ILE:HD13	1.85	0.57
1:A:127:ARG:NH2	5:A:2276:HOH:O	2.35	0.57
1:A:119:THR:CB	5:A:2263:HOH:O	2.06	0.56
1:A:7:THR:HG23	1:A:10:ASP:H	1.68	0.56
1:B:86:LYS:NZ	5:B:2205:HOH:O	2.38	0.56
1:A:81:ILE:HD11	1:A:221:TRP:HB2	1.88	0.56
1:A:120:ILE:CD1	5:A:2268:HOH:O	2.43	0.56
1:A:119:THR:HG21	1:A:122:GLY:HA3	1.86	0.54
1:B:246:GLU:HG3	1:B:259:ILE:HD13	1.91	0.53
1:A:252:LYS:CG	5:A:2425:HOH:O	2.47	0.53
1:B:218:LYS:HG3	5:B:2365:HOH:O	2.09	0.52
1:A:180:GLU:HG2	5:A:2351:HOH:O	2.09	0.52
1:A:184:LEU:HD12	1:A:184:LEU:N	2.26	0.50
1:B:121:MET:HE2	5:B:2154:HOH:O	2.11	0.50
1:B:43:ARG:HD3	5:B:2054:HOH:O	2.13	0.48
1:B:43:ARG:HG2	5:B:2046:HOH:O	2.14	0.47
1:B:246:GLU:HG3	1:B:259:ILE:CD1	2.46	0.46
1:A:62:TYR:CE2	1:A:63[B]:HIS:CD2	2.99	0.45
1:A:7:THR:HG21	5:A:2026:HOH:O	2.16	0.45
1:B:189:ARG:HD3	5:B:2334:HOH:O	2.17	0.44
1:A:180:GLU:CG	5:A:2351:HOH:O	2.63	0.44
1:A:146:LEU:HD23	1:A:146:LEU:HA	1.76	0.43
1:A:264:GLU:HA	1:A:265:ALA:HA	1.81	0.43
1:B:215:ARG:HH21	1:B:226:LYS:HE2	1.82	0.42
1:B:189:ARG:NH1	1:B:189:ARG:HG3	2.34	0.42
1:B:54:HIS:HD2	5:B:2101:HOH:O	2.03	0.42
1:A:144:ILE:HG12	1:A:145:ASN:N	2.35	0.41
1:B:184:LEU:HD13	5:B:2338:HOH:O	2.20	0.41
1:A:265:ALA:C	1:B:178:HIS:ND1	2.74	0.41
1:B:250:LEU:HD22	1:B:258:ALA:HB2	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:184:LEU:HD12	1:B:184:LEU:N	2.36	0.40
1:B:116:GLY:O	1:B:118:PRO:HD3	2.21	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:189:ARG:NH1	5:A:2436:HOH:O[2_556]	1.75	0.45
1:B:189:ARG:NH1	5:A:2439:HOH:O[2_556]	1.88	0.32
5:A:2347:HOH:O	5:B:2402:HOH:O[1_554]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/265 (100%)	257 (97%)	7 (3%)	1 (0%)	34	13
1	B	260/265 (98%)	252 (97%)	8 (3%)	0	100	100
All	All	525/530 (99%)	509 (97%)	15 (3%)	1 (0%)	47	22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	264	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/228 (100%)	225 (99%)	2 (1%)	78	57
1	B	226/228 (99%)	224 (99%)	2 (1%)	78	57
All	All	453/456 (99%)	449 (99%)	4 (1%)	78	57

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	119	THR
1	A	177	PHE
1	B	40	GLU
1	B	177	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	46	ASN
1	B	46	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	IKY	A	1268	2	16,20,20	1.79	4 (25%)	19,29,29	1.89	4 (21%)
3	CO3	A	1267	-	2,3,3	0.12	0	2,3,3	0.20	0
4	IKY	B	1264	2	16,20,20	2.48	3 (18%)	19,29,29	1.81	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IKY	A	1268	2	-	0/8/16/16	0/2/2/2
4	IKY	B	1264	2	-	0/8/16/16	0/2/2/2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1264	IKY	OAB-NAL	-7.89	1.20	1.40
4	A	1268	IKY	CAN-NAL	4.40	1.38	1.32
4	A	1268	IKY	CAO-NAK	-3.51	1.26	1.37
4	B	1264	IKY	CAO-NAK	-3.25	1.27	1.37
4	A	1268	IKY	OAB-NAL	-2.58	1.33	1.40
4	A	1268	IKY	CAQ-NAK	-2.11	1.29	1.35
4	B	1264	IKY	CAI-CAR	2.01	1.42	1.39

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1268	IKY	CAP-CAQ-NAK	4.85	130.04	123.52
4	A	1268	IKY	CAH-CAP-CAR	4.12	121.92	118.47
4	B	1264	IKY	CAH-CAP-CAR	4.12	121.92	118.47
4	B	1264	IKY	CAP-CAQ-NAK	3.78	128.61	123.52
4	B	1264	IKY	FAC-CAS-CAR	-3.40	106.78	112.70
4	A	1268	IKY	OAA-CAN-NAL	-3.08	117.19	122.94
4	A	1268	IKY	FAD-CAS-CAR	-2.21	108.85	112.70


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/265 (99%)	-0.03	4 (1%) 73 74	8, 16, 25, 30	0
1	B	262/265 (98%)	0.00	4 (1%) 73 74	9, 16, 27, 39	0
All	All	526/530 (99%)	-0.01	8 (1%) 73 74	8, 16, 26, 39	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	263	ASP	5.6
1	B	2	ALA	4.9
1	A	2	ALA	4.7
1	A	265	ALA	3.7
1	B	117	LYS	3.3
1	A	192	ASN	3.0
1	B	243	ASN	2.4
1	A	120	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	CO3	A	1267	4/4	0.91	0.13	17,18,20,21	0
4	IKY	A	1268	19/19	0.96	0.07	12,14,17,18	0
4	IKY	B	1264	19/19	0.97	0.08	12,16,20,20	0
2	MN	B	1266	1/1	1.00	0.05	9,9,9,9	0
2	MN	A	1266	1/1	1.00	0.06	9,9,9,9	0
2	MN	A	1269	1/1	1.00	0.06	9,9,9,9	0
2	MN	B	1265	1/1	1.00	0.05	12,12,12,12	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.