



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 27, 2024 – 05:37 PM EST

PDB ID : 1D0Q
Title : STRUCTURE OF THE ZINC-BINDING DOMAIN OF BACILLUS
STEAROTHERMOPHILUS DNA PRIMASE
Authors : Pan, H.; Wigley, D.B.
Deposited on : 1999-09-14
Resolution : 1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5722 (1.74-1.70)
Clashscore	141614	6152 (1.74-1.70)
Ramachandran outliers	138981	6051 (1.74-1.70)
Sidechain outliers	138945	6051 (1.74-1.70)
RSRZ outliers	127900	5629 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	103	 4% (Poor fit), 68% (0 outliers), 20% (1 outlier), 8% (2 outliers), .. (3+ outliers)
1	B	103	 2% (Poor fit), 70% (0 outliers), 18% (1 outlier), 10% (2 outliers), .. (3+ outliers)

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1746 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA PRIMASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	102	791	506	136	145	4	0	0	0
1	B	102	791	506	136	145	4	0	0	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	87	Total	O	0	0
			87	87		
3	B	75	Total	O	0	0
			75	75		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA PRIMASE



- Molecule 1: DNA PRIMASE



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	36.17Å 58.74Å 45.75Å 90.00° 91.73° 90.00°	Depositor
Resolution (Å)	15.00 – 1.71 14.75 – 1.65	Depositor EDS
% Data completeness (in resolution range)	(Not available) (15.00-1.71) 95.8 (14.75-1.65)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 1.65Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.204 , 0.225 0.194 , 0.212	Depositor DCC
R_{free} test set	1132 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	14.0	Xtrriage
Anisotropy	0.364	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 57.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.064 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1746	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/809	2.44	36/1090 (3.3%)
1	B	0.51	0/809	2.15	31/1090 (2.8%)
All	All	0.53	0/1618	2.30	67/2180 (3.1%)

There are no bond length outliers.

All (67) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	88	ARG	CD-NE-CZ	29.77	165.28	123.60
1	B	102	LEU	C-N-CA	19.06	169.35	121.70
1	A	14	ARG	NE-CZ-NH1	17.72	129.16	120.30
1	B	96	ASP	CB-CG-OD1	17.24	133.82	118.30
1	A	88	ARG	NE-CZ-NH2	-13.84	113.38	120.30
1	A	43	HIS	CG-ND1-CE1	13.35	126.89	108.20
1	A	102	LEU	O-C-N	-12.28	103.05	122.70
1	A	102	LEU	C-N-CA	12.22	152.26	121.70
1	A	14	ARG	NH1-CZ-NH2	-11.29	106.98	119.40
1	A	88	ARG	NH1-CZ-NH2	10.53	130.99	119.40
1	B	15	ARG	NE-CZ-NH1	-10.46	115.07	120.30
1	A	76	ASP	CB-CG-OD2	-10.32	109.01	118.30
1	B	36	TYR	CB-CG-CD1	-10.00	115.00	121.00
1	B	102	LEU	O-C-N	-9.96	106.76	122.70
1	B	36	TYR	CB-CG-CD2	9.53	126.72	121.00
1	A	76	ASP	CB-CG-OD1	9.46	126.82	118.30
1	A	88	ARG	NE-CZ-NH1	-9.34	115.63	120.30
1	B	8	GLU	OE1-CD-OE2	9.03	134.14	123.30
1	A	3	HIS	C-N-CA	9.02	144.25	121.70
1	A	43	HIS	ND1-CG-CD2	-8.72	93.79	106.00
1	B	88	ARG	NE-CZ-NH2	-8.67	115.97	120.30
1	A	21	ASP	CB-CG-OD1	8.45	125.90	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	88	ARG	CB-CG-CD	7.93	132.21	111.60
1	B	102	LEU	CA-C-O	7.87	136.63	120.10
1	A	30	LYS	CA-CB-CG	7.75	130.46	113.40
1	B	32	GLN	CA-CB-CG	7.28	129.41	113.40
1	A	33	GLY	O-C-N	7.20	134.22	122.70
1	B	18	ASP	CB-CG-OD1	7.18	124.76	118.30
1	A	14	ARG	NE-CZ-NH2	7.11	123.85	120.30
1	B	34	ARG	NE-CZ-NH1	7.06	123.83	120.30
1	A	34	ARG	NE-CZ-NH1	6.79	123.70	120.30
1	B	96	ASP	OD1-CG-OD2	-6.75	110.47	123.30
1	A	3	HIS	O-C-N	-6.72	111.94	122.70
1	A	102	LEU	CA-C-O	6.58	133.92	120.10
1	A	43	HIS	ND1-CE1-NE2	-6.54	95.50	109.90
1	B	14	ARG	NE-CZ-NH1	-6.46	117.07	120.30
1	B	88	ARG	NH1-CZ-NH2	6.28	126.31	119.40
1	A	3	HIS	N-CA-CB	-6.15	99.53	110.60
1	B	15	ARG	CD-NE-CZ	6.13	132.19	123.60
1	A	51	SER	N-CA-CB	-6.11	101.34	110.50
1	B	8	GLU	CG-CD-OE2	-6.05	106.19	118.30
1	A	33	GLY	CA-C-O	-5.88	110.03	120.60
1	B	87	LYS	CA-CB-CG	5.84	126.26	113.40
1	A	27	VAL	CG1-CB-CG2	-5.77	101.66	110.90
1	A	3	HIS	CA-C-O	5.75	132.18	120.10
1	B	15	ARG	NH1-CZ-NH2	5.70	125.67	119.40
1	B	26	TYR	CZ-CE2-CD2	-5.70	114.67	119.80
1	A	50	PHE	CB-CG-CD1	-5.60	116.88	120.80
1	A	3	HIS	CA-CB-CG	5.60	123.12	113.60
1	A	20	VAL	CA-CB-CG2	-5.60	102.50	110.90
1	A	50	PHE	CB-CG-CD2	5.58	124.71	120.80
1	B	95	VAL	CG1-CB-CG2	-5.54	102.04	110.90
1	A	103	ASP	CB-CG-OD1	5.51	123.26	118.30
1	B	103	ASP	CA-CB-CG	-5.47	101.37	113.40
1	B	24	GLY	O-C-N	-5.39	114.07	122.70
1	A	84	GLU	OE1-CD-OE2	5.28	129.64	123.30
1	B	36	TYR	N-CA-CB	5.26	120.08	110.60
1	A	2	GLY	N-CA-C	-5.25	99.99	113.10
1	B	85	ALA	O-C-N	-5.23	114.33	122.70
1	B	32	GLN	CA-C-O	5.19	131.00	120.10
1	B	36	TYR	O-C-N	-5.17	114.43	122.70
1	A	68	GLY	O-C-N	5.15	130.94	122.70
1	B	88	ARG	NE-CZ-NH1	-5.15	117.73	120.30
1	B	14	ARG	NH1-CZ-NH2	5.11	125.02	119.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	43	HIS	O-C-N	5.11	131.88	123.20
1	A	34	ARG	NH1-CZ-NH2	-5.10	113.79	119.40
1	B	4	ARG	NE-CZ-NH2	-5.07	117.77	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	791	0	770	16	0
1	B	791	0	770	12	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	87	0	0	4	0
3	B	75	0	0	0	0
All	All	1746	0	1540	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:19:ILE:H	1:B:57:GLN:HE22	1.17	0.89
1:A:19:ILE:H	1:A:57:GLN:HE22	1.31	0.78
1:A:3:HIS:NE2	1:A:5:ILE:HG13	2.11	0.64
1:B:19:ILE:H	1:B:57:GLN:NE2	1.93	0.59
1:A:19:ILE:H	1:A:57:GLN:NE2	2.00	0.57
1:B:19:ILE:N	1:B:57:GLN:HE22	1.97	0.56
1:B:32:GLN:HB3	1:B:37:PHE:CD1	2.40	0.56
1:B:87:LYS:HE2	1:B:88:ARG:HG2	1.88	0.56
1:A:3:HIS:CE1	1:A:4:ARG:HB2	2.44	0.53
1:A:2:GLY:N	1:B:96:ASP:OD2	2.44	0.50
1:B:43:HIS:CG	1:B:64:CYS:HB3	2.47	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:29:LEU:HB3	1:B:36:TYR:HB3	1.95	0.49
1:A:3:HIS:CG	1:A:4:ARG:N	2.81	0.48
1:A:88:ARG:HD2	3:A:331:HOH:O	2.14	0.48
1:A:3:HIS:CG	1:A:4:ARG:H	2.32	0.47
1:A:59:PHE:CE2	1:A:73:PHE:HB2	2.50	0.47
1:A:74:LEU:HA	1:A:77:ILE:HG22	1.96	0.46
1:A:30:LYS:HG3	3:A:249:HOH:O	2.16	0.46
1:A:30:LYS:HG2	3:A:208:HOH:O	2.15	0.46
1:A:67:GLY:O	1:B:4:ARG:HA	2.18	0.44
1:A:19:ILE:N	1:A:57:GLN:HE22	2.06	0.44
1:A:11:GLU:HG2	3:A:218:HOH:O	2.18	0.43
1:A:14:ARG:NH2	1:B:101:GLU:OE2	2.51	0.42
1:B:31:ARG:HD3	1:B:33:GLY:O	2.21	0.41
1:B:74:LEU:HD11	1:B:89:LEU:HD13	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	100/103 (97%)	95 (95%)	2 (2%)	3 (3%)	4	0
1	B	100/103 (97%)	97 (97%)	3 (3%)	0	100	100
All	All	200/206 (97%)	192 (96%)	5 (2%)	3 (2%)	10	2

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	102	LEU
1	A	3	HIS
1	A	4	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	82/83 (99%)	71 (87%)	11 (13%)	4 0
1	B	82/83 (99%)	75 (92%)	7 (8%)	10 2
All	All	164/166 (99%)	146 (89%)	18 (11%)	6 1

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	HIS
1	A	11	GLU
1	A	14	ARG
1	A	15	ARG
1	A	30	LYS
1	A	32	GLN
1	A	34	ARG
1	A	43	HIS
1	A	87	LYS
1	A	89	LEU
1	A	103	ASP
1	B	30	LYS
1	B	31	ARG
1	B	45	GLU
1	B	87	LYS
1	B	89	LEU
1	B	102	LEU
1	B	103	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	GLN
1	B	3	HIS
1	B	35	ASN
1	B	57	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	102/103 (99%)	0.18	4 (3%) 39 44	15, 23, 39, 69	0
1	B	102/103 (99%)	0.10	2 (1%) 65 69	17, 22, 39, 75	0
All	All	204/206 (99%)	0.14	6 (2%) 51 56	15, 23, 39, 75	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	103	ASP	7.6
1	A	103	ASP	5.8
1	A	3	HIS	4.1
1	A	2	GLY	3.5
1	B	102	LEU	2.8
1	A	102	LEU	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	B	151	1/1	0.98	0.08	15,15,15,15	0
2	ZN	A	151	1/1	0.99	0.07	15,15,15,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.