



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 19, 2022 – 01:00 PM EST

PDB ID : 8DFK
Title : X-ray crystal structure of Bacillus subtilis ComEA
Authors : Ahmed, I.; Khaja, F.T.; Neiditch, M.B.
Deposited on : 2022-06-22
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.31.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	157	
1	B	157	
1	C	157	
1	D	157	
1	E	157	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	157	 2% 33% 8% 59%
1	G	157	 2% 34% 6% 60%

2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 6679 atoms, of which 3320 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ComE operon protein 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	64	956	296	475	83	101	1	0	0	0
1	B	64	956	296	475	83	101	1	0	0	0
1	C	64	956	296	475	83	101	1	0	0	0
1	D	64	956	296	475	83	101	1	0	0	0
1	E	64	956	296	475	83	101	1	0	0	0
1	F	64	956	296	475	83	101	1	0	0	0
1	G	63	943	292	470	81	99	1	0	0	0

There are 63 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	57	MET	-	initiating methionine	UNP P39694
A	206	LEU	-	expression tag	UNP P39694
A	207	GLU	-	expression tag	UNP P39694
A	208	HIS	-	expression tag	UNP P39694
A	209	HIS	-	expression tag	UNP P39694
A	210	HIS	-	expression tag	UNP P39694
A	211	HIS	-	expression tag	UNP P39694
A	212	HIS	-	expression tag	UNP P39694
A	213	HIS	-	expression tag	UNP P39694
B	57	MET	-	initiating methionine	UNP P39694
B	206	LEU	-	expression tag	UNP P39694
B	207	GLU	-	expression tag	UNP P39694
B	208	HIS	-	expression tag	UNP P39694
B	209	HIS	-	expression tag	UNP P39694
B	210	HIS	-	expression tag	UNP P39694

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	211	HIS	-	expression tag	UNP P39694
B	212	HIS	-	expression tag	UNP P39694
B	213	HIS	-	expression tag	UNP P39694
C	57	MET	-	initiating methionine	UNP P39694
C	206	LEU	-	expression tag	UNP P39694
C	207	GLU	-	expression tag	UNP P39694
C	208	HIS	-	expression tag	UNP P39694
C	209	HIS	-	expression tag	UNP P39694
C	210	HIS	-	expression tag	UNP P39694
C	211	HIS	-	expression tag	UNP P39694
C	212	HIS	-	expression tag	UNP P39694
C	213	HIS	-	expression tag	UNP P39694
D	57	MET	-	initiating methionine	UNP P39694
D	206	LEU	-	expression tag	UNP P39694
D	207	GLU	-	expression tag	UNP P39694
D	208	HIS	-	expression tag	UNP P39694
D	209	HIS	-	expression tag	UNP P39694
D	210	HIS	-	expression tag	UNP P39694
D	211	HIS	-	expression tag	UNP P39694
D	212	HIS	-	expression tag	UNP P39694
D	213	HIS	-	expression tag	UNP P39694
E	57	MET	-	initiating methionine	UNP P39694
E	206	LEU	-	expression tag	UNP P39694
E	207	GLU	-	expression tag	UNP P39694
E	208	HIS	-	expression tag	UNP P39694
E	209	HIS	-	expression tag	UNP P39694
E	210	HIS	-	expression tag	UNP P39694
E	211	HIS	-	expression tag	UNP P39694
E	212	HIS	-	expression tag	UNP P39694
E	213	HIS	-	expression tag	UNP P39694
F	57	MET	-	initiating methionine	UNP P39694
F	206	LEU	-	expression tag	UNP P39694
F	207	GLU	-	expression tag	UNP P39694
F	208	HIS	-	expression tag	UNP P39694
F	209	HIS	-	expression tag	UNP P39694
F	210	HIS	-	expression tag	UNP P39694
F	211	HIS	-	expression tag	UNP P39694
F	212	HIS	-	expression tag	UNP P39694
F	213	HIS	-	expression tag	UNP P39694
G	57	MET	-	initiating methionine	UNP P39694
G	206	LEU	-	expression tag	UNP P39694
G	207	GLU	-	expression tag	UNP P39694


Continued on next page...

Continued from previous page...



Chain	Residue	Modelled	Actual	Comment	Reference
G	208	HIS	-	expression tag	UNP P39694
G	209	HIS	-	expression tag	UNP P39694
G	210	HIS	-	expression tag	UNP P39694
G	211	HIS	-	expression tag	UNP P39694
G	212	HIS	-	expression tag	UNP P39694
G	213	HIS	-	expression tag	UNP P39694





● Molecule 1: ComE operon protein 1



● Molecule 1: ComE operon protein 1



● Molecule 1: ComE operon protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	114.53Å 116.48Å 183.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.01 – 3.20 36.01 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.1 (36.01-3.20) 99.1 (36.01-3.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 3.18Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.254 , 0.276 0.254 , 0.274	Depositor DCC
R_{free} test set	1991 reflections (9.74%)	wwPDB-VP
Wilson B-factor (Å ²)	94.0	Xtrriage
Anisotropy	0.193	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.000 for k,h,-l	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	6679	wwPDB-VP
Average B, all atoms (Å ²)	100.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/485	0.50	0/655
1	B	0.25	0/485	0.51	0/655
1	C	0.27	0/485	0.51	0/655
1	D	0.27	0/485	0.51	0/655
1	E	0.27	0/485	0.50	0/655
1	F	0.25	0/485	0.50	0/655
1	G	0.26	0/477	0.50	0/644
All	All	0.26	0/3387	0.50	0/4574

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	481	475	475	8	0
1	B	481	475	475	18	0
1	C	481	475	475	9	0
1	D	481	475	475	14	0
1	E	481	475	475	9	0
1	F	481	475	475	8	0
1	G	473	470	469	7	0
All	All	3359	3320	3319	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 62 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:65:ASP:OD2	1:A:67:LYS:NZ	2.30	0.65
1:B:60:GLU:HG3	1:B:80:THR:HG21	1.80	0.63
1:D:105:LEU:HD13	1:E:67:LYS:NZ	2.12	0.63
1:F:62:ILE:HD12	1:F:80:THR:HA	1.82	0.61
1:B:105:LEU:N	1:C:65:ASP:OD2	2.30	0.60

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	62/157 (40%)	59 (95%)	3 (5%)	0	100	100
1	B	62/157 (40%)	60 (97%)	2 (3%)	0	100	100
1	C	62/157 (40%)	59 (95%)	3 (5%)	0	100	100
1	D	62/157 (40%)	59 (95%)	3 (5%)	0	100	100
1	E	62/157 (40%)	59 (95%)	3 (5%)	0	100	100
1	F	62/157 (40%)	60 (97%)	2 (3%)	0	100	100
1	G	61/157 (39%)	59 (97%)	2 (3%)	0	100	100
All	All	433/1099 (39%)	415 (96%)	18 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	51/125 (41%)	51 (100%)	0	100	100
1	B	51/125 (41%)	51 (100%)	0	100	100
1	C	51/125 (41%)	49 (96%)	2 (4%)	32	67
1	D	51/125 (41%)	49 (96%)	2 (4%)	32	67
1	E	51/125 (41%)	51 (100%)	0	100	100
1	F	51/125 (41%)	50 (98%)	1 (2%)	55	80
1	G	50/125 (40%)	49 (98%)	1 (2%)	55	80
All	All	356/875 (41%)	350 (98%)	6 (2%)	60	83

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	102	GLN
1	F	85	SER
1	G	85	SER
1	C	85	SER
1	C	62	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	64/157 (40%)	0.28	1 (1%) 72 59	67, 91, 118, 124	0
1	B	64/157 (40%)	0.37	1 (1%) 72 59	60, 83, 114, 123	0
1	C	64/157 (40%)	0.45	3 (4%) 31 19	59, 90, 125, 140	0
1	D	64/157 (40%)	0.35	2 (3%) 49 32	58, 79, 115, 122	0
1	E	64/157 (40%)	0.36	0 100 100	61, 88, 120, 133	0
1	F	64/157 (40%)	0.47	3 (4%) 31 19	69, 92, 122, 144	0
1	G	63/157 (40%)	0.30	1 (1%) 72 59	62, 89, 116, 128	0
All	All	447/1099 (40%)	0.37	11 (2%) 57 43	58, 88, 122, 144	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	65	ASP	3.6
1	C	122	GLU	2.5
1	F	119	LYS	2.5
1	D	65	ASP	2.4
1	C	118	PRO	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.