



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 06:55 am BST

PDB ID : 5FA0
Title : The structure of the beta-3-deoxy-D-manno-oct-2-ulosonic acid transferase domain from WbbB
Authors : Mallette, E.; Ovchinnikova, O.G.; Whitfield, C.; Kimber, M.S.
Deposited on : 2015-12-10
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	410	 85% 11% • •
1	B	410	 86% 9% 5%

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 6801 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative N-acetyl glucosaminyl transferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	394	3140	2012	519	596	7	6	0	1	0
1	B	388	3098	1991	505	590	7	5	0	5	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MSE	-	initiating methionine	UNP Q6U8B0
A	1	GLY	-	expression tag	UNP Q6U8B0
A	402	LEU	-	expression tag	UNP Q6U8B0
A	403	GLU	-	expression tag	UNP Q6U8B0
A	404	HIS	-	expression tag	UNP Q6U8B0
A	405	HIS	-	expression tag	UNP Q6U8B0
A	406	HIS	-	expression tag	UNP Q6U8B0
A	407	HIS	-	expression tag	UNP Q6U8B0
A	408	HIS	-	expression tag	UNP Q6U8B0
A	409	HIS	-	expression tag	UNP Q6U8B0
B	0	MSE	-	initiating methionine	UNP Q6U8B0
B	1	GLY	-	expression tag	UNP Q6U8B0
B	402	LEU	-	expression tag	UNP Q6U8B0
B	403	GLU	-	expression tag	UNP Q6U8B0
B	404	HIS	-	expression tag	UNP Q6U8B0
B	405	HIS	-	expression tag	UNP Q6U8B0
B	406	HIS	-	expression tag	UNP Q6U8B0
B	407	HIS	-	expression tag	UNP Q6U8B0
B	408	HIS	-	expression tag	UNP Q6U8B0
B	409	HIS	-	expression tag	UNP Q6U8B0

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	306	Total O 306 306	0	0
3	B	256	Total O 256 256	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative N-acetyl glucosaminyl transferase



- Molecule 1: Putative N-acetyl glucosaminyl transferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	82.93Å 82.93Å 120.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.47 – 2.30 48.80 – 1.88	Depositor EDS
% Data completeness (in resolution range)	100.0 (41.47-2.30) 96.8 (48.80-1.88)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 1.88Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.185 , 0.229 0.187 , 0.229	Depositor DCC
R_{free} test set	2011 reflections (3.04%)	wwPDB-VP
Wilson B-factor (Å ²)	18.8	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 44.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.034 for k,h,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6801	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.49 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.5948e-03.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/3215	0.39	0/4355
1	B	0.23	0/3183	0.38	0/4315
All	All	0.23	0/6398	0.38	0/8670

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3140	0	3025	30	0
1	B	3098	0	3010	29	0
2	A	1	0	0	1	0
3	A	306	0	0	9	1
3	B	256	0	0	10	1
All	All	6801	0	6035	56	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:ASP:OD1	3:A:601:HOH:O	1.93	0.85
1:B:97:ASP:OD2	3:B:501:HOH:O	1.98	0.81
2:A:501:CL:CL	3:A:867:HOH:O	2.38	0.79
1:A:255:TYR:OH	1:A:294:GLU:OE2	2.01	0.76
1:B:368:GLU:OE1	3:B:502:HOH:O	2.10	0.68
1:B:67:GLU:OE2	3:B:503:HOH:O	2.13	0.66
1:A:125:LYS:NZ	3:A:607:HOH:O	2.26	0.65
1:B:182:THR:O	3:B:505:HOH:O	2.14	0.64
1:B:51:ARG:NH1	3:B:506:HOH:O	2.19	0.64
1:A:99:ILE:HG23	1:A:236:ILE:HD12	1.82	0.61
1:B:163:ARG:NH2	3:B:515:HOH:O	2.30	0.61
1:B:370:VAL:HG21	1:B:392:LEU:HD21	1.82	0.61
1:B:58:PRO:HA	1:B:61:MSE:HE2	1.83	0.58
1:A:96:HIS:ND1	3:A:603:HOH:O	2.17	0.57
1:B:202:GLY:O	1:B:206:ASN:ND2	2.37	0.56
1:A:148:ARG:NH2	3:A:618:HOH:O	2.38	0.56
1:A:382:PRO:HB3	1:A:387:GLN:HG2	1.88	0.56
1:A:210:GLU:OE1	3:A:605:HOH:O	2.18	0.55
1:A:202:GLY:O	1:A:206:ASN:ND2	2.32	0.53
1:A:271:ARG:HH11	1:B:271:ARG:NE	2.07	0.52
1:A:12:ARG:NH2	1:A:231:ASP:OD1	2.42	0.52
1:A:378:SER:OG	3:A:602:HOH:O	2.09	0.52
1:A:122:ASP:OD1	1:A:125:LYS:NZ	2.41	0.51
1:B:343:SER:OG	1:B:366:GLU:OE1	2.29	0.49
1:A:381:MSE:SE	1:A:382:PRO:HD2	2.63	0.49
1:A:407:HIS:N	3:A:622:HOH:O	2.45	0.48
1:B:263:ARG:NE	3:B:504:HOH:O	2.14	0.48
1:B:158:GLU:HB3	1:B:178:VAL:HG21	1.95	0.48
1:B:112:ASP:O	3:B:508:HOH:O	2.20	0.47
1:B:155:MSE:HE3	1:B:361:LEU:HD13	1.95	0.47
1:B:182:THR:HG21	1:B:320:SER:HB3	1.96	0.46
1:B:159:ILE:CG2	1:B:163:ARG:HD2	2.46	0.46
1:B:199:ALA:HB1	1:B:203:ASP:HB2	1.97	0.45
1:A:271:ARG:NH1	1:B:271:ARG:HH21	2.15	0.45
1:A:343:SER:OG	1:A:366:GLU:OE1	2.35	0.45
1:B:354[B]:GLN:HG2	1:B:358:LYS:HE2	1.98	0.45
1:A:124:LYS:O	1:A:128:ILE:HG12	2.16	0.44
1:A:23:TYR:HH	1:A:45:THR:HG1	1.59	0.44
1:A:18:TYR:CD1	1:A:137:ILE:HD13	2.53	0.44
1:B:12:ARG:NH2	1:B:231:ASP:OD1	2.51	0.44
1:A:282:SER:HB3	1:A:287:ALA:HB3	2.00	0.44
1:A:225:VAL:HG22	1:A:251:VAL:HG21	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:124:LYS:O	1:B:128:ILE:HG12	2.18	0.43
1:A:271:ARG:HH11	1:B:271:ARG:CZ	2.32	0.43
1:A:8:PRO:HB3	1:A:38:GLY:HA3	2.01	0.43
1:B:161:PRO:HD3	1:B:172:TYR:CZ	2.54	0.43
1:B:235:LEU:O	3:B:509:HOH:O	2.22	0.42
1:A:158:GLU:HB3	1:A:178:VAL:HG21	2.02	0.42
1:A:271:ARG:NH1	1:B:271:ARG:HE	2.17	0.42
1:A:209:LEU:HD21	1:A:286:LEU:HD11	2.01	0.42
1:B:33:ILE:O	1:B:77:HIS:HB3	2.20	0.41
1:A:254:ARG:NH1	3:A:631:HOH:O	2.53	0.41
1:B:226:VAL:O	3:B:510:HOH:O	2.22	0.41
1:A:195:PHE:CE1	1:A:392:LEU:HD21	2.56	0.41
1:A:53:ARG:HB3	1:A:55:GLU:OE1	2.21	0.41
1:B:159:ILE:HG23	1:B:163:ARG:HD2	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:855:HOH:O	3:B:749:HOH:O[1_655]	1.89	0.31

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	389/410 (95%)	375 (96%)	11 (3%)	3 (1%)	19	23
1	B	387/410 (94%)	376 (97%)	11 (3%)	0	100	100
All	All	776/820 (95%)	751 (97%)	22 (3%)	3 (0%)	34	42

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	381	MSE
1	A	274	ASP
1	A	211	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	338/342 (99%)	334 (99%)	4 (1%)	71	84
1	B	336/342 (98%)	336 (100%)	0	100	100
All	All	674/684 (98%)	670 (99%)	4 (1%)	86	94

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LEU
1	A	211	VAL
1	A	212	LEU
1	A	387	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.