



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2020 – 12:07 am BST

PDB ID : 6G3N
Title : Mycobacterial hydrolase complex 14.
Authors : Cereija, T.B.; Macedo-Ribeiro, S.; Pereira, P.J.B.
Deposited on : 2018-03-26
Resolution : 2.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	448	 2% 92% 6% •
1	B	448	 7% 89% 9% ••

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7503 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Mycobacterial hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	445	Total 3596	C 2292	N 649	O 642	S 13	0	1	0
1	B	445	Total 3596	C 2292	N 649	O 642	S 13	0	1	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP K5BDL0
A	0	ALA	-	expression tag	UNP K5BDL0
A	434	PHE	GLN	engineered mutation	UNP K5BDL0
B	-1	GLY	-	expression tag	UNP K5BDL0
B	0	ALA	-	expression tag	UNP K5BDL0
B	434	PHE	GLN	engineered mutation	UNP K5BDL0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0

- Molecule 3 is SERINE (three-letter code: SER) (formula: C₃H₇NO₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 7 3 1 3	0	0
3	B	1	Total C N O 7 3 1 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	139	Total O 139 139	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	116	Total 116	O 116	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mycobacterial hydrolase



- Molecule 1: Mycobacterial hydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	85.58Å 158.84Å 88.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.32 – 2.32 48.60 – 2.32	Depositor EDS
% Data completeness (in resolution range)	99.3 (41.32-2.32) 99.3 (48.60-2.32)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 2.32Å)	Xtrriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.153 , 0.203 0.153 , 0.204	Depositor DCC
R_{free} test set	2596 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	50.8	Xtrriage
Anisotropy	0.194	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 50.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.029 for l,-k,h	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7503	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/3707	0.61	0/5053
1	B	0.46	0/3707	0.58	0/5053
All	All	0.47	0/7414	0.60	0/10106

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3596	0	3470	18	0
1	B	3596	0	3470	27	0
2	A	24	0	32	2	0
2	B	18	0	24	2	0
3	A	7	0	4	0	0
3	B	7	0	4	0	0
4	A	139	0	0	1	0
4	B	116	0	0	0	0
All	All	7503	0	7004	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:19:LEU:HD11	1:B:433:GLN:HA	1.78	0.65
1:A:19:LEU:HD11	1:A:433:GLN:HA	1.82	0.62
1:A:383:VAL:HG22	1:A:436:TRP:CE2	2.39	0.58
1:B:94:ARG:HD3	1:B:226:LEU:HD12	1.86	0.56
1:A:383:VAL:HG22	1:A:436:TRP:CD2	2.44	0.54
1:B:186:ARG:HD2	1:B:307:ASP:OD2	2.08	0.53
1:A:89:PHE:CD1	1:A:90:PRO:HA	2.45	0.52
1:B:383:VAL:HB	1:B:436:TRP:CZ2	2.44	0.52
1:B:253:VAL:HG21	1:B:305:ALA:HB1	1.92	0.52
1:B:309:ASP:HB2	1:B:316:LEU:HD21	1.93	0.51
1:A:46:PHE:HE1	1:A:386:TRP:CE3	2.28	0.51
1:A:264:GLN:HG2	1:A:394:ARG:NH1	2.28	0.49
1:B:70:GLN:OE1	1:B:154:ARG:NH1	2.46	0.49
1:B:89:PHE:CD1	1:B:90:PRO:HA	2.48	0.49
1:B:365:PRO:HA	1:B:370:PHE:CG	2.48	0.49
1:B:383:VAL:HB	1:B:436:TRP:CE2	2.48	0.49
1:B:264:GLN:HG2	1:B:394:ARG:NH2	2.29	0.47
1:A:383:VAL:HG13	1:A:436:TRP:CZ2	2.49	0.47
1:A:36:TYR:HB3	1:A:40:TRP:CZ3	2.50	0.47
1:B:402:ARG:NH2	2:B:501:GOL:H12	2.29	0.47
1:A:42:TRP:CD2	1:A:436:TRP:HH2	2.33	0.47
1:B:405[B]:ARG:HG3	1:B:444:TRP:CD1	2.51	0.46
1:A:129:HIS:NE2	2:B:501:GOL:H2	2.31	0.46
1:A:135:ARG:HA	1:A:135:ARG:HD2	1.64	0.46
1:A:275:ARG:NH2	4:A:603:HOH:O	2.36	0.45
1:B:117:PRO:HD3	1:B:255:PHE:CE1	2.52	0.45
1:B:71:TRP:CD1	1:B:105:PRO:HD3	2.52	0.45
1:A:315:TRP:CG	2:A:502:GOL:H11	2.52	0.44
1:B:42:TRP:CD2	1:B:436:TRP:HH2	2.36	0.44
1:B:264:GLN:HG2	1:B:394:ARG:HH21	1.83	0.43
1:B:409:LEU:HA	1:B:409:LEU:HD23	1.87	0.43
1:A:254:PHE:CE2	1:A:258:ILE:HD11	2.54	0.43
1:A:60:VAL:HG13	1:A:144:PHE:CD1	2.53	0.43
1:A:266:LEU:HA	1:A:266:LEU:HD12	1.78	0.42
2:A:504:GOL:H2	1:B:129:HIS:NE2	2.34	0.42
1:B:42:TRP:O	1:B:46:PHE:HD2	2.01	0.42
1:A:72:ARG:CZ	1:A:106:ARG:HH12	2.32	0.42
1:B:101:ALA:HB3	1:B:104:ALA:HB2	2.00	0.42
1:B:145:LEU:HA	1:B:145:LEU:HD23	1.88	0.42
1:B:305:ALA:HB2	1:B:326:PRO:HD3	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:198:LYS:HB2	1:B:198:LYS:HE3	1.76	0.41
1:B:204:ARG:HD3	1:B:222:TYR:CD2	2.56	0.41
1:B:199:LEU:HD11	1:B:226:LEU:HB3	2.02	0.41
1:B:95:TRP:CH2	1:B:229:LEU:HD11	2.56	0.41
1:A:264:GLN:HG2	1:A:394:ARG:HH11	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	444/448 (99%)	428 (96%)	16 (4%)	0	100	100
1	B	444/448 (99%)	432 (97%)	12 (3%)	0	100	100
All	All	888/896 (99%)	860 (97%)	28 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	366/366 (100%)	357 (98%)	9 (2%)	47	64
1	B	366/366 (100%)	357 (98%)	9 (2%)	47	64
All	All	732/732 (100%)	714 (98%)	18 (2%)	47	64

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	TRP
1	A	86	ASP
1	A	124	GLN
1	A	182	ASP
1	A	244	SER
1	A	375	TYR
1	A	387	LEU
1	A	394	ARG
1	A	422	GLU
1	B	42	TRP
1	B	222	TYR
1	B	229	LEU
1	B	244	SER
1	B	255	PHE
1	B	368	ARG
1	B	375	TYR
1	B	394	ARG
1	B	422	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	503	-	5,5,5	1.10	0	5,5,5	0.90	0
2	GOL	B	503	-	5,5,5	0.91	0	5,5,5	0.98	0
2	GOL	A	502	-	5,5,5	0.95	0	5,5,5	0.99	0
2	GOL	A	501	-	5,5,5	0.95	0	5,5,5	1.11	0
2	GOL	B	501	-	5,5,5	0.77	0	5,5,5	1.14	0
2	GOL	B	502	-	5,5,5	0.88	0	5,5,5	0.96	0
2	GOL	A	504	-	5,5,5	0.84	0	5,5,5	1.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	503	-	-	2/4/4/4	-
2	GOL	B	503	-	-	2/4/4/4	-
2	GOL	A	502	-	-	0/4/4/4	-
2	GOL	A	501	-	-	0/4/4/4	-
2	GOL	B	501	-	-	2/4/4/4	-
2	GOL	B	502	-	-	4/4/4/4	-
2	GOL	A	504	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	503	GOL	C1-C2-C3-O3
2	B	501	GOL	C1-C2-C3-O3
2	A	504	GOL	C1-C2-C3-O3
2	A	503	GOL	O2-C2-C3-O3
2	A	503	GOL	C1-C2-C3-O3
2	B	502	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	502	GOL	C1-C2-C3-O3
2	A	504	GOL	O1-C1-C2-C3
2	B	501	GOL	O2-C2-C3-O3
2	B	502	GOL	O1-C1-C2-O2
2	B	503	GOL	O2-C2-C3-O3
2	B	502	GOL	O2-C2-C3-O3
2	A	504	GOL	O1-C1-C2-O2
2	A	504	GOL	O2-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	502	GOL	1	0
2	B	501	GOL	2	0
2	A	504	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	445/448 (99%)	0.12	9 (2%) 65 72	34, 52, 85, 135	0
1	B	445/448 (99%)	0.34	33 (7%) 14 19	39, 59, 94, 136	0
All	All	890/896 (99%)	0.23	42 (4%) 31 38	34, 56, 91, 136	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	446	GLY	8.3
1	B	202	TYR	3.9
1	B	226	LEU	3.6
1	B	405[A]	ARG	3.6
1	B	197	GLY	3.5
1	B	222	TYR	3.3
1	A	2	PRO	3.2
1	B	198	LYS	3.1
1	B	195	VAL	2.9
1	B	199	LEU	2.9
1	B	107	ASN	2.7
1	B	397	TRP	2.7
1	B	227	TRP	2.7
1	B	229	LEU	2.6
1	B	196	PRO	2.6
1	A	213	PRO	2.5
1	B	211	THR	2.5
1	A	425	THR	2.5
1	B	218	SER	2.5
1	A	405[A]	ARG	2.4
1	B	3	HIS	2.4
1	B	200	PRO	2.4
1	B	392	PHE	2.4
1	B	201	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	393	ALA	2.4
1	B	245	VAL	2.3
1	B	400	ARG	2.3
1	B	277	HIS	2.3
1	B	327	LEU	2.2
1	A	265	VAL	2.2
1	B	213	PRO	2.2
1	B	398	ALA	2.2
1	B	219	ASP	2.2
1	A	393	ALA	2.1
1	A	446	GLY	2.1
1	B	240	GLU	2.1
1	A	390	TRP	2.1
1	B	223	ASP	2.1
1	B	85	VAL	2.1
1	B	391	CYS	2.1
1	B	95	TRP	2.0
1	A	445	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	B	502	6/6	0.88	0.28	83,92,98,102	0
2	GOL	B	503	6/6	0.91	0.20	77,92,97,100	0
2	GOL	A	503	6/6	0.91	0.16	59,74,82,83	0
2	GOL	A	501	6/6	0.92	0.11	67,80,85,88	0
2	GOL	A	502	6/6	0.92	0.15	76,79,83,87	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	B	501	6/6	0.93	0.33	56,65,71,72	0
2	GOL	A	504	6/6	0.94	0.18	63,88,90,95	0
3	SER	A	505	7/7	0.96	0.11	49,56,59,61	0
3	SER	B	504	7/7	0.97	0.17	60,62,69,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.