



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 25, 2020 – 09:11 am BST

PDB ID : 6H06  
Title : FAB CBTAU-22.1 IN COMPLEX WITH TAU PEPTIDE V1088-5  
Authors : Juraszek, J.; Steinbacher, S.  
Deposited on : 2018-07-06  
Resolution : 2.63 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.









Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1426 (2.66-2.62)
Clashscore	141614	1472 (2.66-2.62)
Ramachandran outliers	138981	1446 (2.66-2.62)
Sidechain outliers	138945	1446 (2.66-2.62)
RSRZ outliers	127900	1408 (2.66-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	224	97%
1	C	224	97%
1	E	224	96%
1	H	224	95%
2	B	219	99%
2	D	219	97%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	219	 99%
2	L	219	 97%
3	G	26	 4% 31% 69%
3	I	26	 8% 27% 12% 62%
3	J	26	 27% 73%
3	K	26	 27% 73%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 13585 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HUMAN FAB ANTIBODY FRAGMENT OF CBTAU-22.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	217	Total 1628	C 1018	N 283	O 317	S 10	0	0	0
1	A	219	Total 1643	C 1027	N 286	O 320	S 10	12	0	0
1	C	219	Total 1641	C 1026	N 286	O 319	S 10	26	0	0
1	E	216	Total 1617	C 1011	N 281	O 315	S 10	16	0	0

- Molecule 2 is a protein called HUMAN FAB ANTIBODY FRAGMENT OF HCBTAU-22.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	218	Total 1686	C 1056	N 290	O 334	S 6	2	0	0
2	B	218	Total 1686	C 1056	N 290	O 334	S 6	16	0	0
2	D	217	Total 1677	C 1051	N 289	O 331	S 6	33	0	0
2	F	218	Total 1686	C 1056	N 290	O 334	S 6	23	0	0

- Molecule 3 is a protein called Microtubule-associated protein tau.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
3	I	10	Total 77	C 44	N 11	O 20	P 1	S 1	0	0	0
3	G	8	Total 61	C 35	N 9	O 16	P 1		0	0	0
3	J	7	Total 54	C 31	N 8	O 14	P 1		0	0	0
3	K	7	Total 54	C 31	N 8	O 14	P 1		0	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	L	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0
4	F	1	Total C O 6 3 3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	H	4	Total O 4 4	0	0
5	L	10	Total O 10 10	0	0
5	A	8	Total O 8 8	0	0
5	B	4	Total O 4 4	0	0
5	D	6	Total O 6 6	0	0
5	E	7	Total O 7 7	0	0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	F	6	Total	O	0	0
			6	6		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HUMAN FAB ANTIBODY FRAGMENT OF CBTAU-22.1

Chain H:  95%



- Molecule 1: HUMAN FAB ANTIBODY FRAGMENT OF CBTAU-22.1

Chain A:  97%



- Molecule 1: HUMAN FAB ANTIBODY FRAGMENT OF CBTAU-22.1

Chain C:  97%




- Molecule 1: HUMAN FAB ANTIBODY FRAGMENT OF CBTAU-22.1

Chain E:  96%




- Molecule 2: HUMAN FAB ANTIBODY FRAGMENT OF HCBTAU-22.1

Chain L:  97%



- Molecule 2: HUMAN FAB ANTIBODY FRAGMENT OF HCBTAU-22.1

Chain B:  99%



- Molecule 2: HUMAN FAB ANTIBODY FRAGMENT OF HCBTAU-22.1

Chain D: 97%




- Molecule 2: HUMAN FAB ANTIBODY FRAGMENT OF HCBTAU-22.1

Chain F: 99%




- Molecule 3: Microtubule-associated protein tau

Chain I: 8% 27% 12% 62%




- Molecule 3: Microtubule-associated protein tau

Chain G: 4% 31% 69%




- Molecule 3: Microtubule-associated protein tau

Chain J: 27% 73%



- Molecule 3: Microtubule-associated protein tau

Chain K: 27% 73%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	108.06 Å 96.21 Å 108.98 Å 90.00° 112.81° 90.00°	Depositor
Resolution (Å)	100.46 – 2.63 47.13 – 2.63	Depositor EDS
% Data completeness (in resolution range)	96.7 (100.46-2.63) 94.1 (47.13-2.63)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.34 (at 2.61 Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.216 , 0.270 0.220 , 0.273	Depositor DCC
$R_{free}$ test set	821 reflections (1.38%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.9	Xtrriage
Anisotropy	0.294	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 15.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.109 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	13585	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.57 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.6843e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PCA, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/1675	0.62	0/2280
1	C	0.36	0/1673	0.61	0/2277
1	E	0.37	0/1649	0.61	0/2247
1	H	0.36	0/1660	0.62	1/2261 (0.0%)
2	B	0.37	0/1725	0.59	0/2343
2	D	0.37	0/1716	0.60	0/2331
2	F	0.37	0/1725	0.60	0/2343
2	L	0.36	0/1725	0.61	0/2343
3	G	0.49	0/50	0.65	0/66
3	I	0.53	0/66	0.80	0/87
3	J	0.44	0/43	0.59	0/56
3	K	0.53	0/43	0.63	0/56
All	All	0.37	0/13750	0.61	1/18690 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	67	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1643	0	1604	0	0
1	C	1641	0	1602	0	0
1	E	1617	0	1571	0	0
1	H	1628	0	1586	1	0
2	B	1686	0	1647	1	0
2	D	1677	0	1641	2	0
2	F	1686	0	1647	1	0
2	L	1686	0	1647	2	0
3	G	61	0	54	0	0
3	I	77	0	67	1	0
3	J	54	0	47	0	0
3	K	54	0	47	0	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
4	D	6	0	8	0	0
4	F	6	0	8	0	0
4	L	6	0	8	0	0
5	A	8	0	0	0	0
5	B	4	0	0	0	0
5	D	6	0	0	0	0
5	E	7	0	0	0	0
5	F	6	0	0	0	0
5	H	4	0	0	0	0
5	L	10	0	0	0	0
All	All	13585	0	13200	8	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

The worst 5 of 8 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:41:TYR:CD1	2:F:49:PRO:HB3	2.51	0.45
2:B:41:TYR:CD1	2:B:49:PRO:HB3	2.53	0.44
2:D:38:LEU:HD13	2:D:76:PHE:CD1	2.53	0.44
2:L:141:LEU:HD21	2:L:201:VAL:HG13	2.00	0.44
3:I:422:SEP:N	3:I:423:PRO:HD2	2.33	0.44

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	215/224 (96%)	209 (97%)	6 (3%)	0	100	100
1	C	215/224 (96%)	208 (97%)	7 (3%)	0	100	100
1	E	212/224 (95%)	204 (96%)	8 (4%)	0	100	100
1	H	213/224 (95%)	209 (98%)	4 (2%)	0	100	100
2	B	216/219 (99%)	210 (97%)	6 (3%)	0	100	100
2	D	215/219 (98%)	205 (95%)	10 (5%)	0	100	100
2	F	216/219 (99%)	209 (97%)	7 (3%)	0	100	100
2	L	216/219 (99%)	213 (99%)	3 (1%)	0	100	100
3	G	5/26 (19%)	5 (100%)	0	0	100	100
3	I	7/26 (27%)	6 (86%)	1 (14%)	0	100	100
3	J	4/26 (15%)	4 (100%)	0	0	100	100
3	K	4/26 (15%)	3 (75%)	1 (25%)	0	100	100
All	All	1738/1876 (93%)	1685 (97%)	53 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/188 (98%)	182 (99%)	2 (1%)	73	85
1	C	183/188 (97%)	182 (100%)	1 (0%)	88	94

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	180/188 (96%)	178 (99%)	2 (1%)	73	85
1	H	182/188 (97%)	181 (100%)	1 (0%)	88	94
2	B	193/194 (100%)	192 (100%)	1 (0%)	88	94
2	D	192/194 (99%)	192 (100%)	0	100	100
2	F	193/194 (100%)	193 (100%)	0	100	100
2	L	193/194 (100%)	192 (100%)	1 (0%)	88	94
3	G	6/21 (29%)	6 (100%)	0	100	100
3	I	8/21 (38%)	7 (88%)	1 (12%)	4	5
3	J	5/21 (24%)	5 (100%)	0	100	100
3	K	5/21 (24%)	5 (100%)	0	100	100
All	All	1524/1612 (94%)	1515 (99%)	9 (1%)	86	93

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	49	PRO
3	I	419	MET
1	E	72	ARG
1	A	72	ARG
1	C	72	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
2	L	142	ASN
2	D	143	ASN
1	C	172	HIS
2	L	27	GLN
2	D	142	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	PCA	E	1	1	7,8,9	0.45	0	9,10,12	1.19	1 (11%)
1	PCA	C	1	1	7,8,9	0.49	0	9,10,12	1.38	1 (11%)
1	PCA	H	1	1	7,8,9	0.55	0	9,10,12	1.42	1 (11%)
3	SEP	I	422	3	8,9,10	0.70	0	8,12,14	1.87	2 (25%)
3	SEP	J	422	3	8,9,10	0.62	0	8,12,14	1.21	1 (12%)
3	SEP	K	422	3	8,9,10	0.67	0	8,12,14	1.42	1 (12%)
3	SEP	G	422	3	8,9,10	0.65	0	8,12,14	1.47	1 (12%)
1	PCA	A	1	1	7,8,9	0.42	0	9,10,12	1.68	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	E	1	1	-	0/0/11/13	0/1/1/1
1	PCA	C	1	1	-	0/0/11/13	0/1/1/1
1	PCA	H	1	1	-	0/0/11/13	0/1/1/1
3	SEP	I	422	3	-	0/5/8/10	-
3	SEP	J	422	3	-	1/5/8/10	-
3	SEP	K	422	3	-	0/5/8/10	-
3	SEP	G	422	3	-	0/5/8/10	-
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	422	SEP	OG-CB-CA	4.05	112.09	108.14

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	PCA	CB-CA-C	-4.00	107.20	112.70
1	H	1	PCA	CB-CA-C	-3.13	108.39	112.70
1	C	1	PCA	CB-CA-C	-2.99	108.59	112.70
3	K	422	SEP	OG-P-O1P	-2.93	98.27	106.47

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	J	422	SEP	CA-CB-OG-P

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	422	SEP	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	F	301	-	5,5,5	0.30	0	5,5,5	0.31	0
4	GOL	B	301	-	5,5,5	0.29	0	5,5,5	0.28	0
4	GOL	D	301	-	5,5,5	0.35	0	5,5,5	0.35	0
4	GOL	A	301	-	5,5,5	0.33	0	5,5,5	0.31	0
4	GOL	L	301	-	5,5,5	0.38	0	5,5,5	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	F	301	-	-	0/4/4/4	-
4	GOL	B	301	-	-	0/4/4/4	-
4	GOL	D	301	-	-	2/4/4/4	-
4	GOL	A	301	-	-	1/4/4/4	-
4	GOL	L	301	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	301	GOL	C1-C2-C3-O3
4	L	301	GOL	O1-C1-C2-C3
4	L	301	GOL	C1-C2-C3-O3
4	D	301	GOL	O2-C2-C3-O3
4	L	301	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	218/224 (97%)	-0.25	1 (0%) 91   90	34, 49, 81, 104	4 (1%)
1	C	218/224 (97%)	-0.26	0 100   100	34, 54, 81, 114	8 (3%)
1	E	215/224 (95%)	-0.28	0 100   100	31, 50, 83, 95	5 (2%)
1	H	216/224 (96%)	-0.33	0 100   100	32, 48, 70, 94	0
2	B	218/219 (99%)	-0.31	1 (0%) 91   90	35, 56, 80, 111	6 (2%)
2	D	217/219 (99%)	-0.17	0 100   100	30, 63, 91, 102	11 (5%)
2	F	218/219 (99%)	-0.26	0 100   100	33, 55, 88, 108	6 (2%)
2	L	218/219 (99%)	-0.32	0 100   100	29, 50, 76, 93	1 (0%)
3	G	7/26 (26%)	0.73	1 (14%) 2   1	66, 74, 104, 123	0
3	I	9/26 (34%)	0.95	2 (22%) 0   0	44, 73, 112, 132	0
3	J	6/26 (23%)	0.18	0 100   100	61, 68, 86, 89	0
3	K	6/26 (23%)	-0.13	0 100   100	69, 86, 91, 95	0
All	All	1766/1876 (94%)	-0.26	5 (0%) 94   95	29, 53, 85, 132	41 (2%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	I	419	MET	4.3
3	G	427	THR	3.5
3	I	427	THR	3.2
2	B	218	GLU	2.4
1	A	199	THR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	PCA	A	1	8/9	0.91	0.20	61,68,76,78	0
3	SEP	G	422	10/11	0.94	0.18	45,62,64,73	0
1	PCA	E	1	8/9	0.94	0.12	40,42,53,54	0
3	SEP	J	422	10/11	0.95	0.19	49,60,71,76	0
1	PCA	C	1	8/9	0.96	0.14	51,60,66,75	0
3	SEP	K	422	10/11	0.97	0.16	46,65,69,69	0
3	SEP	I	422	10/11	0.97	0.16	41,49,55,59	0
1	PCA	H	1	8/9	0.97	0.13	42,51,52,65	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	A	301	6/6	0.87	0.40	74,76,78,81	0
4	GOL	B	301	6/6	0.90	0.30	79,84,95,102	0
4	GOL	F	301	6/6	0.91	0.20	67,84,85,88	0
4	GOL	D	301	6/6	0.91	0.36	60,77,84,87	0
4	GOL	L	301	6/6	0.94	0.18	60,69,77,92	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.