



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 02:45 am BST

PDB ID : 3HJR  
Title : Crystal structure of serine protease of *Aeromonas sobria*  
Authors : Utsunomiya, H.; Tsuge, H.; Kobayashi, H.; Okamoto, K.  
Deposited on : 2009-05-22  
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	600	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5161 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Extracellular serine protease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	593	4458	2747	808	887	16	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	70	GLN	HIS	SEE REMARK 999	UNP Q9L5A4
A	114	ALA	GLY	SEE REMARK 999	UNP Q9L5A4
A	176	GLN	HIS	SEE REMARK 999	UNP Q9L5A4
A	179	GLY	ARG	SEE REMARK 999	UNP Q9L5A4

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Ca	0	0
			3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	700	Total	O	0	0
			700	700		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Extracellular serine protease



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.82Å 112.06Å 51.88Å 90.00° 110.75° 90.00°	Depositor
Resolution (Å)	43.02 – 1.65 43.02 – 1.65	Depositor EDS
% Data completeness (in resolution range)	98.7 (43.02-1.65) 98.7 (43.02-1.65)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.41 (at 1.65Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.168 , 0.202 0.178 , 0.209	Depositor DCC
$R_{free}$ test set	3172 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.1	Xtrriage
Anisotropy	0.562	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 47.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.024 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5161	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.63	1/4541 (0.0%)	0.95	22/6180 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	267	ARG	CD-NE	-5.14	1.37	1.46

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	267	ARG	NE-CZ-NH2	-16.44	112.08	120.30
1	A	267	ARG	NE-CZ-NH1	14.96	127.78	120.30
1	A	256	LEU	CA-CB-CG	7.70	133.00	115.30
1	A	15	ASP	CB-CG-OD2	6.49	124.14	118.30
1	A	373	ASP	CB-CG-OD2	6.45	124.11	118.30
1	A	44	ASP	CB-CG-OD2	6.44	124.10	118.30
1	A	355	ASP	CB-CG-OD2	6.18	123.86	118.30
1	A	256	LEU	CB-CG-CD1	6.08	121.33	111.00
1	A	148	ASP	CB-CG-OD2	5.84	123.56	118.30
1	A	197	ASP	CB-CG-OD2	5.82	123.54	118.30
1	A	327	ASP	CB-CG-OD2	5.82	123.53	118.30
1	A	110	ASP	CB-CG-OD2	5.52	123.27	118.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	310	ASP	CB-CG-OD2	5.51	123.26	118.30
1	A	454	ASP	CB-CG-OD1	5.50	123.25	118.30
1	A	323	ASP	CB-CG-OD2	5.44	123.19	118.30
1	A	33	ASP	CB-CG-OD2	5.39	123.15	118.30
1	A	87	ASP	CB-CG-OD2	5.38	123.14	118.30
1	A	367	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	A	104	ASP	CB-CG-OD2	5.30	123.07	118.30
1	A	164	ASP	CB-CG-OD2	5.28	123.05	118.30
1	A	571	ARG	NE-CZ-NH1	-5.23	117.69	120.30
1	A	267	ARG	CD-NE-CZ	5.06	130.68	123.60

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	208	GLN	Peptide
1	A	318	GLY	Peptide
1	A	355	ASP	Peptide
1	A	9	GLY	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4458	0	4344	76	2
2	A	3	0	0	0	0
3	A	700	0	0	30	1
All	All	5161	0	4344	76	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:THR:HB	3:A:1171:HOH:O	1.04	1.20

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:454:ASP:O	1:A:553:ASP:OD2	1.60	1.17
1:A:3:GLY:HA2	3:A:1258:HOH:O	1.66	0.95
1:A:454:ASP:O	1:A:553:ASP:CG	2.04	0.94
1:A:208:GLN:O	1:A:208:GLN:HG2	1.69	0.90
1:A:267:ARG:HD2	1:A:268:SER:O	1.72	0.89
1:A:569:ARG:N	3:A:1172:HOH:O	2.05	0.87
1:A:302:ASP:HB2	3:A:1070:HOH:O	1.75	0.85
1:A:276:ASN:H	1:A:276:ASN:HD22	1.27	0.83
1:A:70:GLN:H	1:A:70:GLN:HE21	1.26	0.82
1:A:99:VAL:H	1:A:145:ASN:HD21	1.26	0.82
1:A:317:HIS:HE1	1:A:330:GLY:H	1.31	0.78
1:A:580:ARG:HB3	3:A:1053:HOH:O	1.84	0.76
1:A:68:LEU:H	1:A:70:GLN:HE22	1.36	0.73
1:A:208:GLN:O	1:A:208:GLN:CG	2.37	0.72
1:A:243:ASN:HD22	1:A:245:ASN:H	1.37	0.72
1:A:355:ASP:HA	3:A:1176:HOH:O	1.91	0.70
1:A:70:GLN:H	1:A:70:GLN:NE2	1.90	0.69
1:A:565:ASN:HB3	1:A:568:THR:O	1.94	0.68
1:A:454:ASP:HA	3:A:1235:HOH:O	1.95	0.66
1:A:3:GLY:N	3:A:1221:HOH:O	2.29	0.65
1:A:238:LYS:HE3	3:A:664:HOH:O	1.95	0.65
1:A:252:ASN:HD21	1:A:255:ASN:H	1.47	0.63
1:A:99:VAL:N	1:A:145:ASN:HD21	1.98	0.62
1:A:99:VAL:H	1:A:145:ASN:ND2	1.96	0.61
1:A:63:HIS:HE1	1:A:135:ILE:O	1.83	0.61
1:A:568:THR:C	3:A:1172:HOH:O	2.35	0.61
1:A:166:ASN:HB3	3:A:1087:HOH:O	1.99	0.61
1:A:270:TYR:CE1	1:A:334:GLY:HA2	2.36	0.60
1:A:12:SER:HB2	3:A:954:HOH:O	2.01	0.59
1:A:40:ASN:HD22	1:A:42:GLY:H	1.49	0.59
1:A:256:LEU:HD13	1:A:278:PHE:HB3	1.84	0.58
1:A:384:THR:CB	3:A:1171:HOH:O	1.89	0.57
1:A:568:THR:CA	3:A:1172:HOH:O	2.51	0.57
1:A:221:ASN:HD21	1:A:238:LYS:NZ	2.01	0.57
1:A:354:PRO:O	1:A:355:ASP:HB2	2.05	0.57
1:A:145:ASN:H	1:A:145:ASN:HD22	1.54	0.56
1:A:267:ARG:CD	1:A:268:SER:O	2.52	0.56
1:A:41:SER:OG	1:A:43:GLN:HG3	2.07	0.55
1:A:376:HIS:HE1	3:A:629:HOH:O	1.89	0.55
1:A:454:ASP:CA	3:A:1235:HOH:O	2.52	0.55
1:A:570:GLU:HG3	3:A:1238:HOH:O	2.07	0.54

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:580:ARG:HA	3:A:1037:HOH:O	2.07	0.54
1:A:371:ARG:NE	3:A:1300:HOH:O	2.35	0.53
1:A:406:MET:HG2	3:A:1083:HOH:O	2.07	0.53
1:A:373:ASP:O	1:A:376:HIS:HD2	1.91	0.53
1:A:360:ASP:OD2	1:A:429:HIS:HE1	1.93	0.52
1:A:243:ASN:ND2	1:A:245:ASN:H	2.08	0.51
1:A:518:GLN:NE2	3:A:715:HOH:O	2.44	0.51
1:A:40:ASN:ND2	1:A:42:GLY:H	2.08	0.51
1:A:523:VAL:HG12	3:A:986:HOH:O	2.11	0.50
1:A:565:ASN:O	1:A:568:THR:O	2.30	0.49
1:A:384:THR:HG23	3:A:1143:HOH:O	2.12	0.49
1:A:384:THR:CG2	3:A:1171:HOH:O	2.45	0.49
1:A:454:ASP:CB	3:A:1235:HOH:O	2.60	0.49
1:A:568:THR:OG1	1:A:568:THR:O	2.31	0.48
1:A:581:GLN:NE2	3:A:1261:HOH:O	2.45	0.48
1:A:104:ASP:HB3	3:A:1103:HOH:O	2.13	0.47
1:A:145:ASN:N	1:A:145:ASN:HD22	2.13	0.46
1:A:70:GLN:HG2	1:A:138:ARG:NH1	2.30	0.45
1:A:252:ASN:ND2	1:A:255:ASN:H	2.13	0.45
1:A:571:ARG:NE	3:A:1011:HOH:O	2.27	0.45
1:A:40:ASN:C	1:A:40:ASN:HD22	2.19	0.45
1:A:568:THR:C	1:A:570:GLU:H	2.21	0.44
1:A:454:ASP:O	1:A:553:ASP:CB	2.66	0.43
1:A:245:ASN:HD21	1:A:534:LEU:H	1.67	0.42
1:A:221:ASN:HD21	1:A:238:LYS:HZ2	1.67	0.42
1:A:276:ASN:N	1:A:276:ASN:HD22	2.02	0.41
1:A:302:ASP:OD2	3:A:1217:HOH:O	2.22	0.41
1:A:442:LYS:HE3	1:A:444:ASN:OD1	2.19	0.41
1:A:485:ARG:HD2	1:A:576:GLU:HB3	2.02	0.41
1:A:23:ARG:HG3	1:A:60:TRP:CD1	2.55	0.41
1:A:135:ILE:CD1	1:A:341:SER:HA	2.51	0.41
1:A:276:ASN:H	1:A:276:ASN:ND2	2.07	0.40
1:A:371:ARG:NH2	3:A:711:HOH:O	2.54	0.40
1:A:63:HIS:CD2	1:A:133:LYS:NZ	2.89	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:166:ASN:ND2	3:A:1204:HOH:O[1_554]	1.87	0.33
1:A:12:SER:OG	1:A:446:THR:OG1[2_545]	1.89	0.31

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	591/600 (98%)	572 (97%)	18 (3%)	1 (0%)	47 28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	569	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	484/490 (99%)	466 (96%)	18 (4%)	34 10

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	LEU
1	A	40	ASN
1	A	43	GLN
1	A	52	ILE
1	A	59	LEU
1	A	63	HIS
1	A	70	GLN
1	A	145	ASN
1	A	243	ASN
1	A	252	ASN

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	256	LEU
1	A	261	LEU
1	A	276	ASN
1	A	455	VAL
1	A	472	VAL
1	A	556	ASN
1	A	580	ARG
1	A	581	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	40	ASN
1	A	63	HIS
1	A	70	GLN
1	A	73	ASN
1	A	91	ASN
1	A	145	ASN
1	A	150	ASN
1	A	152	GLN
1	A	155	GLN
1	A	166	ASN
1	A	175	ASN
1	A	176	GLN
1	A	193	GLN
1	A	221	ASN
1	A	231	ASN
1	A	243	ASN
1	A	245	ASN
1	A	252	ASN
1	A	276	ASN
1	A	314	ASN
1	A	317	HIS
1	A	376	HIS
1	A	429	HIS
1	A	484	GLN
1	A	518	GLN
1	A	556	ASN
1	A	560	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	593/600 (98%)	0.14	24 (4%) 38 38	3, 9, 16, 24	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	580	ARG	5.8
1	A	9	GLY	4.8
1	A	318	GLY	4.0
1	A	12	SER	3.8
1	A	13	GLY	3.5
1	A	577	ARG	3.4
1	A	208	GLN	3.2
1	A	448	SER	3.0
1	A	302	ASP	3.0
1	A	568	THR	3.0
1	A	3	GLY	3.0
1	A	10	LYS	2.8
1	A	355	ASP	2.6
1	A	581	GLN	2.5
1	A	387	THR	2.5
1	A	235	ASN	2.4
1	A	454	ASP	2.4
1	A	309	ASP	2.4
1	A	166	ASN	2.3
1	A	521	GLY	2.3
1	A	386	SER	2.2
1	A	311	PRO	2.2
1	A	519	GLN	2.2
1	A	569	ARG	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CA	A	603	1/1	0.98	0.11	27,27,27,27	0
2	CA	A	601	1/1	0.98	0.05	22,22,22,22	0
2	CA	A	602	1/1	0.99	0.16	33,33,33,33	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.