



Full wwPDB NMR Structure Validation Report ⓘ

Feb 13, 2022 – 09:50 PM EST

PDB ID : 1HO6
Title : CHIMERIC ARABINONUCLEIC ACID (ANA) HAIRPIN WITH ANA/RNA
HYBRID STEM
Authors : Denisov, A.Y.; Gehring, K.
Deposited on : 2000-12-08

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.26
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.26


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

SOLUTION NMR


The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	12	

2 Ensemble composition and analysis

This entry contains 10 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 384 atoms, of which 135 are hydrogens and 0 are deuteriums.

- Molecule 1 is DNA/RNA hybrid called DNA/RNA (5'-R(*GP*GP*AP*C)-D(P*TP*TP*C P*G)-(P*(GAO)P*(UAR)P*(CAR)P*(CAR)-3').

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
1	A	12	384	115	135	43	80	11	0

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA/RNA (5'-R(*GP*GP*AP*C)-D(P*TP*TP*CP*G)-(P*(GAO)P*(UAR)P*(CAR)P*(CAR)-3')




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: DNA/RNA (5'-R(*GP*GP*AP*C)-D(P*TP*TP*CP*G)-(P*(GAO)P*(UAR)P*(CAR)P*(CAR)-3')



4.2.2 Score per residue for model 2

- Molecule 1: DNA/RNA (5'-R(*GP*GP*AP*C)-D(P*TP*TP*CP*G)-(P*(GAO)P*(UAR)P*(CAR)P*(CAR)-3')



4.2.3 Score per residue for model 3

- Molecule 1: DNA/RNA (5'-R(*GP*GP*AP*C)-D(P*TP*TP*CP*G)-(P*(GAO)P*(UAR)P*(CAR)P*(CAR)-3')

Chain A:  33% 67%



4.2.4 Score per residue for model 4


- Molecule 1: DNA/RNA (5'-R(*GP*GP*AP*C)-D(P*TP*TP*CP*G)-(P*(GAO)P*(UAR)P*(CAR)P*(CAR)-3')

Chain A:  17% 83%



4.2.5 Score per residue for model 5

- Molecule 1: DNA/RNA (5'-R(*GP*GP*AP*C)-D(P*TP*TP*CP*G)-(P*(GAO)P*(UAR)P*(CAR)P*(CAR)-3')

Chain A:  50% 50%



4.2.6 Score per residue for model 6

- Molecule 1: DNA/RNA (5'-R(*GP*GP*AP*C)-D(P*TP*TP*CP*G)-(P*(GAO)P*(UAR)P*(CAR)P*(CAR)-3')

Chain A:  42% 58%



4.2.7 Score per residue for model 7

- Molecule 1: DNA/RNA (5'-R(*GP*GP*AP*C)-D(P*TP*TP*CP*G)-(P*(GAO)P*(UAR)P*(CAR)P*(CAR)-3')


Chain A:  8% 8% 83%



4.2.8 Score per residue for model 8

- Molecule 1: DNA/RNA (5'-R(*GP*GP*AP*C)-D(P*TP*TP*CP*G)-(P*(GAO)P*(UAR)P*(CAR)P*(CAR)-3')


Chain A:  58% 42%



4.2.9 Score per residue for model 9


- Molecule 1: DNA/RNA (5'-R(*GP*GP*AP*C)-D(P*TP*TP*CP*G)-(P*(GAO)P*(UAR)P*(CAR)P*(CAR)-3')


Chain A:  50% 50%



4.2.10 Score per residue for model 10

- Molecule 1: DNA/RNA (5'-R(*GP*GP*AP*C)-D(P*TP*TP*CP*G)-(P*(GAO)P*(UAR)P*(CAR)P*(CAR)-3')

Chain A:  8% 58% 33%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 10 calculated structures, 10 were deposited, based on the following criterion: *all calculated structures submitted*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.843

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: UAR, GAO, CAR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.48±0.02	2±0/185 (0.9± 0.2%)	2.20±0.01	14±1/286 (5.0± 0.4%)
All	All	1.48	17/1850 (0.9%)	2.20	144/2860 (5.0%)

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	5	DT	C5-C7	5.77	1.53	1.50	10	10
1	A	6	DT	C5-C7	5.73	1.53	1.50	1	7

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	2	G	N7-C8-N9	9.98	118.09	113.10	2	10
1	A	8	DG	N7-C8-N9	9.47	117.83	113.10	9	10
1	A	1	G	N7-C8-N9	9.46	117.83	113.10	3	10
1	A	3	A	N7-C8-N9	7.62	117.61	113.80	9	10
1	A	8	DG	C8-N9-C4	-6.93	103.63	106.40	1	10
1	A	1	G	C8-N9-C4	-6.56	103.78	106.40	3	10
1	A	6	DT	C6-C5-C7	-6.41	119.06	122.90	4	10
1	A	2	G	C8-N9-C4	-6.35	103.86	106.40	5	10
1	A	2	G	O4'-C1'-N9	6.00	113.00	108.20	8	4
1	A	4	C	O4'-C1'-N1	5.98	112.99	108.20	10	1
1	A	8	DG	O4'-C1'-N9	5.96	112.17	108.00	1	1
1	A	5	DT	O4'-C1'-N1	5.79	112.06	108.00	4	4
1	A	5	DT	C6-C5-C7	-5.79	119.43	122.90	1	10
1	A	1	G	C5-N7-C8	-5.67	101.47	104.30	4	10
1	A	2	G	C5-N7-C8	-5.61	101.50	104.30	6	10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	6	DT	O4'-C1'-N1	5.41	111.79	108.00	5	2
1	A	3	A	O4'-C1'-N9	5.41	112.53	108.20	4	2
1	A	7	DC	O4'-C1'-N1	5.35	111.75	108.00	3	2
1	A	2	G	C3'-C2'-C1'	5.35	105.78	101.50	9	3
1	A	8	DG	C5-N7-C8	-5.31	101.65	104.30	2	9
1	A	1	G	O4'-C1'-N9	5.19	112.36	108.20	6	1
1	A	3	A	C5-N7-C8	-5.17	101.31	103.90	4	5

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	249	135	135	9±3
All	All	2490	1350	1350	88

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:9:GAO:OP2	1:A:9:GAO:H8	0.98	1.58	2	2
1:A:5:DT:H2'	1:A:6:DT:O4'	0.89	1.68	2	5
1:A:8:DG:N3	1:A:8:DG:H2'	0.78	1.91	2	4
1:A:8:DG:H4'	1:A:9:GAO:O5'	0.72	1.85	5	1
1:A:6:DT:H4'	1:A:7:DC:OP2	0.69	1.86	4	1
1:A:10:UAR:O2'	1:A:11:CAR:H6	0.68	1.89	5	8
1:A:4:C:C5	1:A:5:DT:C4	0.65	2.84	2	4
1:A:6:DT:H4'	1:A:8:DG:O6	0.57	1.99	2	1
1:A:5:DT:H2'	1:A:7:DC:N3	0.57	2.15	8	1
1:A:6:DT:H4'	1:A:7:DC:OP1	0.56	2.00	6	1
1:A:3:A:H2'	1:A:4:C:C6	0.55	2.37	1	5
1:A:2:G:O2'	1:A:3:A:H5'	0.55	2.01	2	2

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:4:C:C4	1:A:5:DT:C4	0.55	2.94	7	2
1:A:4:C:O2'	1:A:5:DT:H5'	0.54	2.02	4	3
1:A:3:A:O2'	1:A:4:C:H5'	0.54	2.03	8	4
1:A:2:G:C6	1:A:3:A:C5	0.54	2.96	2	2
1:A:7:DC:C6	1:A:8:DG:C5	0.51	2.98	8	1
1:A:3:A:H2'	1:A:4:C:O4'	0.51	2.05	10	1
1:A:5:DT:H2'	1:A:6:DT:C6	0.51	2.41	3	3
1:A:9:GAO:OP2	1:A:9:GAO:C8	0.51	2.47	2	1
1:A:11:CAR:O2'	1:A:12:CAR:H6	0.51	2.06	7	2
1:A:4:C:H2'	1:A:5:DT:O4'	0.51	2.05	4	1
1:A:4:C:H2'	1:A:5:DT:C6	0.47	2.45	1	4
1:A:5:DT:C2'	1:A:6:DT:O4'	0.46	2.64	10	1
1:A:8:DG:N3	1:A:8:DG:C2'	0.46	2.79	9	2
1:A:7:DC:H4'	1:A:8:DG:O4'	0.46	2.10	5	1
1:A:3:A:C2	1:A:4:C:C2	0.46	3.04	1	1
1:A:8:DG:H2''	1:A:9:GAO:OP2	0.46	2.10	7	1
1:A:3:A:C5	1:A:4:C:C4	0.46	3.04	1	1
1:A:6:DT:OP2	1:A:6:DT:H73	0.46	2.11	4	3
1:A:3:A:C6	1:A:4:C:C4	0.45	3.04	2	2
1:A:11:CAR:C4	1:A:12:CAR:C4	0.43	2.96	4	2
1:A:10:UAR:O2'	1:A:11:CAR:C6	0.43	2.64	5	2
1:A:1:G:O2'	1:A:2:G:H5'	0.43	2.14	3	2
1:A:1:G:C2	1:A:2:G:C5	0.43	3.07	7	2
1:A:3:A:C6	1:A:4:C:N3	0.43	2.87	1	1
1:A:8:DG:N3	1:A:9:GAO:N7	0.43	2.67	4	1
1:A:1:G:C6	1:A:2:G:C5	0.43	3.06	3	1
1:A:1:G:HO2'	1:A:2:G:C5'	0.43	2.27	2	1
1:A:9:GAO:OP2	1:A:9:GAO:H3'	0.42	2.14	5	1
1:A:9:GAO:H2'	1:A:10:UAR:O5'	0.42	2.14	4	1
1:A:4:C:C4	1:A:5:DT:N3	0.41	2.88	5	1
1:A:2:G:HO2'	1:A:3:A:C5'	0.41	2.28	3	1
1:A:1:G:H2'	1:A:2:G:O4'	0.41	2.16	3	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
1	CAR	A	12	1	15,21,22	0.86±0.01	0±0 (1±2%)
1	GAO	A	9	1	18,25,26	1.29±0.02	2±0 (12±2%)
1	CAR	A	11	1	15,21,22	0.86±0.01	0±0 (0±0%)
1	UAR	A	10	1	14,21,22	1.11±0.02	1±0 (7±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
1	CAR	A	12	1	17,30,33	1.46±0.01	2±0 (11±0%)
1	GAO	A	9	1	20,37,40	2.71±0.02	5±0 (25±0%)
1	CAR	A	11	1	17,30,33	1.50±0.01	2±0 (11±0%)
1	UAR	A	10	1	14,30,33	1.47±0.01	2±0 (14±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	UAR	A	10	1	-	0±0,5,25,26	0±0,2,2,2
1	CAR	A	12	1	-	0±0,5,25,26	0±0,2,2,2
1	CAR	A	11	1	-	0±0,5,25,26	0±0,2,2,2
1	GAO	A	9	1	-	0±0,3,25,26	0±0,3,3,3

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	9	GAO	C6-N1	3.36	1.38	1.33	5	10
1	A	9	GAO	C8-N7	3.23	1.28	1.34	2	10
1	A	10	UAR	C4-N3	3.09	1.38	1.33	7	10
1	A	9	GAO	O4'-C1'	2.30	1.44	1.41	2	2
1	A	12	CAR	O4'-C1'	2.04	1.43	1.41	9	2

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	9	GAO	C5-C6-N1	8.96	111.18	123.43	6	10
1	A	9	GAO	C2-N1-C6	6.10	125.62	115.93	6	10
1	A	11	CAR	C4-N3-C2	4.83	121.24	116.34	8	10
1	A	12	CAR	C4-N3-C2	4.58	120.99	116.34	2	10
1	A	10	UAR	C5-C4-N3	4.21	114.05	123.31	6	10
1	A	9	GAO	C3'-C2'-C1'	3.15	105.72	100.98	4	10
1	A	9	GAO	C2-N3-C4	2.99	111.94	115.36	2	10
1	A	10	UAR	C3'-C2'-C1'	2.98	105.47	100.98	5	10
1	A	11	CAR	C3'-C2'-C1'	2.84	105.26	100.98	10	10
1	A	12	CAR	C3'-C2'-C1'	2.81	105.22	100.98	2	10
1	A	9	GAO	N3-C2-N1	2.52	123.86	127.22	7	10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided