



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 18, 2024 – 08:51 PM EST

PDB ID : 4HVC
Title : Crystal structure of human prolyl-tRNA synthetase in complex with halofuginone and ATP analogue
Authors : Zhou, H.; Sun, L.; Yang, X.L.; Schimmel, P.
Deposited on : 2012-11-06
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	519	 9% 88% 5% 7%
1	B	519	 8% 85% 8% 7%

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 7989 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bifunctional glutamate/proline--tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	482	Total	C	N	O	S	0	5	0
			3844	2462	648	710	24			
1	B	483	Total	C	N	O	S	0	5	0
			3827	2454	644	705	24			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	994	MET	-	expression tag	UNP P07814
A	995	HIS	-	expression tag	UNP P07814
A	996	HIS	-	expression tag	UNP P07814
A	997	HIS	-	expression tag	UNP P07814
A	998	HIS	-	expression tag	UNP P07814
A	999	HIS	-	expression tag	UNP P07814
A	1000	HIS	-	expression tag	UNP P07814
A	1001	GLY	-	expression tag	UNP P07814
A	1002	SER	-	expression tag	UNP P07814
B	994	MET	-	expression tag	UNP P07814
B	995	HIS	-	expression tag	UNP P07814
B	996	HIS	-	expression tag	UNP P07814
B	997	HIS	-	expression tag	UNP P07814
B	998	HIS	-	expression tag	UNP P07814
B	999	HIS	-	expression tag	UNP P07814
B	1000	HIS	-	expression tag	UNP P07814
B	1001	GLY	-	expression tag	UNP P07814
B	1002	SER	-	expression tag	UNP P07814

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	31	10	6	12	3	0	0
2	B	1	31	10	6	12	3	0	0

- Molecule 3 is 7-bromo-6-chloro-3-{3-[(2R,3S)-3-hydroxypiperidin-2-yl]-2-oxopropyl}quinazol in-4(3H)-one (three-letter code: HFG) (formula: C₁₆H₁₇BrClN₃O₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	Br	C	Cl	N			O
3	A	1	24	1	16	1	3	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	Br	C	Cl	N			O
3	B	1	24	1	16	1	3	3	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Zn	0	0
			1	1		
5	B	1	Total	Zn	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	120	Total	O	0	0
			120	120		
6	B	84	Total	O	0	0
			84	84		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional glutamate/proline--tRNA ligase



- Molecule 1: Bifunctional glutamate/proline--tRNA ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	72.54Å 93.10Å 87.00Å 90.00° 107.95° 90.00°	Depositor
Resolution (Å)	46.43 – 2.00 46.43 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.5 (46.43-2.00) 98.5 (46.43-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.53 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.204 , 0.227 0.204 , 0.229	Depositor DCC
R_{free} test set	3678 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	40.2	Xtrriage
Anisotropy	0.288	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7989	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HFG, ANP, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/3950	0.52	0/5355
1	B	0.39	0/3934	0.51	0/5337
All	All	0.39	0/7884	0.51	0/10692

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3844	0	3748	17	0
1	B	3827	0	3730	20	0
2	A	31	0	13	1	0
2	B	31	0	13	1	0
3	A	24	0	17	1	0
3	B	24	0	17	2	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	120	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	84	0	0	0	0
All	All	7989	0	7538	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1415[B]:ARG:CG	1:A:1415[B]:ARG:HH11	1.70	1.04
1:A:1415[B]:ARG:HH11	1:A:1415[B]:ARG:HG2	1.32	0.92
1:A:1415[B]:ARG:HH11	1:A:1415[B]:ARG:HG3	1.52	0.74
1:A:1415[B]:ARG:HG2	1:A:1415[B]:ARG:NH1	2.05	0.71
1:B:1030[B]:GLN:NE2	1:B:1030[B]:GLN:HA	2.12	0.64
1:A:1415[B]:ARG:CG	1:A:1415[B]:ARG:NH1	2.42	0.63
1:B:1442:ILE:HG22	1:B:1509:GLY:HA3	1.91	0.52
1:B:1101:VAL:HG22	3:B:1602:HFG:BR1	2.65	0.52
1:B:1479:LEU:HD12	1:B:1507:LEU:HG	1.92	0.52
1:A:1136:SER:OG	1:A:1138:ARG:HG2	2.09	0.52
1:A:1326:LYS:HG2	1:A:1374:MET:HG2	1.91	0.52
1:B:1090[B]:GLU:HB3	1:B:1251:MET:HE3	1.96	0.48
1:B:1092:THR:HG22	1:B:1248:PHE:HE1	1.79	0.47
1:A:1415[B]:ARG:HG3	1:A:1415[B]:ARG:NH1	2.21	0.47
1:B:1170:GLN:HE21	1:B:1275:LEU:HB3	1.80	0.47
1:B:1092:THR:HG21	1:B:1124:THR:HG22	1.96	0.46
1:B:1082:VAL:HG13	1:B:1117:ALA:HB3	1.98	0.46
1:B:1328:ASN:HA	1:B:1331:ARG:HG2	1.97	0.46
1:A:1479:LEU:HD12	1:A:1507:LEU:HG	1.97	0.46
1:A:1368:GLU:HB2	1:A:1380:VAL:HB	2.00	0.44
1:A:1276:THR:HG23	2:A:1601:ANP:O4'	2.18	0.44
1:A:1101:VAL:HG22	3:A:1602:HFG:BR1	2.73	0.43
1:A:1147:TRP:CE3	1:A:1170:GLN:HB3	2.53	0.43
1:B:1276:THR:HG23	2:B:1601:ANP:O4'	2.18	0.43
1:B:1104:VAL:HB	1:B:1116:ILE:HG13	2.00	0.43
1:B:1093:HIS:HB2	1:B:1097:PHE:HD2	1.84	0.43
1:B:1150:VAL:HG21	1:B:1169:TRP:CE2	2.54	0.42
1:A:1016:LEU:N	6:A:1773:HOH:O	2.52	0.42
1:B:1030[B]:GLN:NE2	1:B:1030[B]:GLN:CA	2.81	0.42
1:A:1146:GLN:O	1:A:1170:GLN:HA	2.20	0.42
1:B:1219:GLY:HA2	1:B:1244:LEU:HA	2.02	0.41
1:B:1258:ASP:HA	1:B:1259:PRO:HD2	1.90	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1100:GLU:HB2	3:B:1602:HFG:BR1	2.75	0.41
1:B:1039:GLU:HB3	1:B:1048:ILE:HB	2.02	0.41
1:A:1150:VAL:HG21	1:A:1169:TRP:CE2	2.55	0.41
1:A:1186:VAL:HG21	1:A:1243:HIS:HB2	2.03	0.40
1:B:1186:VAL:HG21	1:B:1243:HIS:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	479/519 (92%)	471 (98%)	7 (2%)	1 (0%)	47	44
1	B	482/519 (93%)	473 (98%)	7 (2%)	2 (0%)	34	30
All	All	961/1038 (93%)	944 (98%)	14 (2%)	3 (0%)	41	37

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1497	CYS
1	B	1121	THR
1	A	1121	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	406/445 (91%)	400 (98%)	6 (2%)	65	69
1	B	402/445 (90%)	391 (97%)	11 (3%)	44	46
All	All	808/890 (91%)	791 (98%)	17 (2%)	53	57

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1169	TRP
1	A	1240	THR
1	A	1273	TRP
1	A	1324	ILE
1	A	1348	ASN
1	A	1504	TYR
1	B	1021	GLU
1	B	1082	VAL
1	B	1153	TRP
1	B	1169	TRP
1	B	1179	MET
1	B	1232	SER
1	B	1273	TRP
1	B	1323	LEU
1	B	1392	VAL
1	B	1497	CYS
1	B	1504	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1170	GLN
1	B	1170	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HFG	B	1602	-	25,26,26	2.19	5 (20%)	27,37,37	2.77	6 (22%)
2	ANP	A	1601	4	29,33,33	1.86	8 (27%)	31,52,52	1.88	10 (32%)
2	ANP	B	1601	4	29,33,33	1.89	6 (20%)	31,52,52	2.11	8 (25%)
3	HFG	A	1602	-	25,26,26	2.21	6 (24%)	27,37,37	3.12	8 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HFG	B	1602	-	-	0/8/19/19	0/3/3/3
2	ANP	A	1601	4	-	6/14/38/38	0/3/3/3
2	ANP	B	1601	4	-	5/14/38/38	0/3/3/3
3	HFG	A	1602	-	-	0/8/19/19	0/3/3/3

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1602	HFG	C10-C9	6.81	1.49	1.40
3	A	1602	HFG	C10-C9	6.47	1.48	1.40
3	B	1602	HFG	C6-C7	5.10	1.49	1.39
3	A	1602	HFG	C2-N1	5.03	1.35	1.29
3	A	1602	HFG	C6-C7	4.81	1.48	1.39
3	B	1602	HFG	C2-N1	4.79	1.34	1.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1601	ANP	PG-N3B	4.60	1.75	1.63
2	B	1601	ANP	PB-N3B	4.45	1.75	1.63
2	A	1601	ANP	PG-N3B	4.36	1.74	1.63
2	A	1601	ANP	PB-N3B	4.23	1.74	1.63
2	A	1601	ANP	PG-O1G	3.63	1.51	1.46
2	A	1601	ANP	PB-O1B	3.40	1.51	1.46
2	B	1601	ANP	PB-O1B	3.11	1.51	1.46
2	B	1601	ANP	PB-O3A	3.07	1.62	1.59
2	B	1601	ANP	PG-O1G	3.00	1.50	1.46
3	A	1602	HFG	C9-N1	-2.86	1.35	1.40
3	A	1602	HFG	C6-CL1	2.66	1.80	1.73
3	B	1602	HFG	C9-N1	-2.65	1.35	1.40
2	B	1601	ANP	C5-C4	2.59	1.47	1.40
2	A	1601	ANP	PB-O3A	2.52	1.62	1.59
3	B	1602	HFG	C6-CL1	2.32	1.79	1.73
2	A	1601	ANP	C5-C4	2.20	1.46	1.40
2	A	1601	ANP	PG-O2G	-2.15	1.51	1.56
2	A	1601	ANP	PB-O2B	-2.15	1.51	1.56
3	A	1602	HFG	C4-N3	2.02	1.41	1.39

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1602	HFG	C6'-N1'-C2'	10.73	118.31	111.62
3	B	1602	HFG	C6'-N1'-C2'	10.24	118.01	111.62
3	A	1602	HFG	C10-C4-N3	7.20	118.00	113.80
2	B	1601	ANP	O1G-PG-N3B	-7.02	101.44	111.77
3	A	1602	HFG	C9-N1-C2	5.72	122.25	116.62
3	B	1602	HFG	C10-C4-N3	5.59	117.06	113.80
3	B	1602	HFG	C9-N1-C2	5.21	121.76	116.62
3	A	1602	HFG	N3-C2-N1	-4.11	122.77	126.34
2	B	1601	ANP	O2B-PB-O1B	3.86	118.02	109.92
2	A	1601	ANP	N3-C2-N1	-3.75	122.82	128.68
2	B	1601	ANP	N3-C2-N1	-3.63	123.01	128.68
3	A	1602	HFG	C9-C10-C4	-3.50	116.32	119.51
2	A	1601	ANP	O2B-PB-O1B	3.48	117.22	109.92
2	A	1601	ANP	O1G-PG-N3B	-3.35	106.84	111.77
3	B	1602	HFG	N3-C2-N1	-3.12	123.62	126.34
2	A	1601	ANP	O2G-PG-O3G	3.09	115.86	107.64
2	A	1601	ANP	O4'-C4'-C5'	-3.04	99.38	109.37
2	B	1601	ANP	O4'-C4'-C5'	-2.84	100.03	109.37
2	B	1601	ANP	C3'-C2'-C1'	2.79	105.18	100.98

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1602	HFG	BR1-C7-C6	-2.74	117.80	121.29
2	B	1601	ANP	PB-O3A-PA	-2.73	122.99	132.62
2	A	1601	ANP	PB-O3A-PA	-2.71	123.07	132.62
2	A	1601	ANP	O1B-PB-N3B	-2.69	107.80	111.77
2	A	1601	ANP	C3'-C2'-C1'	2.61	104.90	100.98
3	B	1602	HFG	C9-C10-C4	-2.51	117.22	119.51
3	B	1602	HFG	C10-C9-N1	-2.49	118.99	122.54
2	A	1601	ANP	C1'-N9-C4	-2.47	122.31	126.64
2	A	1601	ANP	C4-C5-N7	-2.29	107.01	109.40
2	B	1601	ANP	C2-N1-C6	2.21	122.54	118.75
2	B	1601	ANP	C4-C5-N7	-2.19	107.12	109.40
3	A	1602	HFG	C5-C10-C9	2.09	122.32	120.03
3	A	1602	HFG	C10-C9-N1	-2.05	119.63	122.54

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1601	ANP	PB-N3B-PG-O1G
2	A	1601	ANP	C5'-O5'-PA-O1A
2	A	1601	ANP	C5'-O5'-PA-O3A
2	B	1601	ANP	C5'-O5'-PA-O3A
2	A	1601	ANP	C5'-O5'-PA-O2A
2	B	1601	ANP	C5'-O5'-PA-O1A
2	B	1601	ANP	C5'-O5'-PA-O2A
2	A	1601	ANP	C4'-C5'-O5'-PA
2	B	1601	ANP	C4'-C5'-O5'-PA
2	A	1601	ANP	O4'-C4'-C5'-O5'
2	B	1601	ANP	O4'-C4'-C5'-O5'


There are no ring outliers.


4 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1602	HFG	2	0
2	A	1601	ANP	1	0
2	B	1601	ANP	1	0
3	A	1602	HFG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	482/519 (92%)	0.47	45 (9%) 8 8	29, 48, 79, 99	0
1	B	483/519 (93%)	0.48	44 (9%) 9 8	30, 49, 78, 103	1 (0%)
All	All	965/1038 (92%)	0.48	89 (9%) 9 8	29, 48, 79, 103	1 (0%)

All (89) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1456	TRP	5.8
1	A	1489	LEU	5.3
1	A	1446	PRO	5.1
1	A	1505	TYR	4.4
1	A	1502	ALA	4.3
1	B	1437	LEU	4.2
1	A	1496	VAL	4.1
1	A	1458	LYS	4.1
1	B	1262	PRO	4.0
1	A	1425	VAL	4.0
1	B	1397	ALA	3.8
1	B	1498	GLY	3.6
1	B	1322	ALA	3.6
1	B	1475	GLY	3.5
1	A	1148	CYS	3.5
1	A	1016	LEU	3.5
1	B	1440	GLY	3.5
1	B	1395	ASN	3.5
1	B	1148	CYS	3.4
1	A	1460	THR	3.4
1	A	1447	PHE	3.4
1	B	1433	PHE	3.3
1	B	1456	TRP	3.3
1	B	1438	ASP	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1437	LEU	3.2
1	B	1462	ALA	3.1
1	A	1433	PHE	3.1
1	B	1076	CYS	3.1
1	A	1491	PRO	3.1
1	A	1501	PRO	3.1
1	B	1155	PHE	3.1
1	A	1497	CYS	3.0
1	A	1488	GLU	3.0
1	A	1487	CYS	3.0
1	B	1261	ILE	3.0
1	B	1334	LEU	3.0
1	B	1401	LEU	2.9
1	B	1329	ASP	2.9
1	B	1436	ILE	2.9
1	A	1445	ILE	2.9
1	A	1449	GLY	2.9
1	A	1150	VAL	2.8
1	B	1426	VAL	2.8
1	B	1460	THR	2.8
1	A	1427	ALA	2.7
1	A	1430	MET	2.7
1	A	1428	ASN	2.7
1	B	1458	LYS	2.7
1	A	1431	GLU	2.7
1	A	1452	ASP	2.7
1	A	1147	TRP	2.6
1	B	1496	VAL	2.6
1	A	1451	ILE	2.6
1	B	1080	MET	2.6
1	A	1151	VAL	2.5
1	A	1348	ASN	2.5
1	B	1151	VAL	2.5
1	A	1492	GLY	2.5
1	A	1390	LEU	2.5
1	A	1476	ALA	2.5
1	B	1390	LEU	2.5
1	B	1260	LYS	2.5
1	B	1330	TYR	2.4
1	A	1503	LYS	2.4
1	B	1396	GLU	2.4
1	A	1462	ALA	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1017	GLU	2.3
1	A	1441	LYS	2.3
1	B	1367	LEU	2.3
1	A	1057	TRP	2.3
1	A	1490	GLN	2.3
1	B	1370	GLY	2.3
1	B	1257[A]	GLU	2.3
1	B	1435	LYS	2.2
1	A	1168	LEU	2.2
1	B	1118	ILE	2.1
1	B	1324	ILE	2.1
1	B	1263	GLY	2.1
1	A	1457	ILE	2.1
1	A	1459	LYS	2.1
1	B	1337	VAL	2.1
1	A	1118	ILE	2.1
1	B	1486	LEU	2.1
1	A	1493	ALA	2.1
1	B	1323	LEU	2.0
1	B	1420	LEU	2.0
1	B	1430	MET	2.0
1	B	1425	VAL	2.0
1	B	1497	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	ZN	A	1604	1/1	0.86	0.05	73,73,73,73	0
4	MG	A	1603	1/1	0.87	0.19	39,39,39,39	0
3	HFG	B	1602	24/24	0.91	0.14	35,39,47,53	0
3	HFG	A	1602	24/24	0.92	0.18	34,37,44,49	0
2	ANP	B	1601	31/31	0.94	0.10	41,44,52,54	0
2	ANP	A	1601	31/31	0.95	0.14	35,43,52,53	0
5	ZN	B	1604	1/1	0.97	0.03	65,65,65,65	0
4	MG	B	1603	1/1	0.98	0.07	40,40,40,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around HFG A 1602:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ANP B 1601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.