



wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2020 – 05:27 am BST

PDB ID : 4J4L
Title : Modular evolution and design of the protein binding interface
Authors : Cheong, H.K.; Kim, H.J.
Deposited on : 2013-02-07
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	
1	B	267	
2	C	168	
2	D	168	

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 6751 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Internalin B, REPEAT MODULES, Variable lymphocyte receptor B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	258	Total 2035	C 1297	N 345	O 389	S 4	0	0	0
1	B	251	Total 1982	C 1264	N 336	O 378	S 4	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	59	ALA	ASP	conflict	UNP A4L9V2
A	302	LEU	-	expression tag	UNP Q4G1L3
A	303	GLU	-	expression tag	UNP Q4G1L3
A	304	HIS	-	expression tag	UNP Q4G1L3
A	305	HIS	-	expression tag	UNP Q4G1L3
A	306	HIS	-	expression tag	UNP Q4G1L3
A	307	HIS	-	expression tag	UNP Q4G1L3
A	308	HIS	-	expression tag	UNP Q4G1L3
A	309	HIS	-	expression tag	UNP Q4G1L3
B	59	ALA	ASP	conflict	UNP A4L9V2
B	302	LEU	-	expression tag	UNP Q4G1L3
B	303	GLU	-	expression tag	UNP Q4G1L3
B	304	HIS	-	expression tag	UNP Q4G1L3
B	305	HIS	-	expression tag	UNP Q4G1L3
B	306	HIS	-	expression tag	UNP Q4G1L3
B	307	HIS	-	expression tag	UNP Q4G1L3
B	308	HIS	-	expression tag	UNP Q4G1L3
B	309	HIS	-	expression tag	UNP Q4G1L3

- Molecule 2 is a protein called Interleukin-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	155	Total 1246	C 780	N 216	O 241	S 9	0	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	166	1330	832	229	260	9	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	17	GLY	-	expression tag	UNP P05231
C	18	SER	-	expression tag	UNP P05231
D	17	GLY	-	expression tag	UNP P05231
D	18	SER	-	expression tag	UNP P05231

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	61	61	61	0	0
3	B	28	28	28	0	0
3	C	37	37	37	0	0
3	D	32	32	32	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	46.32Å 134.00Å 148.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	61.09 – 2.30 61.09 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.2 (61.09-2.30) 95.2 (61.09-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.88 (at 2.29Å)	Xtrriage
Refinement program	CNS, REFMAC 5.6.0117	Depositor
R, R_{free}	0.205 , 0.268 0.207 , 0.264	Depositor DCC
R_{free} test set	2017 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	29.6	Xtrriage
Anisotropy	0.204	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 36.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6751	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.81	1/2075 (0.0%)	0.90	2/2821 (0.1%)
1	B	0.69	1/2020 (0.0%)	0.84	3/2745 (0.1%)
2	C	0.81	1/1259 (0.1%)	0.89	0/1687
2	D	0.76	0/1345	0.92	6/1807 (0.3%)
All	All	0.77	3/6699 (0.0%)	0.88	11/9060 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	246	TRP	CD2-CE2	6.02	1.48	1.41
2	C	157	TRP	CD2-CE2	5.96	1.48	1.41
1	B	246	TRP	CD2-CE2	5.13	1.47	1.41

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	30	ARG	NE-CZ-NH1	7.39	124.00	120.30
1	B	151	LEU	CA-CB-CG	6.76	130.84	115.30
2	D	24	ARG	NE-CZ-NH1	6.13	123.36	120.30
2	D	84	LEU	CB-CG-CD1	-6.08	100.66	111.00
2	D	30	ARG	NE-CZ-NH2	-5.94	117.33	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2035	0	2058	46	0
1	B	1982	0	2001	45	0
2	C	1246	0	1261	18	0
2	D	1330	0	1348	27	0
3	A	61	0	0	7	0
3	B	28	0	0	3	0
3	C	37	0	0	2	0
3	D	32	0	0	5	0
All	All	6751	0	6668	130	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 130 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:154:GLN:NE2	3:D:231:HOH:O	1.99	0.89
1:A:208:PRO:HB2	1:A:211:VAL:HG13	1.54	0.87
1:A:226:ASN:HB2	1:A:250:ASN:HD21	1.47	0.78
2:C:111:GLN:HE21	2:C:111:GLN:H	1.31	0.78
2:C:111:GLN:HE21	2:C:111:GLN:N	1.82	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	256/267 (96%)	241 (94%)	14 (6%)	1 (0%)	34 42
1	B	247/267 (92%)	224 (91%)	18 (7%)	5 (2%)	7 6
2	C	149/168 (89%)	146 (98%)	3 (2%)	0	100 100
2	D	164/168 (98%)	156 (95%)	7 (4%)	1 (1%)	25 31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	816/870 (94%)	767 (94%)	42 (5%)	7 (1%)	17 20

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	289	GLY
1	B	160	PRO
1	B	233	ASP
1	B	270	SER
2	D	131	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	233/242 (96%)	212 (91%)	21 (9%)	9 11
1	B	226/242 (93%)	205 (91%)	21 (9%)	9 10
2	C	142/152 (93%)	127 (89%)	15 (11%)	6 7
2	D	151/152 (99%)	135 (89%)	16 (11%)	6 7
All	All	752/788 (95%)	679 (90%)	73 (10%)	8 9

5 of 73 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	155	GLN
1	B	279	SER
2	D	134	ASP
1	B	211	VAL
2	C	37	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	150	GLN
1	B	217	GLN
2	D	127	GLN
1	B	169	ASN
1	B	193	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	258/267 (96%)	-0.24	0 100 100	11, 27, 45, 65	0
1	B	251/267 (94%)	-0.08	1 (0%) 92 95	17, 40, 64, 82	0
2	C	155/168 (92%)	0.02	7 (4%) 33 40	11, 29, 58, 83	1 (0%)
2	D	166/168 (98%)	-0.17	0 100 100	14, 29, 57, 76	1 (0%)
All	All	830/870 (95%)	-0.13	8 (0%) 82 86	11, 31, 58, 83	2 (0%)

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	50	CYS	3.8
2	C	153	ALA	2.7
2	C	106	GLU	2.6
2	C	48	ASN	2.5
2	C	140	ASP	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.