



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2023 – 12:31 PM EDT

PDB ID : 1K1B
Title : Crystal structure of the ankyrin repeat domain of Bcl-3: a unique member of the IkappaB protein family
Authors : Michel, F.; Soler-Lopez, M.; Petosa, C.; Cramer, P.; Siebenlist, U.; Mueller, C.W.
Deposited on : 2001-09-24
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	241	 71% 20% •• 5%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1915 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called B-cell lymphoma 3-encoded protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	228	1717	1058	325	326	8	5	2	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	198	Total	O	0	0
			198	198		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: B-cell lymphoma 3-encoded protein

Chain A: 



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	31.35Å 50.91Å 59.49Å 90.00° 102.04° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90	Depositor
% Data completeness (in resolution range)	97.6 (20.00-1.90)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.177 , 0.217	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1915	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.64	3/1750 (0.2%)	1.23	8/2382 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	343	SER	C-N	18.77	1.77	1.34
1	A	344	ARG	C-N	7.07	1.50	1.34
1	A	342	ARG	NE-CZ	-5.88	1.25	1.33

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	152	GLN	OE1-CD-NE2	-46.95	13.90	121.90
1	A	344	ARG	O-C-N	-8.33	109.38	122.70
1	A	344	ARG	NE-CZ-NH2	7.60	124.10	120.30
1	A	128	ASP	CB-CG-OD2	7.23	124.81	118.30
1	A	344	ARG	CA-C-N	5.80	129.97	117.20
1	A	264	ASP	CB-CG-OD2	5.31	123.08	118.30
1	A	233	ASP	CB-CG-OD2	5.12	122.91	118.30
1	A	126	ASP	CB-CG-OD2	5.10	122.89	118.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	125	GLU	Mainchain
1	A	152	GLN	Sidechain
1	A	344	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1717	0	1743	44	0
2	A	198	0	0	14	0
All	All	1915	0	1743	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:343:SER:C	1:A:344:ARG:N	1.77	1.37
1:A:332:CYS:HB3	2:A:556:HOH:O	1.37	1.25
1:A:322:ARG:NH1	2:A:547:HOH:O	1.84	1.06
1:A:210:PRO:HD2	2:A:373:HOH:O	1.92	0.70
1:A:128:ASP:HA	2:A:516:HOH:O	1.90	0.70
1:A:303:SER:H	1:A:306:HIS:HD2	1.40	0.70
1:A:303:SER:H	1:A:306:HIS:CD2	2.12	0.68
1:A:181:ARG:O	1:A:185:THR:HG23	1.95	0.67
1:A:198:GLN:NE2	1:A:231:ASN:HD22	1.93	0.66
1:A:343:SER:C	1:A:344:ARG:CA	2.64	0.66
1:A:126:ASP:C	1:A:126:ASP:OD1	2.36	0.64
1:A:166:THR:H	1:A:169:HIS:CD2	2.18	0.62
1:A:298:MET:HB3	2:A:374:HOH:O	2.00	0.60
1:A:347:ILE:O	1:A:351:ARG:HG2	2.01	0.59
1:A:182[B]:LEU:HD22	2:A:467:HOH:O	2.01	0.58
1:A:132:HIS:HE1	1:A:165:GLN:O	1.87	0.58
1:A:198:GLN:HE22	1:A:231:ASN:HD22	1.52	0.57
1:A:199:THR:H	1:A:202:HIS:CD2	2.22	0.57
1:A:199:THR:H	1:A:202:HIS:HD2	1.51	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246:CYS:N	2:A:555:HOH:O	2.38	0.55
1:A:343:SER:HB2	2:A:537:HOH:O	2.06	0.55
1:A:314:LEU:HB2	1:A:315:PRO:HD3	1.87	0.55
1:A:276:VAL:HG21	1:A:304:ALA:HB1	1.87	0.55
1:A:247:GLN:HG2	1:A:283:MET:HE1	1.90	0.54
1:A:333:HIS:HE1	2:A:496:HOH:O	1.90	0.53
1:A:236:THR:H	1:A:239:HIS:HD2	1.56	0.53
1:A:348:ASP:HA	1:A:351:ARG:HG3	1.91	0.52
1:A:247:GLN:HG2	1:A:283:MET:CE	2.40	0.50
1:A:217:LEU:HD13	1:A:256:ARG:HG3	1.94	0.49
1:A:166:THR:H	1:A:169:HIS:HD2	1.61	0.47
1:A:132:HIS:HD2	2:A:395:HOH:O	1.97	0.47
1:A:140:LEU:HB3	1:A:141:PRO:HD3	1.97	0.47
1:A:246:CYS:C	2:A:555:HOH:O	2.53	0.47
1:A:297:GLN:NE2	2:A:499:HOH:O	2.43	0.46
1:A:156:GLU:O	1:A:159:ILE:CG2	2.64	0.46
1:A:240:VAL:O	1:A:244:THR:HG23	2.15	0.46
1:A:140:LEU:HD12	1:A:179:VAL:HG22	1.99	0.45
1:A:129:THR:HB	1:A:132:HIS:CD2	2.53	0.44
1:A:247:GLN:HG3	2:A:508:HOH:O	2.18	0.44
1:A:306:HIS:HE1	1:A:335:ASP:O	2.01	0.44
1:A:343:SER:CA	1:A:344:ARG:N	2.74	0.43
1:A:266:LYS:HB3	2:A:447:HOH:O	2.18	0.42
1:A:202:HIS:HE1	1:A:235:LEU:O	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	228/241 (95%)	225 (99%)	3 (1%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	190/197 (96%)	180 (95%)	10 (5%)	22 13

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	126	ASP
1	A	128	ASP
1	A	140	LEU
1	A	149	LEU
1	A	152	GLN
1	A	159	ILE
1	A	313	LEU
1	A	322	ARG
1	A	338	LEU
1	A	351	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	132	HIS
1	A	151	GLN
1	A	169	HIS
1	A	198	GLN
1	A	202	HIS
1	A	239	HIS
1	A	297	GLN
1	A	306	HIS
1	A	334	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	343:SER	C	344:ARG	N	1.77

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.