



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 01:21 am BST

PDB ID : 5L0L
Title : Crystal structure of Uncharacterized protein LPG0439
Authors : Chang, C.; Skarina, T.; Khutoreskaya, G.; Savchenko, A.; Joachimiak, A.
Deposited on : 2016-07-27
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	 8% 93% 7%
1	B	215	 14% 93% 7%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3887 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	215	Total 1770	C 1123	N 301	O 336	S 3	Se 7	0	7	0
1	B	215	Total 1780	C 1129	N 303	O 338	S 3	Se 7	0	9	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	Total 5	O 4	S 1	0	0
2	B	1	Total 5	O 4	S 1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Cl 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	169	Total O 169 169	0	0
4	B	157	Total O 157 157	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uncharacterized protein



- Molecule 1: Uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, α , β , γ	129.69Å 129.69Å 117.14Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.01 – 1.80 39.91 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.01-1.80) 99.9 (39.91-1.80)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.12 (at 1.79Å)	Xtrriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.169 , 0.189 0.169 , 0.190	Depositor DCC
R_{free} test set	2752 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	20.8	Xtrriage
Anisotropy	0.585	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3887	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.59	0/1805	0.77	3/2430 (0.1%)
1	B	0.59	0/1815	0.81	5/2445 (0.2%)
All	All	0.59	0/3620	0.79	8/4875 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	241	ARG	NE-CZ-NH1	6.97	123.79	120.30
1	A	241	ARG	NE-CZ-NH2	-6.88	116.86	120.30
1	B	250	ASP	CB-CG-OD1	6.85	124.47	118.30
1	B	241	ARG	NE-CZ-NH1	6.45	123.53	120.30
1	B	241	ARG	NE-CZ-NH2	-5.61	117.50	120.30
1	B	125	ARG	NE-CZ-NH1	5.51	123.05	120.30
1	B	250	ASP	CB-CG-OD2	-5.29	113.54	118.30
1	A	125	ARG	NE-CZ-NH1	5.02	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1770	0	1729	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1780	0	1742	11	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	B	1	0	0	0	0
4	A	169	0	0	1	0
4	B	157	0	0	0	1
All	All	3887	0	3471	16	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288[B]:ASN:ND2	4:A:501:HOH:O	2.21	0.65
1:A:232[A]:MSE:HE2	1:B:95:LYS:HD2	1.83	0.59
1:A:232[B]:MSE:SE	1:B:95:LYS:CD	3.01	0.58
1:A:232[B]:MSE:SE	1:B:95:LYS:HE2	2.60	0.52
1:A:95:LYS:HE2	1:B:232[B]:MSE:SE	2.62	0.49
1:A:238:VAL:HG11	1:A:275[B]:CYS:SG	2.52	0.49
1:B:238:VAL:HG21	1:B:275[B]:CYS:SG	2.54	0.48
1:B:276[B]:LEU:HG	1:B:280:LYS:HE3	1.98	0.46
1:A:295:GLU:OE1	1:B:170:ARG:HD2	2.17	0.45
1:A:276[B]:LEU:HG	1:A:280:LYS:HE3	1.98	0.45
1:B:283:LEU:O	1:B:287[B]:VAL:HG23	2.17	0.44
1:A:276[B]:LEU:HA	1:A:276[B]:LEU:HD12	1.79	0.41
1:B:238:VAL:HG11	1:B:275[B]:CYS:SG	2.60	0.41
1:A:124:VAL:HG22	1:A:300:LEU:CD2	2.51	0.41
1:A:232[B]:MSE:SE	1:B:95:LYS:HD2	2.71	0.41
1:A:211:ILE:HD11	1:B:189:ILE:HD11	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:618:HOH:O	4:B:618:HOH:O[11_655]	1.43	0.77

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	220/215 (102%)	215 (98%)	4 (2%)	1 (0%)	29	15
1	B	222/215 (103%)	218 (98%)	3 (1%)	1 (0%)	29	15
All	All	442/430 (103%)	433 (98%)	7 (2%)	2 (0%)	29	15

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	252	ILE
1	A	252	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/182 (107%)	190 (98%)	4 (2%)	53	42
1	B	196/182 (108%)	194 (99%)	2 (1%)	76	71
All	All	390/364 (107%)	384 (98%)	6 (2%)	69	56

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	147	ARG
1	A	232[A]	MSE
1	A	232[B]	MSE
1	A	298	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	102	GLU
1	B	147	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	305	HIS
1	B	261	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	401	-	4,4,4	0.35	0	6,6,6	1.79	2 (33%)
2	SO4	B	401	-	4,4,4	0.55	0	6,6,6	1.01	0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	SO4	O4-S-O1	-2.49	96.30	109.31
2	A	401	SO4	O4-S-O3	2.23	118.57	109.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	209/215 (97%)	0.30	17 (8%) 12 9	13, 21, 50, 65	0
1	B	209/215 (97%)	0.50	31 (14%) 2 1	13, 23, 58, 77	0
All	All	418/430 (97%)	0.40	48 (11%) 4 3	13, 22, 56, 77	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	94	GLY	11.0
1	B	258	HIS	6.9
1	B	108	TRP	5.9
1	B	105	PRO	5.5
1	B	106	GLY	4.7
1	A	94	GLY	4.6
1	A	308	LYS	4.5
1	B	104	SER	4.5
1	A	108	TRP	4.4
1	B	259	PRO	4.4
1	A	106	GLY	4.0
1	B	298	ASP	3.8
1	A	298	ASP	3.7
1	B	103	TYR	3.6
1	A	105	PRO	3.6
1	B	102	GLU	3.5
1	B	214	GLU	3.3
1	B	112	TYR	3.2
1	B	101	HIS	3.2
1	B	114	ARG	3.2
1	B	153	ASP	3.2
1	B	215	GLU	3.2
1	A	153	ASP	3.0
1	B	257	THR	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	126	ASN	2.9
1	A	114	ARG	2.9
1	A	102	GLU	2.9
1	B	308	LYS	2.8
1	B	192	PHE	2.8
1	B	294	GLU	2.7
1	B	111	ASP	2.7
1	B	115	ALA	2.7
1	B	100	LYS	2.6
1	B	107	HIS	2.6
1	A	104	SER	2.6
1	B	290	PRO	2.5
1	B	260	ILE	2.4
1	A	294	GLU	2.4
1	B	222	LEU	2.3
1	A	214	GLU	2.2
1	A	111	ASP	2.2
1	A	107	HIS	2.2
1	B	110	ILE	2.1
1	B	256	LYS	2.1
1	A	192	PHE	2.1
1	B	109	SER	2.1
1	B	176	ALA	2.0
1	A	307	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	SO4	A	401	5/5	0.98	0.11	21,23,24,29	0
2	SO4	B	401	5/5	0.98	0.09	19,20,20,24	0
3	CL	B	402	1/1	0.99	0.04	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.