



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 11, 2023 – 06:28 AM EDT

PDB ID : 7L50
Title : Crystal structure of human monoacylglycerol lipase in complex with compound 4f
Authors : Qin, L.; Lane, W.; Skene, R.J.
Deposited on : 2020-12-21
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	 3% 76% 12% 11%
1	B	320	 4% 79% 12% 9%
1	C	320	 7% 77% 12% 11%
1	D	320	 6% 81% 8% 11%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9153 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Monoglyceride lipase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	285	2205	1408	381	403	13	0	0	0
1	B	290	2238	1427	389	409	13	0	0	0
1	C	286	2213	1412	385	403	13	0	0	0
1	D	285	2208	1409	384	402	13	0	0	0

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-16	MET	-	initiating methionine	UNP A0A0C4DFN3
A	-15	HIS	-	expression tag	UNP A0A0C4DFN3
A	-14	HIS	-	expression tag	UNP A0A0C4DFN3
A	-13	HIS	-	expression tag	UNP A0A0C4DFN3
A	-12	HIS	-	expression tag	UNP A0A0C4DFN3
A	-11	HIS	-	expression tag	UNP A0A0C4DFN3
A	-10	HIS	-	expression tag	UNP A0A0C4DFN3
A	-9	GLY	-	expression tag	UNP A0A0C4DFN3
A	-8	SER	-	expression tag	UNP A0A0C4DFN3
A	-7	GLU	-	expression tag	UNP A0A0C4DFN3
A	-6	ASN	-	expression tag	UNP A0A0C4DFN3
A	-5	LEU	-	expression tag	UNP A0A0C4DFN3
A	-4	TYR	-	expression tag	UNP A0A0C4DFN3
A	-3	PHE	-	expression tag	UNP A0A0C4DFN3
A	-2	GLN	-	expression tag	UNP A0A0C4DFN3
A	-1	GLY	-	expression tag	UNP A0A0C4DFN3
A	36	ALA	LYS	conflict	UNP A0A0C4DFN3
A	169	SER	LEU	conflict	UNP A0A0C4DFN3
A	176	SER	LEU	conflict	UNP A0A0C4DFN3
B	-16	MET	-	initiating methionine	UNP A0A0C4DFN3
B	-15	HIS	-	expression tag	UNP A0A0C4DFN3

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP A0A0C4DFN3
B	-13	HIS	-	expression tag	UNP A0A0C4DFN3
B	-12	HIS	-	expression tag	UNP A0A0C4DFN3
B	-11	HIS	-	expression tag	UNP A0A0C4DFN3
B	-10	HIS	-	expression tag	UNP A0A0C4DFN3
B	-9	GLY	-	expression tag	UNP A0A0C4DFN3
B	-8	SER	-	expression tag	UNP A0A0C4DFN3
B	-7	GLU	-	expression tag	UNP A0A0C4DFN3
B	-6	ASN	-	expression tag	UNP A0A0C4DFN3
B	-5	LEU	-	expression tag	UNP A0A0C4DFN3
B	-4	TYR	-	expression tag	UNP A0A0C4DFN3
B	-3	PHE	-	expression tag	UNP A0A0C4DFN3
B	-2	GLN	-	expression tag	UNP A0A0C4DFN3
B	-1	GLY	-	expression tag	UNP A0A0C4DFN3
B	36	ALA	LYS	conflict	UNP A0A0C4DFN3
B	169	SER	LEU	conflict	UNP A0A0C4DFN3
B	176	SER	LEU	conflict	UNP A0A0C4DFN3
C	-16	MET	-	initiating methionine	UNP A0A0C4DFN3
C	-15	HIS	-	expression tag	UNP A0A0C4DFN3
C	-14	HIS	-	expression tag	UNP A0A0C4DFN3
C	-13	HIS	-	expression tag	UNP A0A0C4DFN3
C	-12	HIS	-	expression tag	UNP A0A0C4DFN3
C	-11	HIS	-	expression tag	UNP A0A0C4DFN3
C	-10	HIS	-	expression tag	UNP A0A0C4DFN3
C	-9	GLY	-	expression tag	UNP A0A0C4DFN3
C	-8	SER	-	expression tag	UNP A0A0C4DFN3
C	-7	GLU	-	expression tag	UNP A0A0C4DFN3
C	-6	ASN	-	expression tag	UNP A0A0C4DFN3
C	-5	LEU	-	expression tag	UNP A0A0C4DFN3
C	-4	TYR	-	expression tag	UNP A0A0C4DFN3
C	-3	PHE	-	expression tag	UNP A0A0C4DFN3
C	-2	GLN	-	expression tag	UNP A0A0C4DFN3
C	-1	GLY	-	expression tag	UNP A0A0C4DFN3
C	36	ALA	LYS	conflict	UNP A0A0C4DFN3
C	169	SER	LEU	conflict	UNP A0A0C4DFN3
C	176	SER	LEU	conflict	UNP A0A0C4DFN3
D	-16	MET	-	initiating methionine	UNP A0A0C4DFN3
D	-15	HIS	-	expression tag	UNP A0A0C4DFN3
D	-14	HIS	-	expression tag	UNP A0A0C4DFN3
D	-13	HIS	-	expression tag	UNP A0A0C4DFN3
D	-12	HIS	-	expression tag	UNP A0A0C4DFN3
D	-11	HIS	-	expression tag	UNP A0A0C4DFN3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-10	HIS	-	expression tag	UNP A0A0C4DFN3
D	-9	GLY	-	expression tag	UNP A0A0C4DFN3
D	-8	SER	-	expression tag	UNP A0A0C4DFN3
D	-7	GLU	-	expression tag	UNP A0A0C4DFN3
D	-6	ASN	-	expression tag	UNP A0A0C4DFN3
D	-5	LEU	-	expression tag	UNP A0A0C4DFN3
D	-4	TYR	-	expression tag	UNP A0A0C4DFN3
D	-3	PHE	-	expression tag	UNP A0A0C4DFN3
D	-2	GLN	-	expression tag	UNP A0A0C4DFN3
D	-1	GLY	-	expression tag	UNP A0A0C4DFN3
D	36	ALA	LYS	conflict	UNP A0A0C4DFN3
D	169	SER	LEU	conflict	UNP A0A0C4DFN3
D	176	SER	LEU	conflict	UNP A0A0C4DFN3

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is (2s,4R)-2-{3-[(3-chloro-4-methylphenyl)methoxy]azetidine-1-carbonyl}-7-oxa-5-azaspiro[3.4]octan-6-one (three-letter code: XOM) (formula: C₁₈H₂₁ClN₂O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total	C	Cl	N	O	0	0
			25	18	1	2	4		
3	B	1	Total	C	Cl	N	O	0	0
			25	18	1	2	4		
3	C	1	Total	C	Cl	N	O	0	0
			25	18	1	2	4		
3	D	1	Total	C	Cl	N	O	0	0
			25	18	1	2	4		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	77	Total	O	0	0
			77	77		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	60	Total O 60 60	0	0
4	C	14	Total O 14 14	0	0
4	D	6	Total O 6 6	0	0

- Molecule 1: Monoglyceride lipase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.89Å 127.17Å 119.67Å 90.00° 73.67° 90.00°	Depositor
Resolution (Å)	48.19 – 2.30 48.15 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.7 (48.19-2.30) 97.7 (48.15-2.30)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.8.0257	Depositor
R, R_{free}	0.239 , 0.269 0.250 , 0.280	Depositor DCC
R_{free} test set	3834 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	45.8	Xtrriage
Anisotropy	0.345	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 49.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.056 for h,-k,h-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9153	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.19 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0283e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, XOM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/2257	0.68	0/3063
1	B	0.33	0/2291	0.69	0/3110
1	C	0.27	0/2265	0.55	0/3073
1	D	0.27	0/2260	0.54	0/3066
All	All	0.30	0/9073	0.62	0/12312

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2205	0	2204	33	0
1	B	2238	0	2235	24	0
1	C	2213	0	2216	23	0
1	D	2208	0	2211	17	0
2	A	8	0	6	0	0
2	B	8	0	6	0	0
2	C	8	0	6	0	0
2	D	8	0	6	0	0
3	A	25	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	25	0	0	1	0
3	C	25	0	0	0	0
3	D	25	0	0	2	0
4	A	77	0	0	2	0
4	B	60	0	0	0	0
4	C	14	0	0	0	0
4	D	6	0	0	0	0
All	All	9153	0	8890	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 96 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:LEU:HD12	1:A:22:LEU:O	1.82	0.78
1:A:116:VAL:HG13	1:A:139:PHE:HD1	1.50	0.76
1:B:202:ARG:HG3	1:B:202:ARG:HH21	1.51	0.76
1:B:116:VAL:HG13	1:B:139:PHE:CD1	2.28	0.68
1:A:22:LEU:HD12	1:A:22:LEU:C	2.13	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	281/320 (88%)	269 (96%)	12 (4%)	0	100 100
1	B	286/320 (89%)	269 (94%)	17 (6%)	0	100 100
1	C	282/320 (88%)	270 (96%)	11 (4%)	1 (0%)	34 42
1	D	281/320 (88%)	272 (97%)	9 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1130/1280 (88%)	1080 (96%)	49 (4%)	1 (0%)	51 64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	177	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	240/271 (89%)	234 (98%)	6 (2%)	47 65
1	B	244/271 (90%)	241 (99%)	3 (1%)	71 84
1	C	241/271 (89%)	237 (98%)	4 (2%)	60 76
1	D	241/271 (89%)	238 (99%)	3 (1%)	71 84
All	All	966/1084 (89%)	950 (98%)	16 (2%)	60 76

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	274	GLU
1	D	251	MET
1	B	274	GLU
1	C	291	SER
1	B	218	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	HIS
1	A	212	GLN
1	A	215	ASN
1	B	108	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	212	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ACT	A	601	-	3,3,3	1.02	0	3,3,3	0.82	0
3	XOM	C	603	-	22,28,28	1.47	5 (22%)	23,42,42	1.91	6 (26%)
3	XOM	D	603	-	22,28,28	1.32	4 (18%)	23,42,42	1.61	4 (17%)
2	ACT	A	602	-	3,3,3	1.03	0	3,3,3	0.75	0
3	XOM	A	603	-	22,28,28	1.66	3 (13%)	23,42,42	2.25	6 (26%)
2	ACT	B	601	-	3,3,3	1.04	0	3,3,3	0.79	0
2	ACT	C	602	-	3,3,3	1.02	0	3,3,3	0.77	0
2	ACT	B	602	-	3,3,3	1.05	0	3,3,3	0.73	0
2	ACT	D	601	-	3,3,3	1.05	0	3,3,3	0.77	0
2	ACT	C	601	-	3,3,3	0.98	0	3,3,3	0.80	0
2	ACT	D	602	-	3,3,3	0.97	0	3,3,3	0.84	0
3	XOM	B	603	-	22,28,28	1.62	4 (18%)	23,42,42	2.25	8 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	XOM	A	603	-	-	3/13/42/42	0/4/4/4
3	XOM	D	603	-	-	2/13/42/42	0/4/4/4
3	XOM	C	603	-	-	5/13/42/42	0/4/4/4
3	XOM	B	603	-	-	3/13/42/42	0/4/4/4

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	603	XOM	O23-C7	5.06	1.39	1.35
3	B	603	XOM	O23-C7	4.79	1.39	1.35
3	C	603	XOM	C7-N19	3.67	1.38	1.34
3	B	603	XOM	C7-N19	3.67	1.38	1.34
3	A	603	XOM	C7-N19	3.47	1.38	1.34

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	603	XOM	O23-C7-N19	-5.73	106.38	109.87
3	C	603	XOM	O23-C7-N19	-5.32	106.63	109.87
3	A	603	XOM	O23-C7-O21	5.04	125.61	121.30
3	B	603	XOM	C18-O24-C15	-4.91	104.91	113.78
3	B	603	XOM	C14-C8-N20	-4.35	113.81	118.80

There are no chirality outliers.

5 of 13 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	603	XOM	C12-C15-O24-C18
3	D	603	XOM	C12-C15-O24-C18
3	B	603	XOM	C4-C18-O24-C15
3	A	603	XOM	C4-C18-O24-C15
3	A	603	XOM	C10-C14-C8-O22


There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	603	XOM	2	0
3	A	603	XOM	1	0
3	B	603	XOM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	285/320 (89%)	0.23	11 (3%) 39 46	34, 57, 109, 162	0
1	B	290/320 (90%)	0.29	12 (4%) 37 44	35, 58, 121, 169	0
1	C	286/320 (89%)	0.50	23 (8%) 12 16	68, 96, 146, 186	0
1	D	285/320 (89%)	0.44	18 (6%) 20 25	73, 96, 144, 182	0
All	All	1146/1280 (89%)	0.37	64 (5%) 24 30	34, 84, 138, 186	0

The worst 5 of 64 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	161	VAL	7.2
1	D	178	PRO	4.8
1	C	165	LYS	4.8
1	B	295	ALA	4.5
1	C	177	GLY	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ACT	B	601	4/4	0.78	0.17	92,97,98,101	0
2	ACT	A	601	4/4	0.79	0.19	95,96,98,101	0
2	ACT	A	602	4/4	0.82	0.27	98,100,101,101	0
2	ACT	D	601	4/4	0.82	0.23	107,110,112,114	0
2	ACT	D	602	4/4	0.84	0.24	126,131,133,134	0
2	ACT	C	602	4/4	0.86	0.21	111,115,117,118	0
3	XOM	C	603	25/25	0.88	0.20	89,94,122,133	0
2	ACT	C	601	4/4	0.89	0.24	121,126,126,126	0
2	ACT	B	602	4/4	0.92	0.18	90,92,92,94	0
3	XOM	A	603	25/25	0.95	0.17	45,56,93,111	0
3	XOM	B	603	25/25	0.96	0.17	44,55,93,103	0
3	XOM	D	603	25/25	0.96	0.15	78,87,134,150	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around XOM A 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around XOM B 603:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.