



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2021 – 01:10 PM EDT

PDB ID : 2MEA
Title : CHANGES IN CONFORMATIONAL STABILITY OF A SERIES OF MUTANT HUMAN LYSOZYMES AT CONSTANT POSITIONS
Authors : Funahashi, J.; Takano, K.; Yamagata, Y.; Yutani, K.
Deposited on : 1998-05-02
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	130	
1	B	130	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2265 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LYSOZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	130	1032	636	200	186	10	0	0	0
1	B	130	1032	636	200	186	10	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	56	PHE	ILE	engineered mutation	UNP P61626
B	56	PHE	ILE	engineered mutation	UNP P61626

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	106	106	106	0	0
2	B	95	95	95	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: LYSOZYME

Chain A:  79% 18%



- Molecule 1: LYSOZYME

Chain B:  81% 18%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.74Å 110.55Å 43.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.20	Depositor
% Data completeness (in resolution range)	82.9 (8.00-2.20)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.172 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2265	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/1053	0.69	0/1421
1	B	0.41	0/1053	0.68	0/1421
All	All	0.41	0/2106	0.68	0/2842

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1032	0	990	18	0
1	B	1032	0	990	22	0
2	A	106	0	0	3	0
2	B	95	0	0	2	0
All	All	2265	0	1980	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:107:ARG:HH11	1:B:117:GLN:HE22	1.33	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:122:ARG:O	1:A:126:GLN:HG2	1.88	0.72
1:B:50:ARG:HD3	2:B:300:HOH:O	1.93	0.68
1:B:123:GLN:HA	1:B:126:GLN:HG2	1.76	0.66
1:A:33:LYS:HG3	1:B:101:ARG:HH12	1.60	0.65
1:B:123:GLN:HG3	2:B:178:HOH:O	1.96	0.64
1:A:110:VAL:HG23	2:A:168:HOH:O	1.97	0.64
1:A:34:TRP:CE3	1:A:115:ARG:HD3	2.35	0.61
1:B:107:ARG:NH1	1:B:117:GLN:HE22	2.00	0.59
1:A:33:LYS:HG3	1:B:101:ARG:NH1	2.18	0.58
1:A:62:ARG:HG2	1:A:62:ARG:HH11	1.68	0.57
1:B:107:ARG:HH11	1:B:117:GLN:NE2	2.02	0.56
1:A:33:LYS:HE2	2:A:394:HOH:O	2.08	0.54
1:A:62:ARG:HG2	1:A:62:ARG:NH1	2.23	0.53
1:A:34:TRP:CD2	1:A:115:ARG:HD3	2.44	0.52
1:B:39:ASN:OD1	1:B:41:ARG:HD3	2.09	0.52
1:A:95:CYS:O	1:A:99:VAL:HG23	2.09	0.52
1:A:64:TRP:O	1:A:77:CYS:HB2	2.10	0.51
1:B:128:CYS:HB2	1:B:130:VAL:HG22	1.92	0.50
1:A:13:LYS:HZ3	1:A:25:LEU:HD22	1.75	0.49
1:B:122:ARG:O	1:B:126:GLN:HG2	2.14	0.48
1:B:98:ARG:HA	1:B:101:ARG:HE	1.78	0.48
1:A:31:LEU:O	1:A:35:GLU:HG2	2.14	0.47
1:B:111:ALA:O	1:B:115:ARG:HG2	2.14	0.46
1:A:21:ARG:HG3	2:A:284:HOH:O	2.15	0.45
1:B:33:LYS:C	1:B:33:LYS:HD3	2.37	0.45
1:B:15:LEU:CD1	1:B:93:VAL:HG21	2.48	0.44
1:B:39:ASN:HD21	1:B:41:ARG:NH1	2.15	0.44
1:B:64:TRP:CE3	1:B:76:ALA:HB3	2.55	0.42
1:B:15:LEU:HD12	1:B:93:VAL:HG21	2.02	0.42
1:B:63:TYR:HA	1:B:74:VAL:HG22	2.01	0.42
1:A:6:CYS:O	1:A:10:ARG:HG3	2.20	0.42
1:A:107:ARG:NH1	1:A:117:GLN:HE22	2.18	0.41
1:B:64:TRP:O	1:B:77:CYS:HB2	2.21	0.41
1:B:123:GLN:HA	1:B:126:GLN:CG	2.47	0.41
1:B:84:LEU:CD2	1:B:92:ALA:HA	2.51	0.40
1:A:36:SER:OG	1:A:56:PHE:HA	2.22	0.40
1:A:37:GLY:O	1:A:38:TYR:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	128/130 (98%)	122 (95%)	6 (5%)	0	100	100
1	B	128/130 (98%)	124 (97%)	4 (3%)	0	100	100
All	All	256/260 (98%)	246 (96%)	10 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	105/105 (100%)	99 (94%)	6 (6%)	20	24
1	B	105/105 (100%)	102 (97%)	3 (3%)	42	54
All	All	210/210 (100%)	201 (96%)	9 (4%)	29	36

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	LYS
1	A	21	ARG
1	A	50	ARG
1	A	79	LEU
1	A	115	ARG
1	A	119	ARG
1	B	41	ARG
1	B	102	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	123	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	HIS
1	A	117	GLN
1	A	118	ASN
1	B	86	GLN
1	B	117	GLN
1	B	118	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.