



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 04:11 pm BST

PDB ID : 5MS2  
Title : Crystal structure of the Legionella pneumophila effector protein RavZ in complex with human LC3B  
Authors : Pantoom, S.; Vetter, I.R.; Wu, Y.W.  
Deposited on : 2016-12-30  
Resolution : 2.47 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5857 (2.50-2.46)
Clashscore	141614	6594 (2.50-2.46)
Ramachandran outliers	138981	6469 (2.50-2.46)
Sidechain outliers	138945	6471 (2.50-2.46)
RSRZ outliers	127900	5738 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	
2	B	124	

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 4273 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Legionella pneumophila effector protein RavZ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	405	3202	2029	518	645	10	0	2	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q5ZUV9
A	0	PRO	-	expression tag	UNP Q5ZUV9

- Molecule 2 is a protein called Microtubule-associated proteins 1A/1B light chain 3B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	115	960	614	168	175	3	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	GLY	-	expression tag	UNP Q9GZQ8
B	-3	HIS	-	expression tag	UNP Q9GZQ8
B	-2	MET	-	expression tag	UNP Q9GZQ8
B	-1	GLY	-	expression tag	UNP Q9GZQ8
B	0	CYS	-	expression tag	UNP Q9GZQ8

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	106	Total	O	0	0
			106	106		
3	B	5	Total	O	0	0
			5	5		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.55Å 69.55Å 90.17Å 90.00° 101.08° 90.00°	Depositor
Resolution (Å)	48.08 – 2.47 48.08 – 2.47	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.08-2.47) 99.9 (48.08-2.47)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.24 (at 2.48Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.177 , 0.238 0.180 , 0.237	Depositor DCC
$R_{free}$ test set	1131 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.3	Xtrriage
Anisotropy	0.585	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 57.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4273	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/3267	0.70	3/4413 (0.1%)
2	B	0.41	0/978	0.72	0/1316
All	All	0.49	0/4245	0.70	3/5729 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	4
All	All	0	5

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	251	ASN	N-CA-C	-9.06	86.54	111.00
1	A	29	PHE	CA-C-O	7.62	136.11	120.10
1	A	29	PHE	CA-C-N	-7.27	101.21	117.20

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	250	GLN	Peptide
2	B	21	ARG	Peptide
2	B	22	LEU	Peptide
2	B	48	ASP	Peptide
2	B	73	LEU	Peptide

## 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3202	0	3173	53	1
2	B	960	0	979	46	0
3	A	106	0	0	0	0
3	B	5	0	0	3	0
All	All	4273	0	4152	97	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:24:ARG:NH1	2:B:25:GLU:OE1	1.98	0.97
2:B:82:LEU:HD12	2:B:87:SER:HB3	1.61	0.82
2:B:45:PRO:HG2	2:B:71:LEU:HD22	1.61	0.81
1:A:107:ASN:HB3	1:A:109:GLN:H	1.46	0.80
2:B:87:SER:OG	3:B:201:HOH:O	2.02	0.78
2:B:76:ASN:OD1	3:B:202:HOH:O	2.03	0.76
1:A:251:ASN:OD1	1:A:289:SER:OG	2.04	0.75
1:A:34:GLY:HA3	2:B:76:ASN:HD21	1.51	0.74
1:A:63:MET:HG3	1:A:259:GLY:HA3	1.71	0.72
2:B:70:ARG:O	2:B:72:GLN:NE2	2.23	0.71
2:B:13:PHE:HE2	2:B:105:GLU:HG3	1.58	0.69
2:B:26:GLN:N	2:B:26:GLN:OE1	2.28	0.67
2:B:10:ARG:HH22	2:B:50:THR:HG21	1.60	0.66
1:A:25:ASP:OD1	1:A:26:ILE:N	2.27	0.66
2:B:37:ARG:NH2	2:B:41:GLU:O	2.29	0.66
1:A:34:GLY:HA3	2:B:76:ASN:ND2	2.09	0.66
2:B:68:ARG:NH1	2:B:75:ALA:HA	2.14	0.63
2:B:37:ARG:HD3	2:B:46:VAL:HG22	1.81	0.61
2:B:22:LEU:HD11	2:B:25:GLU:H	1.68	0.59
2:B:11:ARG:HD3	2:B:15:GLN:HB3	1.84	0.59
2:B:24:ARG:HH11	2:B:25:GLU:CD	2.05	0.59
1:A:90:VAL:HG11	1:A:320:THR:O	2.04	0.57
2:B:25:GLU:HA	2:B:28:PRO:HA	1.85	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:109:GLN:NE2	1:A:158:THR:OG1	2.38	0.57
1:A:83:TYR:CZ	1:A:85:GLY:HA3	2.39	0.57
1:A:204:TYR:CD2	1:A:218:THR:HG21	2.40	0.57
2:B:22:LEU:CD1	2:B:25:GLU:H	2.19	0.55
1:A:294[A]:THR:HB	1:A:297:HIS:ND1	2.22	0.55
2:B:83:VAL:HG22	2:B:111:MET:HG2	1.89	0.54
1:A:387:TYR:CZ	1:A:423:LEU:HD13	2.43	0.54
1:A:29:PHE:CD1	1:A:31:LEU:HB2	2.43	0.54
2:B:73:LEU:HD23	2:B:74:ASN:H	1.72	0.54
1:A:294[A]:THR:HG22	1:A:296:GLN:H	1.72	0.54
1:A:423:LEU:O	1:A:426:VAL:HG12	2.07	0.53
1:A:163:PHE:CE1	1:A:178:VAL:HG13	2.43	0.53
2:B:67:ILE:HD12	2:B:81:LEU:HD11	1.89	0.53
2:B:8:LYS:NZ	2:B:104:ASP:OD2	2.35	0.52
1:A:126:VAL:HG11	1:A:208:LEU:HD13	1.91	0.52
1:A:201:ARG:HG2	1:A:252:ARG:HG3	1.92	0.52
1:A:33:GLU:HB2	3:B:202:HOH:O	2.10	0.51
1:A:28:GLU:HG2	1:A:29:PHE:H	1.74	0.51
1:A:294[B]:THR:HG22	1:A:297:HIS:ND1	2.26	0.51
2:B:116:GLN:HG2	2:B:118:THR:H	1.77	0.50
2:B:22:LEU:HG	2:B:26:GLN:OE1	2.12	0.50
2:B:66:ILE:O	2:B:70:ARG:HG3	2.12	0.50
2:B:30:LYS:HE3	2:B:55:PRO:HG3	1.94	0.50
2:B:20:VAL:HG11	2:B:107:GLY:HA3	1.94	0.50
1:A:294[B]:THR:HG23	1:A:296:GLN:H	1.78	0.49
2:B:117:GLU:O	2:B:117:GLU:HG2	2.13	0.49
1:A:201:ARG:NH1	1:A:254:THR:HG22	2.28	0.49
2:B:30:LYS:HB3	2:B:53:LEU:HG	1.95	0.49
2:B:11:ARG:NH1	2:B:19:ASP:OD1	2.46	0.49
2:B:97:GLU:O	2:B:101:SER:OG	2.28	0.49
1:A:343:ARG:HH12	1:A:362:LYS:HD2	1.78	0.48
2:B:10:ARG:NH2	2:B:50:THR:HG21	2.28	0.48
1:A:62:GLY:HA2	1:A:256:GLY:HA2	1.96	0.48
2:B:25:GLU:HA	2:B:28:PRO:CA	2.44	0.48
2:B:99:TYR:HD1	2:B:109:LEU:HB2	1.78	0.48
1:A:294[B]:THR:HG22	1:A:297:HIS:H	1.78	0.47
1:A:201:ARG:CG	1:A:252:ARG:HG3	2.42	0.47
1:A:333:ILE:O	1:A:337:GLU:HG3	2.14	0.47
2:B:11:ARG:HH11	2:B:15:GLN:HB3	1.79	0.47
1:A:57:TRP:CH2	1:A:59:VAL:HA	2.50	0.47
1:A:152:LYS:HB2	1:A:152:LYS:HE2	1.69	0.47

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:294[B]:THR:HG23	1:A:296:GLN:N	2.30	0.47
2:B:27:HIS:HB3	2:B:30:LYS:HB2	1.98	0.46
1:A:387:TYR:O	1:A:391:VAL:HG23	2.16	0.45
2:B:22:LEU:HA	2:B:22:LEU:HD12	1.72	0.45
1:A:189:LYS:HD3	1:A:236:ASP:HB3	1.99	0.45
1:A:59:VAL:HG21	1:A:298:VAL:HG13	1.98	0.45
2:B:21:ARG:HG3	2:B:24:ARG:HH21	1.82	0.44
2:B:56:ASP:N	2:B:56:ASP:OD1	2.50	0.43
2:B:63:LEU:HD22	2:B:95:ILE:HD13	2.00	0.43
1:A:67:ARG:HB3	1:A:305:ILE:HD12	2.00	0.43
1:A:70:LYS:HA	1:A:315:ILE:HD11	1.99	0.43
1:A:29:PHE:CE1	1:A:31:LEU:HB2	2.53	0.43
2:B:21:ARG:O	2:B:24:ARG:NH2	2.52	0.43
1:A:189:LYS:NZ	1:A:236:ASP:OD2	2.49	0.43
1:A:325:LEU:HA	1:A:326:PRO:HD2	1.88	0.43
1:A:394:VAL:O	1:A:398:MET:HG2	2.19	0.43
1:A:54:GLU:O	1:A:306:LYS:HA	2.19	0.43
1:A:104:SER:CB	1:A:107:ASN:HB2	2.49	0.42
1:A:330:SER:O	1:A:334:VAL:HG23	2.18	0.42
1:A:185:ASP:HA	1:A:186:PRO:HD3	1.77	0.42
2:B:5:LYS:HE2	2:B:5:LYS:HB2	1.86	0.42
2:B:74:ASN:O	2:B:75:ALA:HB3	2.20	0.42
1:A:63:MET:HE1	1:A:68:LEU:N	2.35	0.41
2:B:8:LYS:HE3	2:B:110:TYR:CZ	2.55	0.41
1:A:163:PHE:CZ	1:A:178:VAL:HG13	2.54	0.41
1:A:165:GLU:OE1	1:A:244:ARG:NH2	2.52	0.41
1:A:100:TYR:CD2	1:A:319:LEU:HD12	2.55	0.41
1:A:126:VAL:HG13	1:A:216:TYR:CE1	2.55	0.41
1:A:382:ILE:HD13	1:A:382:ILE:HA	1.78	0.41
1:A:394:VAL:HG22	1:A:418:LEU:HD12	2.02	0.41
2:B:62:GLU:O	2:B:66:ILE:HG13	2.21	0.41
1:A:45:ASP:OD1	1:A:45:ASP:N	2.43	0.40
1:A:95:PRO:HB2	1:A:149:PHE:HE2	1.86	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:ASP:OD2	1:A:207:SER:OG[2_455]	2.03	0.17

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	405/433 (94%)	398 (98%)	7 (2%)	0	100	100
2	B	113/124 (91%)	107 (95%)	6 (5%)	0	100	100
All	All	518/557 (93%)	505 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	366/388 (94%)	361 (99%)	5 (1%)	67	84
2	B	109/116 (94%)	102 (94%)	7 (6%)	17	31
All	All	475/504 (94%)	463 (98%)	12 (2%)	47	71

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	140	ASP
1	A	251	ASN
1	A	383	SER
1	A	394	VAL
1	A	429	ASP
2	B	21	ARG
2	B	29	THR
2	B	56	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	76	ASN
2	B	87	SER
2	B	100	GLU
2	B	118	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	76	ASN
2	B	77	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	405/433 (93%)	0.10	11 (2%) 54 56	17, 33, 74, 122	0
2	B	115/124 (92%)	2.45	60 (52%) 0 0	46, 89, 134, 147	0
All	All	520/557 (93%)	0.62	71 (13%) 3 2	17, 40, 106, 147	0

All (71) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	119	PHE	9.8
2	B	89	VAL	9.5
1	A	44	LEU	8.3
2	B	23	ILE	7.3
2	B	57	HIS	6.1
2	B	86	HIS	5.8
2	B	118	THR	5.8
2	B	90	SER	5.3
2	B	72	GLN	4.9
2	B	28	PRO	4.8
2	B	21	ARG	4.8
2	B	95	ILE	4.8
2	B	38	TYR	4.6
2	B	88	MET	4.6
2	B	22	LEU	4.5
2	B	25	GLU	4.4
2	B	47	LEU	4.4
2	B	24	ARG	4.3
2	B	93	THR	4.3
2	B	83	VAL	4.3
2	B	53	LEU	4.2
1	A	29	PHE	4.2
2	B	84	ASN	4.1
2	B	50	THR	4.0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	B	52	PHE	4.0
1	A	47	THR	3.9
2	B	48	ASP	3.9
1	A	352	LEU	3.9
2	B	73	LEU	3.8
2	B	117	GLU	3.7
2	B	71	LEU	3.7
2	B	26	GLN	3.5
2	B	39	LYS	3.5
2	B	56	ASP	3.4
2	B	33	VAL	3.4
2	B	92	SER	3.4
1	A	350	SER	3.4
2	B	29	THR	3.3
2	B	58	VAL	3.3
1	A	48	THR	3.2
2	B	114	ALA	3.2
2	B	49	LYS	3.2
2	B	116	GLN	3.0
2	B	115	SER	3.0
1	A	93	ASP	3.0
2	B	78	ALA	2.9
2	B	82	LEU	2.9
2	B	42	LYS	2.8
2	B	85	GLY	2.8
2	B	54	VAL	2.7
1	A	428	PHE	2.7
2	B	19	ASP	2.7
2	B	102	GLU	2.7
2	B	77	GLN	2.6
1	A	92	GLU	2.6
1	A	423	LEU	2.5
2	B	18	GLU	2.5
2	B	51	LYS	2.5
2	B	91	VAL	2.4
2	B	11	ARG	2.4
2	B	75	ALA	2.3
1	A	351	MET	2.3
2	B	79	PHE	2.3
2	B	27	HIS	2.2
2	B	113	TYR	2.2
2	B	100	GLU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	46	VAL	2.2
2	B	62	GLU	2.1
2	B	30	LYS	2.1
2	B	107	GLY	2.1
2	B	108	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.