



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 12, 2023 – 09:07 PM EDT

PDB ID : 4MZY  
Title : Crystal structure of enterococcus faecalis nicotinate phosphoribosyltransferase with malonate and phosphate bound  
Authors : Patskovsky, Y.; Toro, R.; Wasserman, S.R.; Almo, S.C.; Burley, S.K.; New York SGX Research Center for Structural Genomics (NYSGXRC); New York Structural Genomics Research Consortium (NYSGRC)  
Deposited on : 2013-09-30  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	494	 7% 91% 6% ..

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 4252 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NICOTINATE PHOSPHORIBOSYLTRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	486	3971	2529	666	759	17	0	10	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	448	HIS	TRP	engineered mutation	UNP Q830Y8

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0

- Molecule 4 is MALONIC ACID (three-letter code: MLA) (formula: C<sub>3</sub>H<sub>4</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 3 4	0	0
4	A	1	Total C O 7 3 4	0	0
4	A	1	Total C O 7 3 4	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	4	Total Cl 4 4	0	0

- Molecule 6 is FORMIC ACID (three-letter code: FMT) (formula: CH<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 3 1 2	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	170	Total O 171 171	0	1

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NICOTINATE PHOSPHORIBOSYLTRANSFERASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.25Å 111.38Å 154.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.78 – 1.95 25.77 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.6 (25.78-1.95) 99.6 (25.77-1.95)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.09 (at 1.95Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.154 , 0.186 0.169 , 0.196	Depositor DCC
$R_{free}$ test set	1396 reflections (3.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.1	Xtrriage
Anisotropy	0.541	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 64.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4252	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMT, MLA, GOL, CL, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/4084	0.64	1/5525 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	30	ARG	NE-CZ-NH1	5.82	123.21	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3971	0	3916	19	0
2	A	10	0	0	0	0
3	A	72	0	96	1	0
4	A	21	0	6	0	0
5	A	4	0	0	0	0
6	A	3	0	1	0	0
7	A	171	0	0	1	0
All	All	4252	0	4019	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:152:ILE:HG23	1:A:155:ASP:HB2	1.75	0.69
1:A:26:TRP:CH2	1:A:85:ASP:HB2	2.40	0.57
1:A:292:SER:O	1:A:293:ASN:CB	2.52	0.57
1:A:26:TRP:HH2	1:A:85:ASP:HB2	1.72	0.54
1:A:46:PHE:H	1:A:325:GLN:NE2	2.07	0.52
1:A:48:HIS:CD2	1:A:147:ARG:HH11	2.28	0.52
1:A:390[B]:GLN:HA	1:A:390[B]:GLN:HE21	1.74	0.52
1:A:152:ILE:CG2	1:A:155:ASP:HB2	2.41	0.49
1:A:48:HIS:HD2	1:A:147:ARG:HH11	1.62	0.47
1:A:160[A]:PHE:CD1	1:A:187:SER:HB3	2.50	0.47
1:A:59:LEU:C	1:A:59:LEU:HD12	2.35	0.47
1:A:136:ASN:HB3	7:A:681:HOH:O	2.16	0.45
1:A:132:LEU:O	1:A:136:ASN:HB2	2.17	0.44
1:A:311:ASP:OD1	3:A:514:GOL:H12	2.18	0.43
1:A:346:ASP:OD2	1:A:471:HIS:NE2	2.49	0.43
1:A:95:PHE:CZ	1:A:97:CYS:HB2	2.54	0.43
1:A:292:SER:O	1:A:293:ASN:HB3	2.17	0.43
1:A:378:GLY:HA2	1:A:460:TYR:CZ	2.54	0.42
1:A:390[B]:GLN:HE21	1:A:390[B]:GLN:CA	2.32	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	494/494 (100%)	487 (99%)	6 (1%)	1 (0%)	47 38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	293	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	430/428 (100%)	418 (97%)	12 (3%)	43 33

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	LEU
1	A	59	LEU
1	A	85	ASP
1	A	87	LEU
1	A	131	LEU
1	A	133	ASN
1	A	239	LYS
1	A	268	MET
1	A	332	PHE
1	A	353	ASN
1	A	448	HIS
1	A	485	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	ASN
1	A	48	HIS
1	A	325	GLN
1	A	456	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 4 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	FMT	A	520	-	2,2,2	0.70	0	1,1,1	0.76	0
3	GOL	A	506	-	5,5,5	0.23	0	5,5,5	0.31	0
3	GOL	A	505	-	5,5,5	0.29	0	5,5,5	0.29	0
3	GOL	A	508	-	5,5,5	0.28	0	5,5,5	1.08	0
3	GOL	A	510	-	5,5,5	0.17	0	5,5,5	0.59	0
3	GOL	A	504	-	5,5,5	0.37	0	5,5,5	0.17	0
2	PO4	A	501	-	4,4,4	0.99	0	6,6,6	0.59	0
3	GOL	A	503	-	5,5,5	0.46	0	5,5,5	0.41	0
3	GOL	A	507	-	5,5,5	0.33	0	5,5,5	0.31	0
2	PO4	A	502	-	4,4,4	0.94	0	6,6,6	0.42	0
3	GOL	A	512	-	5,5,5	0.29	0	5,5,5	0.57	0
3	GOL	A	513	-	5,5,5	0.42	0	5,5,5	0.25	0
4	MLA	A	515	-	6,6,6	1.13	0	7,7,7	1.18	0
4	MLA	A	516	-	6,6,6	1.21	0	7,7,7	1.24	0
3	GOL	A	509	-	5,5,5	0.26	0	5,5,5	0.28	0
4	MLA	A	517	-	6,6,6	1.14	0	7,7,7	1.12	0
3	GOL	A	511	-	5,5,5	0.18	0	5,5,5	0.25	0
3	GOL	A	514	-	5,5,5	0.27	0	5,5,5	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	506	-	-	2/4/4/4	-
3	GOL	A	505	-	-	1/4/4/4	-
3	GOL	A	508	-	-	2/4/4/4	-
3	GOL	A	510	-	-	2/4/4/4	-
3	GOL	A	504	-	-	0/4/4/4	-
3	GOL	A	503	-	-	0/4/4/4	-
3	GOL	A	507	-	-	2/4/4/4	-
4	MLA	A	515	-	-	0/4/4/4	-
3	GOL	A	512	-	-	2/4/4/4	-
3	GOL	A	513	-	-	2/4/4/4	-
4	MLA	A	516	-	-	2/4/4/4	-
4	MLA	A	517	-	-	2/4/4/4	-
3	GOL	A	509	-	-	2/4/4/4	-
3	GOL	A	511	-	-	2/4/4/4	-
3	GOL	A	514	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	506	GOL	C1-C2-C3-O3
3	A	507	GOL	O1-C1-C2-C3
3	A	508	GOL	C1-C2-C3-O3
3	A	509	GOL	C1-C2-C3-O3
3	A	510	GOL	C1-C2-C3-O3
3	A	511	GOL	C1-C2-C3-O3
3	A	511	GOL	O2-C2-C3-O3
3	A	512	GOL	C1-C2-C3-O3
3	A	512	GOL	O2-C2-C3-O3
3	A	513	GOL	C1-C2-C3-O3
3	A	514	GOL	O1-C1-C2-C3
3	A	506	GOL	O2-C2-C3-O3
3	A	513	GOL	O2-C2-C3-O3
3	A	514	GOL	C1-C2-C3-O3
3	A	507	GOL	O1-C1-C2-O2
3	A	510	GOL	O2-C2-C3-O3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	A	514	GOL	O1-C1-C2-O2
3	A	514	GOL	O2-C2-C3-O3
3	A	508	GOL	O2-C2-C3-O3
3	A	509	GOL	O2-C2-C3-O3
3	A	505	GOL	O1-C1-C2-C3
4	A	516	MLA	C1-C2-C3-O3A
4	A	516	MLA	C1-C2-C3-O3B
4	A	517	MLA	C1-C2-C3-O3A
4	A	517	MLA	C1-C2-C3-O3B

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	514	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	486/494 (98%)	0.19	35 (7%) <b>15</b> <b>23</b>	19, 30, 64, 119	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	488	GLU	7.6
1	A	353	ASN	7.2
1	A	485	HIS	6.3
1	A	486	LEU	6.0
1	A	3	TYR	4.7
1	A	487	THR	4.4
1	A	352	SER	4.2
1	A	340	GLU	4.2
1	A	354	ALA	4.1
1	A	341	ASP	4.1
1	A	343	GLN	3.9
1	A	373	ASP	3.5
1	A	481[A]	LYS	3.5
1	A	482	ASP	3.3
1	A	484	LYS	3.2
1	A	375	LYS	3.0
1	A	342	GLY	3.0
1	A	477	GLU	2.9
1	A	356	LYS	2.9
1	A	355	GLU	2.8
1	A	392	GLU	2.8
1	A	293	ASN	2.6
1	A	294	ASP	2.6
1	A	407	TYR	2.5
1	A	70	GLU	2.4
1	A	409	ARG	2.4
1	A	281	GLU	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	160[A]	PHE	2.2
1	A	339	GLY	2.2
1	A	445	ASP	2.2
1	A	208	VAL	2.2
1	A	17	TYR	2.1
1	A	438	GLN	2.1
1	A	154	ASP	2.0
1	A	371	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	MLA	A	515	7/7	0.70	0.20	41,56,76,109	0
3	GOL	A	513	6/6	0.74	0.31	40,70,76,81	0
3	GOL	A	514	6/6	0.83	0.25	58,68,96,101	0
3	GOL	A	512	6/6	0.85	0.23	43,76,92,96	0
3	GOL	A	509	6/6	0.86	0.18	56,60,73,81	0
3	GOL	A	507	6/6	0.88	0.12	50,71,80,84	0
3	GOL	A	506	6/6	0.88	0.14	45,62,83,97	0
5	CL	A	522	1/1	0.88	0.09	72,72,72,72	0
3	GOL	A	510	6/6	0.90	0.20	52,75,85,95	0
4	MLA	A	517	7/7	0.90	0.15	45,57,63,80	0
5	CL	A	519	1/1	0.90	0.09	64,64,64,64	0
3	GOL	A	508	6/6	0.90	0.16	49,55,61,85	0
6	FMT	A	520	3/3	0.93	0.28	43,43,56,73	0
3	GOL	A	504	6/6	0.94	0.17	29,38,40,62	0
3	GOL	A	505	6/6	0.94	0.15	32,46,66,73	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PO4	A	502	5/5	0.94	0.18	25,28,47,54	5
5	CL	A	521	1/1	0.95	0.15	63,63,63,63	0
3	GOL	A	511	6/6	0.95	0.22	44,57,68,79	0
4	MLA	A	516	7/7	0.95	0.13	41,56,79,88	0
3	GOL	A	503	6/6	0.96	0.09	26,35,36,42	0
2	PO4	A	501	5/5	0.97	0.13	44,46,62,93	0
5	CL	A	518	1/1	0.97	0.06	38,38,38,38	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.