



wwPDB NMR Structure Validation Summary Report ⓘ

Jun 4, 2023 – 10:47 AM EDT

PDB ID : 2NB1
BMRB ID : 25958
Title : P63/p73 hetero-tetramerisation domain
Authors : Gebel, J.; Buchner, L.; Loehr, F.M.; Luh, L.M.; Coutandin, D.; Guentert, P.;
Doetsch, V.
Deposited on : 2016-01-19

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.33


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 86%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	60	
1	C	60	
2	B	50	
2	D	50	

2 Ensemble composition and analysis

This entry contains 20 models. Model 3 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:4-A:32, B:98-B:140, C:1004-C:1032, D:1098- D:1140 (144)	0.84	3
2	A:35-A:46 (12)	0.30	17
3	C:1035-C:1046 (12)	0.32	3

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 5, 6, 7, 8, 9, 11, 14, 15, 16, 17
2	3, 4, 10, 13, 19, 20
3	12, 18

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 3732 atoms, of which 1862 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Tumor protein 63.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	60	1016	320	503	89	102	2	0
1	C	60	1016	320	503	89	102	2	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP Q9H3D4
A	21	GLU	LYS	engineered mutation	UNP Q9H3D4
C	1001	SER	-	expression tag	UNP Q9H3D4
C	1021	GLU	LYS	engineered mutation	UNP Q9H3D4

- Molecule 2 is a protein called Tumor protein p73.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
2	B	50	850	266	428	73	81	2	0
2	D	50	850	266	428	73	81	2	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	93	GLY	-	expression tag	UNP O15350
B	94	SER	-	expression tag	UNP O15350
B	107	LYS	GLU	engineered mutation	UNP O15350
D	1093	GLY	-	expression tag	UNP O15350
D	1094	SER	-	expression tag	UNP O15350
D	1107	LYS	GLU	engineered mutation	UNP O15350

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Tumor protein 63




- Molecule 1: Tumor protein 63



- Molecule 2: Tumor protein p73



- Molecule 2: Tumor protein p73



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 3. Colouring as in section 4.1 above.

- Molecule 1: Tumor protein 63

Chain A:  53% 15% 32%



- Molecule 1: Tumor protein 63

Chain C:  57% 12% 32%



- Molecule 2: Tumor protein p73

Chain B:  64% 22% 14%



- Molecule 2: Tumor protein p73

Chain D:  60% 26% 14%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	3.97
OPALp	refinement	1.4

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	2706
Number of shifts mapped to atoms	2706
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	86%

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.64±0.01	0±0/364 (0.0± 0.0%)	0.95±0.04	0±0/492 (0.0± 0.1%)
1	C	0.64±0.01	0±0/364 (0.0± 0.0%)	0.96±0.04	0±1/492 (0.1± 0.1%)
2	B	0.63±0.01	0±0/373 (0.0± 0.0%)	0.99±0.04	0±0/502 (0.1± 0.1%)
2	D	0.63±0.01	0±0/373 (0.0± 0.0%)	0.98±0.05	0±1/502 (0.1± 0.1%)
All	All	0.64	0/29480 (0.0%)	0.97	26/39760 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.7±0.7
1	C	0.0±0.0	0.5±0.6
2	B	0.0±0.0	0.5±0.6
2	D	0.0±0.0	0.6±0.7
All	All	0	45

There are no bond-length outliers.

5 of 16 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	104	ARG	NE-CZ-NH2	-7.33	116.63	120.30	12	3
2	D	1104	ARG	NE-CZ-NH2	-7.27	116.67	120.30	4	4
2	B	106	ARG	NE-CZ-NH2	-6.52	117.04	120.30	6	1
2	D	1125	VAL	CA-CB-CG1	6.03	119.95	110.90	13	2
1	C	1011	ARG	NE-CZ-NH2	-5.66	117.47	120.30	12	1

There are no chirality outliers.

5 of 19 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	11	ARG	Sidechain	5
2	B	99	TYR	Sidechain	4
2	D	1099	TYR	Sidechain	4
2	B	106	ARG	Sidechain	3
1	C	1013	ARG	Sidechain	3

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	358	358	358	0±0
1	C	358	358	358	0±0
2	D	367	385	385	0±0
2	B	367	385	385	0±0
All	All	29000	29720	29720	11

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

5 of 11 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:119:LEU:HD12	2:D:1108:ASN:HD21	0.49	1.67	14	1
1:C:1037:ILE:H	1:C:1037:ILE:HD12	0.47	1.70	5	1
2:B:101:LEU:HD22	2:D:1116:LYS:HE2	0.46	1.87	6	1
1:C:1028:LEU:HD22	2:D:1130:VAL:HG23	0.46	1.86	13	1
1:C:1036:THR:HG23	2:D:1125:VAL:HG21	0.46	1.87	10	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	41/60 (68%)	39±1 (95±2%)	2±1 (5±2%)	0±0 (0±1%)	54	85
1	C	41/60 (68%)	39±1 (96±2%)	2±1 (4±2%)	0±0 (0±1%)	54	85
2	B	43/50 (86%)	40±1 (93±2%)	2±1 (5±2%)	1±1 (2±2%)	10	50
2	D	43/50 (86%)	40±1 (93±2%)	2±1 (5±2%)	1±1 (2±2%)	11	52
All	All	3360/4400 (76%)	3166 (94%)	157 (5%)	37 (1%)	18	66

5 of 8 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	128	PRO	13
2	D	1128	PRO	12
2	B	126	PRO	4
2	D	1126	PRO	4
2	B	127	GLN	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	40/59 (68%)	34±2 (86±6%)	6±2 (14±6%)	7	47
1	C	40/59 (68%)	34±3 (86±6%)	6±3 (14±6%)	6	45
2	B	42/48 (88%)	36±2 (86±5%)	6±2 (14±5%)	6	46
2	D	42/48 (88%)	35±2 (84±5%)	7±2 (16±5%)	5	42
All	All	3280/4280 (77%)	2799 (85%)	481 (15%)	6	45

5 of 128 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	D	1104	ARG	14
2	D	1134	ARG	11
2	B	102	GLN	10
1	C	1020	LEU	10
2	D	1137	GLN	10

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 86% for the well-defined parts and 83% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	2706
Number of shifts mapped to atoms	2706
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	10

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	214	-0.52 ± 0.04	Should be checked
$^{13}\text{C}_\beta$	212	0.09 ± 0.08	None needed (< 0.5 ppm)
$^{13}\text{C}'$	210	-0.17 ± 0.10	None needed (< 0.5 ppm)
^{15}N	202	-0.49 ± 0.21	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 86%, i.e. 2214 atoms were assigned a chemical shift out of a possible 2582. 0 out of 42 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	832/832 (100%)	334/334 (100%)	336/336 (100%)	162/162 (100%)
Sidechain	1280/1590 (81%)	866/1028 (84%)	398/492 (81%)	16/70 (23%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	102/160 (64%)	68/74 (92%)	34/84 (40%)	0/2 (0%)
Overall	2214/2582 (86%)	1268/1436 (88%)	768/912 (84%)	178/234 (76%)

7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	41	ARG	NE	136.90	76.53 – 92.65	32.5
1	C	1041	ARG	NE	136.90	76.53 – 92.65	32.5
1	B	104	ARG	NE	136.28	76.53 – 92.65	32.1
1	D	1104	ARG	NE	136.28	76.53 – 92.65	32.1
1	B	134	ARG	NE	135.12	76.53 – 92.65	31.4
1	D	1134	ARG	NE	135.12	76.53 – 92.65	31.4
1	A	11	ARG	NE	134.86	76.53 – 92.65	31.2
1	C	1011	ARG	NE	134.86	76.53 – 92.65	31.2
1	A	13	ARG	NE	134.05	76.53 – 92.65	30.7
1	C	1013	ARG	NE	134.05	76.53 – 92.65	30.7


7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.


Random coil index (RCI) for chain A:




Random coil index (RCI) for chain B:



Random coil index (RCI) for chain C:



Random coil index (RCI) for chain D:



8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	3020
Intra-residue ($ i-j =0$)	412
Sequential ($ i-j =1$)	706
Medium range ($ i-j >1$ and $ i-j <5$)	614
Long range ($ i-j \geq 5$)	52
Inter-chain	796
Hydrogen bond restraints	440
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	13.7
Number of long range restraints per residue ¹	0.2

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	38.0	0.2
0.2-0.5 (Medium)	56.3	0.5
>0.5 (Large)	13.8	0.58

8.2.2 Average number of dihedral-angle violations per model

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations

9 Distance violation analysis [i](#)


9.1 Summary of distance violations [i](#)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue ($i-j =0$)	412	13.6	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	376	12.5	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	36	1.2	0	0.0	0.0	0	0.0	0.0
Sequential ($i-j =1$)	706	23.4	5	0.7	0.2	0	0.0	0.0
Backbone-Backbone	160	5.3	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	464	15.4	1	0.2	0.0	0	0.0	0.0
Sidechain-Sidechain	82	2.7	4	4.9	0.1	0	0.0	0.0
Medium range ($i-j >1$ & $i-j <5$)	614	20.3	17	2.8	0.6	0	0.0	0.0
Backbone-Backbone	172	5.7	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	264	8.7	3	1.1	0.1	0	0.0	0.0
Sidechain-Sidechain	178	5.9	14	7.9	0.5	0	0.0	0.0
Long range ($i-j \geq 5$)	52	1.7	11	21.2	0.4	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	18	0.6	2	11.1	0.1	0	0.0	0.0
Sidechain-Sidechain	34	1.1	9	26.5	0.3	0	0.0	0.0
Inter-chain	796	26.4	66	8.3	2.2	0	0.0	0.0
Backbone-Backbone	34	1.1	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	317	10.5	26	8.2	0.9	0	0.0	0.0
Sidechain-Sidechain	445	14.7	40	9.0	1.3	0	0.0	0.0
Hydrogen bond	440	14.6	110	25.0	3.6	51	11.6	1.7
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	3020	100.0	209	6.9	6.9	51	1.7	1.7
Backbone-Backbone	806	26.7	110	13.6	3.6	51	6.3	1.7
Backbone-Sidechain	1439	47.6	32	2.2	1.1	0	0.0	0.0
Sidechain-Sidechain	775	25.7	67	8.6	2.2	0	0.0	0.0

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfid bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	0	2	83	0	22	107	0.28	0.57	0.15	0.24
2	0	0	88	1	18	107	0.32	0.57	0.15	0.3
3	0	0	90	1	15	106	0.29	0.58	0.13	0.29
4	0	0	84	1	18	103	0.29	0.58	0.13	0.27
5	0	0	88	0	17	105	0.3	0.57	0.15	0.28
6	0	0	89	1	20	110	0.28	0.57	0.14	0.25
7	0	0	93	1	23	117	0.29	0.57	0.15	0.26
8	0	1	88	2	19	110	0.28	0.58	0.13	0.25
9	0	1	89	0	22	112	0.3	0.57	0.15	0.25
10	0	0	86	1	19	106	0.32	0.56	0.15	0.32
11	0	0	90	0	16	106	0.33	0.58	0.15	0.3


Continued on next page...

Continued from previous page...

Model ID	Number of violations					Total	Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵					
12	0	0	92	1	16	109	0.3	0.58	0.15	0.28
13	0	0	87	0	18	105	0.29	0.58	0.15	0.24
14	0	0	85	1	20	106	0.31	0.58	0.16	0.3
15	0	0	89	0	18	107	0.3	0.58	0.14	0.28
16	0	0	90	0	17	107	0.32	0.58	0.15	0.29
17	0	2	94	3	16	115	0.3	0.58	0.14	0.27
18	0	0	88	1	19	108	0.3	0.58	0.14	0.29
19	0	0	84	0	22	106	0.29	0.58	0.15	0.25
20	0	0	89	0	22	111	0.28	0.58	0.14	0.27

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model [\(i\)](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right


9.3 Distance violation statistics for the ensemble

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 2481(IR:412, SQ:701, MR:597, LR:41, IC:730) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
0	4	12	8	39	63	1	5.0
0	1	3	3	11	18	2	10.0
0	0	0	0	8	8	3	15.0
0	0	0	0	2	2	4	20.0
0	0	0	0	2	2	5	25.0
0	0	0	0	0	0	6	30.0
0	0	0	0	2	2	7	35.0
0	0	1	0	1	2	8	40.0
0	0	1	0	0	1	9	45.0
0	0	0	0	0	0	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	0	1	1	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
0	0	0	0	0	0	16	80.0
0	0	0	0	0	0	17	85.0
0	0	0	0	0	0	18	90.0
0	0	0	0	0	0	19	95.0
0	0	0	0	0	0	20	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶ Number of models with violations


9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)



9.4 Most violated distance restraints in the ensemble [i](#)

9.4.1 Histogram : Distribution of mean distance violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(4,95)	2:D:1106:ARG:O	2:D:1110:GLU:N	20	0.56	0.02	0.56
(4,71)	2:B:106:ARG:O	2:B:110:GLU:N	20	0.55	0.04	0.56
(4,87)	2:D:1128:PRO:O	2:D:1132:SER:N	20	0.52	0.06	0.55
(4,63)	2:B:128:PRO:O	2:B:132:SER:N	20	0.51	0.09	0.56
(4,70)	2:B:135:GLN:O	2:B:139:LEU:N	20	0.49	0.05	0.5
(4,19)	1:A:22:ILE:O	1:A:26:LEU:N	20	0.47	0.11	0.53
(4,10)	1:A:13:ARG:O	1:A:17:GLU:N	20	0.46	0.1	0.48
(4,94)	2:D:1135:GLN:O	2:D:1139:LEU:N	20	0.46	0.08	0.49
(4,99)	2:D:1110:GLU:O	2:D:1114:LYS:N	20	0.43	0.12	0.45
(4,75)	2:B:110:GLU:O	2:B:114:LYS:N	20	0.43	0.13	0.46

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(4,99)	2:D:1110:GLU:O	2:D:1114:LYS:N	15	0.58
(4,95)	2:D:1106:ARG:O	2:D:1110:GLU:N	4	0.58
(4,95)	2:D:1106:ARG:O	2:D:1110:GLU:N	8	0.58
(4,95)	2:D:1106:ARG:O	2:D:1110:GLU:N	13	0.58
(4,95)	2:D:1106:ARG:O	2:D:1110:GLU:N	14	0.58
(4,95)	2:D:1106:ARG:O	2:D:1110:GLU:N	15	0.58
(4,95)	2:D:1106:ARG:O	2:D:1110:GLU:N	20	0.58
(4,9)	1:A:42:GLN:O	1:A:46:GLN:N	14	0.58
(4,87)	2:D:1128:PRO:O	2:D:1132:SER:N	19	0.58
(4,75)	2:B:110:GLU:O	2:B:114:LYS:N	18	0.58

10 Dihedral-angle violation analysis

Dihedral angle analysis failed due to data error in the dihedral angle restraints, possibly missing target value