



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 09:23 pm BST

PDB ID : 1NES  
Title : STRUCTURE OF THE PRODUCT COMPLEX OF ACETYL-ALA-PRO-ALA WITH PORCINE PANCREATIC ELASTASE AT 1.65 ANGSTROMS RESOLUTION  
Authors : Meyer Junior, E.F.; Radhakrishnan, R.; M Cole, G.; Presta, L.G.  
Deposited on : 1995-07-31  
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	E	240	
2	I	4	
2	J	4	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 1998 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ELASTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	E	240	1812	1133	324	345	10	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	77	ASN	ASP	CONFLICT	UNP P00772

- Molecule 2 is a protein called ACETYL-ALA-PRO-ALA.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	I	4	21	13	3	5	0	0	0
2	J	4	21	13	3	5	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	1	Total	Ca	0	0
			1	1		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	E	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	137	Total	O	0	0
			137	137		
5	J	1	Total	O	0	0
			1	1		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ELASTASE

Chain E: 



- Molecule 2: ACETYL-ALA-PRO-ALA

Chain I: 



- Molecule 2: ACETYL-ALA-PRO-ALA

Chain J: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.40Å 58.20Å 75.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.00 – 1.65 7.05 – 1.66	Depositor EDS
% Data completeness (in resolution range)	72.0 (7.00-1.65) 76.1 (7.05-1.66)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtrriage
Refinement program	EREF	Depositor
R, $R_{free}$	0.184 , (Not available) 0.281 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.9	Xtrriage
Anisotropy	0.134	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 66.5	EDS
L-test for twinning <sup>1</sup>	$\langle  L  \rangle = 0.39$ , $\langle L^2 \rangle = 0.22$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	1998	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.57% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CA, SO4, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	E	1.10	4/1852 (0.2%)	1.62	22/2531 (0.9%)
2	I	0.91	0/19	1.38	0/25
2	J	0.75	0/19	2.34	1/25 (4.0%)
All	All	1.10	4/1890 (0.2%)	1.63	23/2581 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	25

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	223	ARG	NE-CZ	6.27	1.41	1.33
1	E	107	ARG	NE-CZ	6.01	1.40	1.33
1	E	189	SER	CB-OG	5.35	1.49	1.42
1	E	36	ARG	CZ-NH1	5.13	1.39	1.33

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	188(A)	ARG	NE-CZ-NH2	-19.11	110.75	120.30
1	E	188(A)	ARG	NE-CZ-NH1	12.74	126.67	120.30
1	E	80	GLU	OE1-CD-OE2	-8.91	112.61	123.30
1	E	107	ARG	CD-NE-CZ	8.28	135.20	123.60
1	E	145	ARG	NE-CZ-NH1	8.04	124.32	120.30
2	J	606	ALA	N-CA-CB	-8.00	98.90	110.10

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	107	ARG	NE-CZ-NH1	7.71	124.16	120.30
1	E	217(A)	ARG	NE-CZ-NH2	-7.52	116.54	120.30
1	E	70	GLU	OE1-CD-OE2	-7.28	114.57	123.30
1	E	48	ARG	NE-CZ-NH2	-7.10	116.75	120.30
1	E	223	ARG	NE-CZ-NH1	6.77	123.69	120.30
1	E	164	ASP	CB-CG-OD1	6.74	124.36	118.30
1	E	24	ARG	CG-CD-NE	-6.71	97.71	111.80
1	E	145	ARG	NE-CZ-NH2	-6.35	117.13	120.30
1	E	137	TYR	CB-CG-CD2	-6.22	117.27	121.00
1	E	24	ARG	NE-CZ-NH2	-6.15	117.23	120.30
1	E	206	GLN	N-CA-CB	-6.07	99.67	110.60
1	E	24	ARG	CB-CA-C	-6.07	98.26	110.40
1	E	65(A)	ARG	NE-CZ-NH2	5.99	123.29	120.30
1	E	207	TYR	CB-CG-CD2	-5.75	117.55	121.00
1	E	21	GLU	OE1-CD-OE2	-5.63	116.54	123.30
1	E	97	ASP	CB-CG-OD1	5.54	123.28	118.30
1	E	36	ARG	NE-CZ-NH2	-5.21	117.70	120.30

There are no chirality outliers.

All (25) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	127	GLY	Mainchain
1	E	132	ASN	Sidechain
1	E	142	GLY	Mainchain
1	E	148	ASN	Sidechain
1	E	153	GLN	Sidechain
1	E	159	TYR	Sidechain
1	E	188(A)	ARG	Sidechain
1	E	189	SER	Mainchain
1	E	204	ASN	Sidechain
1	E	206	GLN	Sidechain
1	E	21	GLU	Sidechain
1	E	213	THR	Mainchain
1	E	219	GLY	Mainchain
1	E	239	ASN	Sidechain
1	E	245	ASN	Sidechain
1	E	25	ASN	Sidechain
1	E	30	GLN	Sidechain
1	E	36(A)	SER	Mainchain
1	E	49	GLN	Sidechain
1	E	60	ASP	Sidechain

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	E	70	GLU	Sidechain
1	E	80	GLU	Sidechain
1	E	86	GLN	Sidechain
1	E	93	TYR	Sidechain
1	E	95	ASN	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1812	0	1739	16	1
2	I	21	0	20	1	0
2	J	21	0	20	4	0
3	E	1	0	0	0	0
4	E	5	0	0	0	0
5	E	137	0	0	1	0
5	J	1	0	0	0	0
All	All	1998	0	1779	20	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:E:447:HOH:O	2:J:607:PRO:HD3	1.82	0.79
2:J:606:ALA:HB1	2:J:607:PRO:HD2	1.71	0.70
1:E:83:VAL:HG13	1:E:110:GLN:HG2	1.75	0.67
1:E:48:ARG:HB2	1:E:51:TRP:HB2	1.80	0.62
1:E:91:HIS:HD2	1:E:93:TYR:H	1.51	0.57
1:E:24:ARG:HH12	1:E:77:ASN:HD22	1.55	0.55
1:E:68:VAL:HG22	1:E:81:GLN:HB2	1.89	0.54
1:E:83:VAL:CG1	1:E:110:GLN:HG2	2.38	0.54
1:E:24:ARG:HH12	1:E:77:ASN:ND2	2.09	0.51
1:E:50:ASN:H	1:E:50:ASN:HD22	1.61	0.48
1:E:23:GLN:HB2	1:E:26:SER:OG	2.14	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:606:ALA:HB1	2:J:607:PRO:CD	2.42	0.47
1:E:60:ASP:OD2	1:E:96:THR:HG23	2.16	0.46
1:E:105:LEU:HD12	1:E:241:VAL:HG11	1.97	0.46
1:E:176:VAL:HG21	1:E:182:CYS:SG	2.57	0.44
1:E:195:SER:OG	2:I:600:ACE:H2	2.18	0.43
1:E:213:THR:HG1	1:E:228:PHE:HE2	1.65	0.42
2:J:606:ALA:CB	2:J:607:PRO:HD2	2.44	0.42
1:E:31:ILE:HG22	1:E:68:VAL:HG12	2.00	0.42
1:E:237:TRP:O	1:E:241:VAL:HG23	2.20	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:62:GLU:O	1:E:175:THR:OG1[4_556]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	238/240 (99%)	230 (97%)	8 (3%)	0	100	100
2	I	2/4 (50%)	2 (100%)	0	0	100	100
2	J	2/4 (50%)	1 (50%)	0	1 (50%)	0	0
All	All	242/248 (98%)	233 (96%)	8 (3%)	1 (0%)	34	16

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	J	606	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	194/198 (98%)	188 (97%)	6 (3%)	40	14
2	I	1/1 (100%)	1 (100%)	0	100	100
2	J	1/1 (100%)	1 (100%)	0	100	100
All	All	196/200 (98%)	190 (97%)	6 (3%)	40	14

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	31	ILE
1	E	48	ARG
1	E	50	ASN
1	E	83	VAL
1	E	86	GLN
1	E	110	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	23	GLN
1	E	50	ASN
1	E	75	GLN
1	E	91	HIS
1	E	204	ASN
1	E	239	ASN
1	E	240	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	SO4	E	510	-	4,4,4	0.13	0	6,6,6	0.40	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.