



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 10:45 am BST

PDB ID : 4OC8  
Title : DNA modification-dependent restriction endonuclease AspBHI  
Authors : Horton, J.R.  
Deposited on : 2014-01-08  
Resolution : 2.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2691 (2.90-2.86)
Clashscore	141614	2947 (2.90-2.86)
Ramachandran outliers	138981	2868 (2.90-2.86)
Sidechain outliers	138945	2871 (2.90-2.86)
RSRZ outliers	127900	2629 (2.90-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	<p>0% 87% 12%</p>
1	B	388	<p>2% 88% 12%</p>
1	C	388	<p>3% 48% 7% 45%</p>
1	D	388	<p>3% 47% 8% 45%</p>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 9516 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called restriction endonuclease AspBHI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	387	Total 3023	C 1919	N 549	O 551	S 4	0	1	0
1	B	387	Total 3023	C 1919	N 549	O 551	S 4	0	1	0
1	C	215	Total 1691	C 1085	N 301	O 303	S 2	0	1	0
1	D	215	Total 1686	C 1082	N 301	O 301	S 2	0	0	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	Total 5	O 4	P 1	0	0
2	A	1	Total 5	O 4	P 1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	22	Total	O	0	0
			22	22		
3	B	20	Total	O	0	0
			20	20		
3	C	6	Total	O	0	0
			6	6		
3	D	5	Total	O	0	0
			5	5		



ILE  
THR  
ALA  
SER  
VAL

● Molecule 1: restriction endonuclease AspBHI



ILE  
THR  
ALA  
SER  
VAL

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	195.09 Å 195.09 Å 81.45 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.74 – 2.88 30.74 – 2.88	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.74-2.88) 93.7 (30.74-2.88)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.40 (at 2.90 Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, $R_{free}$	0.197 , 0.235 0.199 , 0.236	Depositor DCC
$R_{free}$ test set	2013 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	60.3	Xtrriage
Anisotropy	0.333	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 31.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.027 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	9516	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/3096	0.48	1/4218 (0.0%)
1	B	0.28	0/3096	0.49	0/4218
1	C	0.25	0/1742	0.46	0/2379
1	D	0.25	0/1734	0.46	0/2368
All	All	0.27	0/9668	0.48	1/13183 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	267	VAL	N-CA-C	-5.86	95.17	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	266	ASP	Peptide



## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3023	0	3024	30	0
1	B	3023	0	3024	27	0
1	C	1691	0	1666	15	0
1	D	1686	0	1662	19	0
2	A	15	0	0	0	0
2	B	10	0	0	0	0
2	C	5	0	0	0	0
2	D	10	0	0	0	0
3	A	22	0	0	0	0
3	B	20	0	0	0	0
3	C	6	0	0	0	0
3	D	5	0	0	0	0
All	All	9516	0	9376	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:155:ALA:HB3	1:B:155:ALA:HB3	1.69	0.74
1:A:266:ASP:HB2	1:A:287:LEU:HD12	1.75	0.69
1:B:25:THR:HG22	1:B:26:GLU:HG2	1.77	0.67
1:A:35:VAL:HG21	1:A:42:ARG:HG2	1.78	0.66
1:A:48:TYR:OH	1:A:52[B]:ARG:NE	2.31	0.64
1:D:2:THR:HG23	1:D:18:ALA:HA	1.80	0.63
1:A:267:VAL:HG13	1:B:146:ALA:HB2	1.81	0.61
1:C:140:ALA:HB2	1:C:174:ILE:HD13	1.83	0.61
1:D:103:LEU:O	1:D:107:ILE:HG12	2.01	0.59
1:D:61:THR:HG22	1:D:125:PHE:HB2	1.85	0.58
1:C:115:GLN:HB3	1:C:118:LEU:HD12	1.85	0.58
1:A:386:ALA:O	1:B:211:ARG:NH2	2.39	0.56
1:D:140:ALA:HB2	1:D:174:ILE:HD13	1.87	0.56
1:A:253:GLU:OE1	1:A:281:ARG:NH1	2.40	0.54
1:D:43:GLN:O	1:D:98:ARG:NH1	2.34	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:298:VAL:HB	1:A:329:GLU:HG3	1.90	0.53
1:A:337:SER:OG	1:A:338:TYR:N	2.42	0.53
1:C:32:ASP:OD2	1:C:47:ARG:NH1	2.32	0.53
1:B:27:ARG:HG3	1:B:29:GLY:H	1.74	0.52
1:B:24:LYS:HE2	1:B:28:GLY:HA2	1.91	0.52
1:D:107:ILE:HD12	1:D:120:PRO:HD2	1.90	0.52
1:C:93:LEU:HB2	1:C:166:GLN:HB3	1.91	0.51
1:D:52:ARG:HD2	1:D:52:ARG:O	2.11	0.50
1:A:267:VAL:O	1:A:269:ARG:NH1	2.45	0.50
1:B:27:ARG:CZ	1:B:29:GLY:HA3	2.42	0.50
1:B:321:LEU:O	1:B:325:ILE:HG13	2.11	0.50
1:C:176:ASP:HA	1:C:213:LEU:HB2	1.94	0.50
1:A:281:ARG:NH2	1:C:151:GLU:OE2	2.45	0.49
1:C:125:PHE:HB3	1:C:134:PHE:HB3	1.94	0.49
1:A:67:PRO:O	1:A:372:ARG:NH2	2.45	0.49
1:B:32:ASP:OD1	1:B:132:ARG:NE	2.37	0.49
1:C:103:LEU:O	1:C:107:ILE:HG12	2.13	0.49
1:A:312:ALA:HB2	1:A:338:TYR:CZ	2.48	0.48
1:C:91:ARG:NH1	1:C:95:ASP:OD2	2.48	0.47
1:A:75:GLU:OE1	1:A:217:ARG:NH2	2.48	0.47
1:A:367:ARG:HD2	1:A:372:ARG:HH21	1.79	0.47
1:A:25:THR:OG1	1:A:26:GLU:OE1	2.33	0.47
1:D:115:GLN:HB3	1:D:118:LEU:HD12	1.96	0.47
1:B:215:ALA:HB2	1:C:157:TRP:HB3	1.96	0.46
1:B:281:ARG:NH2	1:D:151:GLU:OE1	2.48	0.46
1:A:140:ALA:HB2	1:A:174:ILE:HD13	1.97	0.46
1:C:208:GLY:N	1:C:209:GLY:HA3	2.30	0.45
1:B:303:GLU:HB2	1:B:321:LEU:HD13	1.97	0.45
1:D:38:VAL:HG21	1:D:59:VAL:HG21	1.99	0.45
1:B:380:TRP:CZ2	1:B:384:ILE:HD11	2.51	0.45
1:A:381:LEU:HD23	1:A:384:ILE:HD12	1.99	0.45
1:B:337:SER:OG	1:B:338:TYR:N	2.49	0.45
1:D:52:ARG:C	1:D:52:ARG:HD2	2.37	0.45
1:A:109:ASP:OD1	1:B:164:ARG:HD3	2.18	0.44
1:D:105:ARG:NH1	1:D:109:ASP:OD1	2.50	0.44
1:A:128:GLU:HB2	1:A:133:THR:OG1	2.17	0.44
1:A:25:THR:HG23	1:A:26:GLU:H	1.82	0.44
1:C:113:LEU:HB2	1:C:115:GLN:HG3	2.00	0.44
1:A:91:ARG:NH1	1:A:95:ASP:O	2.50	0.43
1:C:43:GLN:CG	1:C:44:GLY:H	2.31	0.43
1:D:2:THR:N	1:D:19:VAL:H	2.16	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:286:ARG:HD3	1:B:297:GLU:OE1	2.19	0.42
1:A:264:LEU:HD23	1:A:264:LEU:HA	1.73	0.42
1:B:265:PRO:HD2	1:B:287:LEU:HD11	2.00	0.42
1:A:146:ALA:CB	1:B:267:VAL:HG22	2.49	0.42
1:B:283:GLY:O	1:B:324:ARG:NH2	2.52	0.42
1:B:313:VAL:HG11	1:B:333:LEU:HD21	2.00	0.42
1:C:107:ILE:HD12	1:C:120:PRO:HD2	2.01	0.42
1:A:350:ASP:HB2	1:A:352:HIS:CD2	2.55	0.42
1:B:32:ASP:OD2	1:B:34:LEU:HB2	2.20	0.42
1:A:380:TRP:CZ2	1:A:384:ILE:HD11	2.54	0.42
1:B:285:GLY:HA3	1:B:300:PHE:CZ	2.55	0.42
1:C:89:PRO:HA	1:C:165:PHE:HB3	2.00	0.42
1:D:38:VAL:HG23	1:D:40:VAL:HG13	2.02	0.42
1:D:46:PHE:CZ	1:D:58:LEU:HD11	2.55	0.41
1:D:80:PHE:CE2	1:D:82:TYR:HB2	2.54	0.41
1:A:91:ARG:HH12	1:A:96:THR:HG22	1.85	0.41
1:A:32:ASP:CG	1:A:47:ARG:HH12	2.23	0.41
1:B:125:PHE:HB3	1:B:134:PHE:HB3	2.02	0.41
1:D:93:LEU:HD21	1:D:168:TYR:CZ	2.55	0.41
1:B:103:LEU:HA	1:B:103:LEU:HD12	1.96	0.41
1:B:17:ASP:N	1:B:136:PHE:O	2.46	0.40
1:D:128:GLU:HB2	1:D:133:THR:OG1	2.21	0.40
1:A:9:LEU:O	1:A:182:ARG:NH1	2.53	0.40
1:D:107:ILE:HG23	1:D:119:VAL:HG13	2.01	0.40
1:A:252:PHE:CE2	1:A:304:ALA:HB1	2.55	0.40
1:B:180:ILE:HA	1:B:181:PRO:HD3	1.84	0.40
1:B:68:GLU:HG2	1:B:69:TRP:CD1	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/388 (100%)	373 (97%)	13 (3%)	0	100	100
1	B	386/388 (100%)	371 (96%)	14 (4%)	1 (0%)	41	70
1	C	214/388 (55%)	204 (95%)	10 (5%)	0	100	100
1	D	213/388 (55%)	203 (95%)	10 (5%)	0	100	100
All	All	1199/1552 (77%)	1151 (96%)	47 (4%)	1 (0%)	51	80

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	27	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/313 (100%)	304 (97%)	9 (3%)	42	74
1	B	313/313 (100%)	308 (98%)	5 (2%)	62	85
1	C	173/313 (55%)	171 (99%)	2 (1%)	71	89
1	D	172/313 (55%)	168 (98%)	4 (2%)	50	79
All	All	971/1252 (78%)	951 (98%)	20 (2%)	53	80

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	THR
1	A	102	GLN
1	A	130	THR
1	A	168	TYR
1	A	218	SER
1	A	269	ARG
1	A	281	ARG
1	A	333	LEU
1	A	344	TYR
1	B	133	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	168	TYR
1	B	269	ARG
1	B	344	TYR
1	B	373	THR
1	C	168	TYR
1	C	199	VAL
1	D	27	ARG
1	D	34	LEU
1	D	52	ARG
1	D	193	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PO4	C	401	-	4,4,4	0.90	0	6,6,6	0.42	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PO4	A	402	-	4,4,4	0.85	0	6,6,6	0.46	0
2	PO4	D	401	-	4,4,4	0.85	0	6,6,6	0.48	0
2	PO4	A	401	-	4,4,4	0.88	0	6,6,6	0.37	0
2	PO4	B	401	-	4,4,4	0.85	0	6,6,6	0.53	0
2	PO4	A	403	-	4,4,4	0.85	0	6,6,6	0.41	0
2	PO4	B	402	-	4,4,4	0.87	0	6,6,6	0.44	0
2	PO4	D	402	-	4,4,4	0.94	0	6,6,6	0.50	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	387/388 (99%)	-0.32	5 (1%) 77 77	27, 48, 82, 146	0
1	B	387/388 (99%)	-0.27	8 (2%) 63 62	29, 49, 83, 128	0
1	C	215/388 (55%)	0.03	13 (6%) 21 17	34, 56, 99, 127	0
1	D	215/388 (55%)	-0.22	10 (4%) 31 27	33, 50, 86, 135	0
All	All	1204/1552 (77%)	-0.22	36 (2%) 50 46	27, 51, 86, 146	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	27	ARG	7.6
1	A	28	GLY	7.5
1	B	28	GLY	7.0
1	B	27	ARG	6.2
1	C	30	MET	6.2
1	C	216	PRO	4.7
1	B	29	GLY	4.6
1	D	29	GLY	4.5
1	D	26	GLU	4.3
1	C	129	ALA	4.3
1	C	130	THR	4.1
1	D	216	PRO	3.9
1	D	28	GLY	3.7
1	D	3	PHE	3.5
1	C	131	GLY	3.4
1	B	23	TYR	3.1
1	C	29	GLY	3.0
1	C	65	ALA	2.8
1	C	27	ARG	2.8
1	C	68	GLU	2.7
1	A	25	THR	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	130	THR	2.6
1	D	215	ALA	2.6
1	D	2	THR	2.6
1	B	30	MET	2.6
1	C	24	LYS	2.5
1	C	28	GLY	2.4
1	C	31	ALA	2.3
1	C	48	TYR	2.3
1	D	30	MET	2.3
1	B	193	THR	2.3
1	A	26	GLU	2.2
1	B	372	ARG	2.2
1	B	2	THR	2.1
1	A	23	TYR	2.0
1	D	25	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	PO4	D	402	5/5	0.83	0.28	72,95,104,115	0
2	PO4	A	402	5/5	0.86	0.38	67,68,103,115	0
2	PO4	A	403	5/5	0.88	0.22	74,78,87,104	0
2	PO4	B	402	5/5	0.94	0.19	65,75,79,101	0
2	PO4	D	401	5/5	0.96	0.08	60,65,77,86	0
2	PO4	C	401	5/5	0.97	0.10	66,71,79,90	0
2	PO4	B	401	5/5	0.98	0.10	40,44,49,54	0
2	PO4	A	401	5/5	0.98	0.11	44,49,58,63	0



## 6.5 Other polymers

There are no such residues in this entry.