



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 03:18 am BST

PDB ID : 5OQF  
Title : Crystal structure of a single chain trimer composed of the MHC 1 heavy chain H2-Kb WT, beta-2microglobulin, and ovalbumin derived peptide  
Authors : Mikolajek, H.; Werner, J.M.  
Deposited on : 2017-08-11  
Resolution : 2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6980 (2.30-2.26)
Clashscore	141614	7711 (2.30-2.26)
Ramachandran outliers	138981	7597 (2.30-2.26)
Sidechain outliers	138945	7598 (2.30-2.26)
RSRZ outliers	127900	6849 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	431	
1	B	431	

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 6727 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-2-microglobulin,H-2 class I histocompatibility antigen, K-B alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	388	3150	1993	548	591	18	0	0	0
1	B	388	3158	1999	549	592	18	0	1	0

There are 102 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP P01887
A	2	ILE	-	expression tag	UNP P01887
A	3	ILE	-	expression tag	UNP P01887
A	4	ASN	-	expression tag	UNP P01887
A	5	PHE	-	expression tag	UNP P01887
A	6	GLU	-	expression tag	UNP P01887
A	7	LYS	-	expression tag	UNP P01887
A	8	LEU	-	expression tag	UNP P01887
A	9	GLY	-	expression tag	UNP P01887
A	10	CYS	-	expression tag	UNP P01887
A	11	GLY	-	expression tag	UNP P01887
A	12	ALA	-	expression tag	UNP P01887
A	13	SER	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	SER	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	GLY	-	expression tag	UNP P01887
A	?	SER	-	expression tag	UNP P01887
A	?	GLY	-	linker	UNP P01887

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	SER	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	SER	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	SER	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	GLY	-	linker	UNP P01887
A	?	SER	-	linker	UNP P01887
A	143	GLY	-	linker	UNP P01887
A	226	CYS	TYR	engineered mutation	UNP P01901
A	426	HIS	-	expression tag	UNP P01901
A	427	HIS	-	expression tag	UNP P01901
A	428	HIS	-	expression tag	UNP P01901
A	429	HIS	-	expression tag	UNP P01901
A	430	HIS	-	expression tag	UNP P01901
A	431	HIS	-	expression tag	UNP P01901
B	1	SER	-	expression tag	UNP P01887
B	2	ILE	-	expression tag	UNP P01887
B	3	ILE	-	expression tag	UNP P01887
B	4	ASN	-	expression tag	UNP P01887
B	5	PHE	-	expression tag	UNP P01887
B	6	GLU	-	expression tag	UNP P01887
B	7	LYS	-	expression tag	UNP P01887
B	8	LEU	-	expression tag	UNP P01887
B	9	GLY	-	expression tag	UNP P01887
B	10	CYS	-	expression tag	UNP P01887
B	11	GLY	-	expression tag	UNP P01887
B	12	ALA	-	expression tag	UNP P01887
B	13	SER	-	expression tag	UNP P01887
B	?	GLY	-	expression tag	UNP P01887
B	?	GLY	-	expression tag	UNP P01887

*Continued on next page...*




*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
2	B	200	Total 200	O 200	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-2-microglobulin,H-2 class I histocompatibility antigen, K-B alpha chain



- Molecule 1: Beta-2-microglobulin,H-2 class I histocompatibility antigen, K-B alpha chain



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.65Å 90.08Å 89.33Å 90.00° 111.23° 90.00°	Depositor
Resolution (Å)	83.27 - 2.27 83.27 - 2.27	Depositor EDS
% Data completeness (in resolution range)	98.5 (83.27-2.27) 98.5 (83.27-2.27)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.31 (at 2.27Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, $R_{free}$	0.200 , 0.250 0.200 , 0.250	Depositor DCC
$R_{free}$ test set	2266 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.1	Xtrriage
Anisotropy	0.838	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 46.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6727	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.56 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.6214e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/3237	0.59	0/4392
1	B	0.43	0/3245	0.58	0/4403
All	All	0.43	0/6482	0.59	0/8795

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3150	0	3016	46	0
1	B	3158	0	3026	42	0
2	A	219	0	0	5	0
2	B	200	0	0	11	0
All	All	6727	0	6042	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:TYR:HE1	1:A:209:ALA:HB2	1.35	0.89

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:ILE:O	1:A:208:LYS:NZ	2.06	0.87
1:B:8:LEU:HD23	1:B:285:THR:HG21	1.59	0.85
1:B:336:ARG:O	1:B:338:GLU:N	2.16	0.78
1:A:166:GLU:OE1	1:A:187:TYR:OH	2.04	0.75
1:B:8:LEU:CD2	1:B:285:THR:HG21	2.16	0.75
1:B:7:LYS:H	1:B:7:LYS:HD2	1.54	0.71
1:B:161:GLU:OE1	1:B:217:ARG:NH1	2.24	0.70
1:B:11:GLY:HA3	1:B:284:ILE:HG13	1.72	0.70
1:A:39:GLU:H	1:A:42:LYS:CE	2.05	0.69
1:B:147:LEU:HB2	1:B:310:LEU:HD13	1.76	0.66
1:B:122:MET:HG3	1:B:346:TRP:HH2	1.61	0.65
1:B:148:ARG:NH1	2:B:502:HOH:O	2.32	0.62
1:A:42:LYS:HZ2	1:A:44:ASN:HB2	1.64	0.62
1:A:336:ARG:HB3	1:A:337:PRO:HD2	1.82	0.61
1:A:8:LEU:HD21	1:A:258:TYR:CE2	2.37	0.60
1:A:8:LEU:HD12	1:A:265:TYR:CE2	2.36	0.60
1:A:39:GLU:H	1:A:42:LYS:HE2	1.67	0.59
1:A:357:LEU:HD22	1:A:403:VAL:HG22	1.85	0.59
1:B:156:ARG:NH2	2:B:508:HOH:O	2.35	0.58
1:B:81:LYS:NZ	2:B:505:HOH:O	2.31	0.57
1:B:8:LEU:HD23	1:B:285:THR:CG2	2.34	0.57
1:A:10:CYS:SG	1:A:222:THR:HG22	2.46	0.56
1:B:8:LEU:HD22	1:B:265:TYR:HE2	1.70	0.56
1:B:257:GLN:NE2	2:B:504:HOH:O	2.30	0.55
1:B:314:LEU:O	1:B:318:ASN:HB2	2.06	0.55
1:B:362:ASN:HB2	2:B:627:HOH:O	2.07	0.55
1:A:42:LYS:NZ	1:A:44:ASN:HD22	2.05	0.55
1:B:118:TRP:HE1	1:B:122:MET:HE1	1.72	0.55
1:A:83:TRP:CE2	1:A:259:ALA:HB2	2.43	0.54
1:B:29:GLN:NE2	2:B:501:HOH:O	2.15	0.54
1:B:197:GLU:OE2	1:B:312:ARG:NH2	2.39	0.53
1:A:39:GLU:H	1:A:42:LYS:HE3	1.73	0.53
1:A:339:ASP:N	1:A:339:ASP:OD1	2.32	0.52
1:A:217:ARG:NH1	2:A:508:HOH:O	2.42	0.52
1:A:360:GLN:HB2	1:A:400:THR:HG22	1.91	0.52
1:B:318:ASN:O	1:B:322:LEU:HD12	2.11	0.51
1:A:8:LEU:HD12	1:A:265:TYR:HE2	1.76	0.51
1:B:186:ARG:HH22	1:B:203:GLU:HG2	1.76	0.51
1:A:219:ASP:O	1:A:223:LEU:HD23	2.11	0.50
1:A:121:ASP:HB3	1:A:334:HIS:CE1	2.46	0.50
1:A:287:HIS:O	1:A:291:GLN:HG2	2.12	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:GLN:HG2	1:A:55:PRO:HD3	1.95	0.49
1:B:104:ARG:HD3	1:B:115:THR:OG1	2.13	0.49
1:A:145:HIS:ND1	1:A:171:ASP:OD2	2.47	0.48
1:B:357:LEU:HD22	1:B:403:VAL:HG22	1.96	0.48
1:A:61:GLN:HG3	2:A:599:HOH:O	2.14	0.48
1:A:336:ARG:HD3	1:A:342:THR:OG1	2.14	0.47
1:A:403:VAL:HB	1:A:412:LEU:HB2	1.96	0.47
1:B:163:ARG:HD2	1:B:181:ASP:OD2	2.14	0.47
1:A:199:PRO:HB3	1:B:279:ASP:HA	1.96	0.47
1:A:197:GLU:OE1	1:A:312:ARG:NH2	2.44	0.47
1:A:223:LEU:HD12	1:A:227:TYR:CE1	2.50	0.47
1:B:118:TRP:HE1	1:B:122:MET:CE	2.27	0.46
1:A:42:LYS:HD2	1:A:42:LYS:O	2.14	0.46
1:B:254:GLY:HA3	1:B:302:LEU:HD13	1.97	0.46
1:A:8:LEU:HD13	1:A:285:THR:HG21	1.97	0.46
1:A:155:SER:HB3	1:A:220:LEU:HD13	1.97	0.46
1:A:336:ARG:HG3	1:A:341:VAL:HA	1.96	0.46
1:B:183:GLU:HG3	1:B:184:ASN:H	1.80	0.45
1:A:187:TYR:CE1	1:A:209:ALA:HB2	2.28	0.45
1:A:360:GLN:HB2	1:A:400:THR:CG2	2.47	0.45
1:B:145:HIS:HA	1:B:171:ASP:OD1	2.17	0.45
1:B:26:LYS:HE2	1:B:84:SER:HB3	2.00	0.44
1:B:51:THR:HG22	1:B:86:TYR:HB2	1.99	0.44
1:B:146:SER:O	1:B:170:VAL:HA	2.18	0.44
1:B:67:LYS:NZ	2:B:514:HOH:O	2.42	0.44
1:B:156:ARG:NH2	2:B:520:HOH:O	2.50	0.44
1:B:287:HIS:HB3	2:B:656:HOH:O	2.17	0.44
1:A:257:GLN:NE2	2:A:511:HOH:O	2.49	0.44
1:A:42:LYS:NZ	1:A:44:ASN:ND2	2.66	0.44
1:A:223:LEU:HD12	1:A:227:TYR:HE1	1.83	0.43
1:B:223:LEU:HD13	1:B:260:TYR:CD1	2.53	0.43
1:A:286:LYS:NZ	2:A:512:HOH:O	2.51	0.43
1:B:250:ARG:HD3	2:B:583:HOH:O	2.19	0.42
1:A:64:LYS:HB2	1:A:69:ILE:HD11	2.00	0.42
1:A:39:GLU:O	1:A:42:LYS:HE3	2.19	0.42
1:B:8:LEU:HD22	1:B:265:TYR:CE2	2.53	0.42
1:B:287:HIS:HD2	2:B:672:HOH:O	2.03	0.42
1:A:42:LYS:HZ3	1:A:44:ASN:HD22	1.69	0.41
1:B:333:HIS:NE2	1:B:341:VAL:HG11	2.35	0.41
1:B:187:TYR:HE1	1:B:209:ALA:HB2	1.86	0.41
1:A:42:LYS:NZ	1:A:44:ASN:HB2	2.33	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:231:LYS:HE3	1:B:231:LYS:HB2	1.77	0.41
1:A:192:ARG:NE	1:A:195:GLU:OE2	2.54	0.40
1:A:323:ARG:NH2	2:A:504:HOH:O	2.30	0.40
1:A:399:TYR:O	1:A:415:ARG:HG2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	382/431 (89%)	370 (97%)	12 (3%)	0	100	100
1	B	383/431 (89%)	370 (97%)	11 (3%)	2 (0%)	29	34
All	All	765/862 (89%)	740 (97%)	23 (3%)	2 (0%)	41	49

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	337	PRO
1	B	70	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/356 (95%)	333 (99%)	4 (1%)	71	82
1	B	338/356 (95%)	332 (98%)	6 (2%)	59	72
All	All	675/712 (95%)	665 (98%)	10 (2%)	69	77

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	LYS
1	A	93	PHE
1	A	163	ARG
1	A	181	ASP
1	B	93	PHE
1	B	181	ASP
1	B	187	TYR
1	B	215	SER
1	B	414[A]	LEU
1	B	414[B]	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	388/431 (90%)	0.09	12 (3%) 49 54	22, 37, 57, 79	0
1	B	388/431 (90%)	0.09	8 (2%) 63 69	22, 36, 55, 77	0
All	All	776/862 (90%)	0.09	20 (2%) 56 62	22, 37, 57, 79	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	367	ILE	6.2
1	B	369	ASP	4.0
1	B	368	GLN	3.9
1	A	337	PRO	3.9
1	B	317	GLY	3.6
1	A	12	ALA	3.6
1	A	122	MET	3.4
1	A	318	ASN	3.2
1	B	121	ASP	2.7
1	A	336	ARG	2.7
1	B	337	PRO	2.6
1	A	13	SER	2.4
1	A	339	ASP	2.4
1	B	183	GLU	2.4
1	B	318	ASN	2.3
1	A	369	ASP	2.1
1	B	336	ARG	2.1
1	A	322	LEU	2.1
1	A	121	ASP	2.0
1	A	143	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.