



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2020 – 04:52 AM BST

PDB ID : 3PAY  
Title : Crystal structure of a putative adhesin (BACOVA\_04077) from *Bacteroides ovatus* at 2.50 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2010-10-19  
Resolution : 2.50 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	 4% 89% 7%
1	B	314	 5% 90% 8%
1	C	314	 3% 86% 11%
1	D	314	 3% 91% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PEG	C	333	-	-	X	-

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10020 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called putative adhesin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	302	2400	1543	398	450	2	7	0	2	0
1	B	309	2451	1571	403	468	2	7	0	0	0
1	C	307	2433	1562	393	469	2	7	0	1	0
1	D	313	2476	1582	405	480	2	7	0	1	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP A7M1U3
B	0	GLY	-	expression tag	UNP A7M1U3
C	0	GLY	-	expression tag	UNP A7M1U3
D	0	GLY	-	expression tag	UNP A7M1U3

- Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total C O 7 4 3	0	0
2	C	1	Total C O 7 4 3	0	0
2	D	1	Total C O 7 4 3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	47	Total O 47 47	0	0
3	B	64	Total O 64 64	0	0
3	C	74	Total O 74 74	0	0
3	D	54	Total O 54 54	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: putative adhesin



- Molecule 1: putative adhesin




- Molecule 1: putative adhesin



- Molecule 1: putative adhesin





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.11Å 78.78Å 84.11Å 93.88° 102.08° 103.54°	Depositor
Resolution (Å)	29.46 – 2.50 29.47 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (29.46-2.50) 96.4 (29.47-2.50)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.46 (at 2.51Å)	Xtrriage
Refinement program	BUSTER-TNT BUSTER 2.8.0, BUSTER 2.8.0	Depositor
R, $R_{free}$	0.214 , 0.239 0.221 , 0.242	Depositor DCC
$R_{free}$ test set	2758 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.1	Xtrriage
Anisotropy	0.312	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 45.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	10020	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.86% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.48	0/2451	0.70	0/3326
1	B	0.51	0/2504	0.71	0/3400
1	C	0.54	0/2487	0.74	0/3377
1	D	0.50	0/2533	0.71	0/3443
All	All	0.51	0/9975	0.72	0/13546

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2400	0	2270	14	0
1	B	2451	0	2328	13	0
1	C	2433	0	2293	23	0
1	D	2476	0	2319	14	0
2	C	14	0	19	8	0
2	D	7	0	10	2	0
3	A	47	0	0	0	0
3	B	64	0	0	0	0
3	C	74	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	54	0	0	0	0
All	All	10020	0	9239	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:295:ARG:HH22	2:C:334:PEG:H41	1.51	0.75
1:A:285:THR:HB	2:C:333:PEG:H41	1.80	0.64
1:B:59:LEU:HD11	1:B:109:LEU:HD21	1.79	0.64
1:C:134:GLU:CD	1:C:134:GLU:H	2.01	0.63
1:D:135:THR:HG22	1:D:137:GLY:H	1.64	0.62
1:C:245:ASN:O	2:C:333:PEG:H32	2.02	0.58
1:A:73:VAL:HG13	1:A:77:ALA:HB3	1.88	0.56
1:C:188:VAL:HG13	1:C:233:VAL:HG21	1.87	0.55
1:B:73:VAL:HG13	1:B:77:ALA:HB3	1.88	0.55
1:D:282:TYR:CD1	2:D:335:PEG:H22	2.42	0.55
1:D:73:VAL:HG13	1:D:77:ALA:HB3	1.88	0.54
1:C:73:VAL:HG13	1:C:77:ALA:HB3	1.90	0.54
1:C:246:ARG:HE	2:C:333:PEG:H11	1.72	0.54
1:C:59:LEU:HD11	1:C:109:LEU:HD21	1.90	0.53
1:A:197:ILE:HD12	1:A:238:LEU:HD11	1.92	0.52
1:D:197:ILE:HD12	1:D:238:LEU:HD11	1.90	0.52
1:A:59:LEU:HD11	1:A:109:LEU:HD21	1.92	0.51
1:D:45:TYR:HB2	2:D:335:PEG:H21	1.92	0.51
1:C:41:TYR:HD1	2:C:334:PEG:H22	1.76	0.51
1:A:221:PRO:HA	1:A:238:LEU:HB3	1.93	0.51
1:C:265:TRP:CE2	1:C:316:ILE:HG21	2.47	0.50
1:C:295:ARG:NH2	2:C:334:PEG:H41	2.24	0.50
1:D:58:VAL:O	1:D:70:THR:HA	2.12	0.50
1:D:58:VAL:HG13	1:D:97:VAL:HG13	1.93	0.49
1:A:144:HIS:HE2	1:A:175:ARG:NE	2.11	0.49
1:B:318:ILE:HG23	1:B:327:LEU:HD11	1.95	0.49
1:C:58:VAL:HG13	1:C:97:VAL:HG13	1.94	0.48
1:C:197:ILE:HD12	1:C:238:LEU:HD11	1.95	0.48
1:D:59:LEU:HD11	1:D:109:LEU:HD21	1.95	0.48
1:B:197:ILE:HD12	1:B:238:LEU:HD11	1.96	0.48
1:C:245:ASN:HB3	2:C:333:PEG:H31	1.95	0.48
1:C:257:ASP:OD1	1:C:260:THR:HG22	2.14	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:275:ILE:HD12	1:D:318:ILE:HD11	1.96	0.47
1:D:318:ILE:HG23	1:D:327:LEU:HD11	1.95	0.47
1:B:221:PRO:HA	1:B:238:LEU:HB3	1.97	0.47
1:C:290:GLN:HB2	2:C:333:PEG:H42	1.98	0.46
1:C:221:PRO:HA	1:C:238:LEU:HB3	1.97	0.46
1:A:109:LEU:HG	1:A:117:LEU:HD11	1.98	0.46
1:B:109:LEU:HG	1:B:117:LEU:HD11	1.97	0.46
1:B:174:ASN:ND2	1:B:241:MSE:O	2.49	0.46
1:A:144:HIS:HE2	1:A:175:ARG:CZ	2.28	0.46
1:A:318:ILE:HG23	1:A:327:LEU:HD11	1.98	0.46
1:C:60:VAL:HG11	1:C:90:LEU:HD21	1.99	0.44
1:D:109:LEU:HG	1:D:117:LEU:HD11	1.99	0.44
1:D:207:TRP:CE2	1:D:208:GLU:HG3	2.52	0.44
1:C:23:SER:HA	1:C:30:ARG:HA	1.99	0.44
1:B:207:TRP:CE2	1:B:208:GLU:HG3	2.52	0.44
1:D:125:GLN:HA	1:D:207:TRP:CD1	2.52	0.44
1:C:207:TRP:CE2	1:C:208:GLU:HG3	2.53	0.44
1:A:207:TRP:CE2	1:A:208:GLU:HG3	2.52	0.43
1:C:20:SER:HB3	1:C:161:ASN:OD1	2.18	0.43
1:A:125:GLN:HA	1:A:207:TRP:CD1	2.53	0.43
1:A:58:VAL:O	1:A:70:THR:HA	2.19	0.43
1:C:53:VAL:O	1:C:75:VAL:HG21	2.18	0.43
1:C:109:LEU:HG	1:C:117:LEU:HD11	2.01	0.43
1:A:323:TRP:NE1	1:B:276:ALA:O	2.52	0.43
1:B:58:VAL:HG21	1:B:84:MSE:SE	2.69	0.43
1:C:125:GLN:HA	1:C:207:TRP:CD1	2.55	0.42
1:B:260:THR:HG23	1:B:262:ALA:H	1.85	0.42
1:D:260:THR:HG23	1:D:262:ALA:H	1.85	0.42
1:B:125:GLN:HA	1:B:207:TRP:CD1	2.54	0.41
1:A:260:THR:HG23	1:A:262:ALA:H	1.86	0.41
1:B:20:SER:HB3	1:B:161:ASN:HA	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	298/314 (95%)	281 (94%)	17 (6%)	0	100	100
1	B	305/314 (97%)	293 (96%)	12 (4%)	0	100	100
1	C	300/314 (96%)	289 (96%)	11 (4%)	0	100	100
1	D	312/314 (99%)	297 (95%)	13 (4%)	2 (1%)	25	43
All	All	1215/1256 (97%)	1160 (96%)	53 (4%)	2 (0%)	47	68

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	229	ILE
1	D	228	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/271 (93%)	251 (99%)	2 (1%)	81	93
1	B	263/271 (97%)	259 (98%)	4 (2%)	65	85
1	C	261/271 (96%)	256 (98%)	5 (2%)	57	80
1	D	263/271 (97%)	260 (99%)	3 (1%)	73	89
All	All	1040/1084 (96%)	1026 (99%)	14 (1%)	67	87

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	239	ASN
1	A	278	PRO
1	B	134	GLU
1	B	185	GLU
1	B	201	GLU
1	B	239	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	22	ASP
1	C	27	ASP
1	C	112	ASN
1	C	134	GLU
1	C	239	ASN
1	D	50	ASN
1	D	239	ASN
1	D	267	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	168	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PEG	C	334	-	6,6,6	0.69	0	5,5,5	0.42	0
2	PEG	D	335	-	6,6,6	0.70	0	5,5,5	0.96	1 (20%)
2	PEG	C	333	-	6,6,6	1.07	0	5,5,5	1.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	C	334	-	-	1/4/4/4	-
2	PEG	D	335	-	-	0/4/4/4	-
2	PEG	C	333	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	335	PEG	O2-C2-C1	2.06	119.11	110.07

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	334	PEG	C4-C3-O2-C2

There are no ring outliers.

3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	334	PEG	3	0
2	D	335	PEG	2	0
2	C	333	PEG	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	295/314 (93%)	0.09	13 (4%) 34 37	30, 50, 74, 92	0
1	B	302/314 (96%)	0.11	17 (5%) 24 25	26, 41, 67, 93	0
1	C	300/314 (95%)	-0.09	8 (2%) 54 58	19, 34, 58, 74	0
1	D	306/314 (97%)	-0.01	8 (2%) 56 59	22, 38, 72, 97	0
All	All	1203/1256 (95%)	0.03	46 (3%) 40 43	19, 41, 70, 97	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	229	ILE	6.4
1	D	228	GLU	4.7
1	D	230	SER	4.7
1	C	28	LEU	4.4
1	B	28	LEU	4.3
1	D	28	LEU	4.0
1	D	27	ASP	4.0
1	B	311	GLY	3.9
1	B	51	PRO	3.5
1	A	232	VAL	3.5
1	A	223	TYR	3.4
1	A	51	PRO	3.2
1	B	191	THR	3.1
1	B	310	GLY	3.1
1	A	27	ASP	3.0
1	D	311	GLY	3.0
1	A	310	GLY	2.8
1	A	309	ASN	2.8
1	C	187	LYS	2.8
1	C	230	SER	2.8
1	A	197	ILE	2.7

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	233	VAL	2.7
1	A	259	ASN	2.7
1	C	184	GLU	2.6
1	A	113	ALA	2.6
1	B	20	SER	2.6
1	B	135	THR	2.6
1	A	311	GLY	2.5
1	C	231	ASP	2.4
1	B	81	GLY	2.4
1	B	100	VAL	2.4
1	A	207	TRP	2.3
1	B	309	ASN	2.3
1	B	75	VAL	2.3
1	B	188	VAL	2.3
1	C	240	THR	2.3
1	D	226	PRO	2.2
1	A	240	THR	2.2
1	C	186	ASN	2.1
1	B	80	GLY	2.1
1	B	189	ASP	2.1
1	D	29	PRO	2.1
1	B	240	THR	2.1
1	B	147	PHE	2.0
1	B	27	ASP	2.0
1	C	81	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PEG	C	334	7/7	0.77	0.34	56,57,60,63	0
2	PEG	C	333	7/7	0.88	0.53	40,43,44,49	0
2	PEG	D	335	7/7	0.91	0.28	43,44,45,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.