



## wwPDB EM Validation Summary Report ⓘ

Jan 29, 2022 – 06:22 am GMT

PDB ID : 7QJ9  
EMDB ID : EMD-14014  
Title : Structure of recombinant human gamma-Tubulin Ring Complex 10-spoked assembly intermediate (spokes 3-12, homogeneous dataset)  
Authors : Zupa, E.; Pfeffer, S.  
Deposited on : 2021-12-16  
Resolution : 8.10 Å (reported)  
Based on initial models : 6L81, 6V6S, 7AS4, 6X0U

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.0.dev97  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.26


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 8.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	J	1024	
1	l	1024	
2	e	375	
3	b	82	
3	d	82	
3	i	82	
3	k	82	
3	m	82	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	907	55% 9% 36%
4	F	907	56% 9% 34%
4	H	907	55% 9% 35%
4	a	907	12% 87%
4	h	907	11% 89%
4	j	907	12% 88%
5	C	902	6% 57% 11% 31%
5	E	902	62% 9% 29%
5	G	902	61% 10% 29%
6	I	667	66% 12% 22%
6	K	667	70% 13% 16%
7	L	1819	27% 69%
7	c	1819	9% 91%
8	Q	451	27% 63% 26% 7% 7%
8	R	451	12% 63% 25% 7% 7%
8	S	451	64% 25% 7% 7%
8	T	451	62% 27% 7% 7%
8	U	451	64% 25% 7% 7%
8	V	451	63% 26% 7% 7%
8	W	451	63% 26% 7% 7%
8	X	451	63% 26% 7% 7%
8	Y	451	63% 26% 7% 7%
8	Z	451	5% 64% 25% 7% 7%

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 91535 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma-tubulin complex component 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	l	108	Total 847	C 539	N 150	O 157	S 1	0	0
1	J	534	Total 4429	C 2893	N 737	O 776	S 23	0	0

- Molecule 2 is a protein called actin, cytoplasmic 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	e	364	Total 2847	C 1803	N 476	O 548	S 20	0	0

- Molecule 3 is a protein called Mitotic-spindle organizing protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	b	65	Total 484	C 299	N 85	O 96	S 4	0	0
3	i	65	Total 484	C 299	N 85	O 96	S 4	0	0
3	m	65	Total 484	C 299	N 85	O 96	S 4	0	0
3	k	65	Total 484	C 299	N 85	O 96	S 4	0	0
3	d	59	Total 454	C 281	N 79	O 90	S 4	0	0

- Molecule 4 is a protein called Gamma-tubulin complex component 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	a	116	Total 933	C 591	N 171	O 169	S 2	0	0
4	D	581	Total 4796	C 3061	N 842	O 868	S 25	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf	Trace
4	F	599	Total	C	N	O	S	0	0
			4941	3151	871	894	25		
4	H	594	Total	C	N	O	S	0	0
			4907	3130	864	888	25		
4	h	99	Total	C	N	O	S	0	0
			803	509	148	144	2		
4	j	107	Total	C	N	O	S	0	0
			843	533	156	152	2		

- Molecule 5 is a protein called Gamma-tubulin complex component 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	C	620	Total	C	N	O	S	0	0
			5044	3257	845	910	32		
5	E	638	Total	C	N	O	S	0	0
			5202	3354	873	942	33		
5	G	640	Total	C	N	O	S	0	0
			5206	3354	875	944	33		

- Molecule 6 is a protein called Gamma-tubulin complex component 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	I	521	Total	C	N	O	S	0	0
			4225	2737	720	750	18		
6	K	562	Total	C	N	O	S	0	0
			4579	2964	781	816	18		

- Molecule 7 is a protein called Gamma-tubulin complex component 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	L	566	Total	C	N	O	S	0	0
			4587	3000	773	789	25		
7	c	158	Total	C	N	O	S	0	0
			1220	771	209	232	8		

- Molecule 8 is a protein called Tubulin gamma-1 chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	Q	420	Total	C	N	O	S	0	0
			3373	2134	586	638	15		
8	R	420	Total	C	N	O	S	0	0
			3373	2134	586	638	15		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf	Trace
8	S	420	Total	C	N	O	S	0	0
			3373	2134	586	638	15		
8	T	420	Total	C	N	O	S	0	0
			3373	2134	586	638	15		
8	U	420	Total	C	N	O	S	0	0
			3373	2134	586	638	15		
8	V	420	Total	C	N	O	S	0	0
			3373	2134	586	638	15		
8	W	420	Total	C	N	O	S	0	0
			3373	2134	586	638	15		
8	X	420	Total	C	N	O	S	0	0
			3373	2134	586	638	15		
8	Y	420	Total	C	N	O	S	0	0
			3373	2134	586	638	15		
8	Z	420	Total	C	N	O	S	0	0
			3373	2134	586	638	15		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		AltConf
9	1	6	Total	O	0
			6	6	

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Gamma-tubulin complex component 5















• Molecule 3: Mitotic-spindle organizing protein 1




• Molecule 3: Mitotic-spindle organizing protein 1



• Molecule 3: Mitotic-spindle organizing protein 1



• Molecule 3: Mitotic-spindle organizing protein 1


















• Molecule 5: Gamma-tubulin complex component 2




• Molecule 5: Gamma-tubulin complex component 2











• Molecule 7: Gamma-tubulin complex component 6













• Molecule 8: Tubulin gamma-1 chain




• Molecule 8: Tubulin gamma-1 chain




• Molecule 8: Tubulin gamma-1 chain

Chain U:  64% 25% 7%




• Molecule 8: Tubulin gamma-1 chain


Chain V:  63% 26% 7%




• Molecule 8: Tubulin gamma-1 chain

Chain W:  63% 26% 7%







• Molecule 8: Tubulin gamma-1 chain







• Molecule 8: Tubulin gamma-1 chain





• Molecule 8: Tubulin gamma-1 chain





## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	21860	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	35	Depositor
Minimum defocus (nm)	2000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.329	Depositor
Minimum map value	-0.148	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.010	Depositor
Recommended contour level	0.0299	Depositor
Map size (Å)	532.0, 532.0, 532.0	wwPDB
Map dimensions	200, 200, 200	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	2.66, 2.66, 2.66	Depositor

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	J	0.47	3/4525 (0.1%)	0.73	2/6119 (0.0%)
1	l	0.39	0/863	0.71	2/1166 (0.2%)
2	e	0.54	1/2908 (0.0%)	0.74	3/3938 (0.1%)
3	b	0.40	0/484	0.69	0/653
3	d	0.43	0/454	0.80	1/611 (0.2%)
3	i	0.41	0/484	0.70	0/653
3	k	0.41	0/484	0.63	1/653 (0.2%)
3	m	0.34	0/484	0.67	1/653 (0.2%)
4	D	0.39	0/4897	0.71	3/6610 (0.0%)
4	F	0.38	0/5044	0.66	4/6809 (0.1%)
4	H	0.45	1/5009 (0.0%)	0.70	2/6761 (0.0%)
4	a	0.45	0/948	0.75	1/1277 (0.1%)
4	h	0.37	0/815	0.65	1/1096 (0.1%)
4	j	0.36	0/855	0.70	2/1152 (0.2%)
5	C	0.42	0/5151	0.73	8/6955 (0.1%)
5	E	0.41	1/5311 (0.0%)	0.68	3/7169 (0.0%)
5	G	0.41	0/5315	0.69	3/7175 (0.0%)
6	I	0.49	1/4322 (0.0%)	0.71	4/5853 (0.1%)
6	K	0.49	3/4683 (0.1%)	0.78	15/6338 (0.2%)
7	L	0.41	1/4697 (0.0%)	0.69	7/6348 (0.1%)
7	c	0.40	0/1235	0.77	2/1664 (0.1%)
8	Q	0.35	0/3441	0.66	3/4661 (0.1%)
8	R	0.35	0/3441	0.66	3/4661 (0.1%)
8	S	0.35	0/3441	0.66	3/4661 (0.1%)
8	T	0.35	0/3441	0.66	3/4661 (0.1%)
8	U	0.35	0/3441	0.66	3/4661 (0.1%)
8	V	0.35	0/3441	0.66	3/4661 (0.1%)
8	W	0.35	0/3441	0.66	3/4661 (0.1%)
8	X	0.35	0/3441	0.66	3/4661 (0.1%)
8	Y	0.35	0/3441	0.66	3/4661 (0.1%)
8	Z	0.35	0/3441	0.66	3/4661 (0.1%)
All	All	0.40	11/93378 (0.0%)	0.69	95/126263 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if

the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	J	0	4
2	e	0	1
4	D	0	1
4	F	0	1
4	H	0	5
5	C	0	1
5	E	0	1
5	G	0	3
6	I	0	2
6	K	0	4
8	Q	0	2
8	R	0	2
8	S	0	2
8	T	0	2
8	U	0	2
8	V	0	2
8	W	0	2
8	X	0	2
8	Y	0	2
8	Z	0	2
All	All	0	43

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	e	243	PRO	N-CD	18.00	1.73	1.47
6	I	361	TYR	CD2-CE2	-9.36	1.25	1.39
6	K	651	TYR	CB-CG	-7.87	1.39	1.51
6	K	651	TYR	CD1-CE1	-7.14	1.28	1.39
6	K	651	TYR	CD2-CE2	-6.94	1.28	1.39

The worst 5 of 95 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	S	339	ARG	NE-CZ-NH2	11.98	126.29	120.30
8	V	339	ARG	NE-CZ-NH2	11.97	126.28	120.30
8	T	339	ARG	NE-CZ-NH2	11.96	126.28	120.30
8	W	339	ARG	NE-CZ-NH2	11.92	126.26	120.30
8	Z	339	ARG	NE-CZ-NH2	11.91	126.26	120.30

There are no chirality outliers.

5 of 43 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
5	C	464	GLY	Peptide
4	D	888	ALA	Peptide
5	E	370	SER	Peptide
4	F	269	ASN	Peptide
2	e	200	PHE	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	J	4429	0	4482	43	0
1	l	847	0	789	0	0
2	e	2847	0	2810	0	0
3	b	484	0	512	0	0
3	d	454	0	482	0	0
3	i	484	0	512	0	0
3	k	484	0	512	0	0
3	m	484	0	512	0	0
4	D	4796	0	4775	50	0
4	F	4941	0	4935	48	0
4	H	4907	0	4896	59	0
4	a	933	0	953	0	0
4	h	803	0	831	0	0
4	j	843	0	846	0	0
5	C	5044	0	5081	67	0
5	E	5202	0	5241	51	0
5	G	5206	0	5230	61	0
6	I	4225	0	4259	49	0
6	K	4579	0	4586	53	0
7	L	4587	0	4636	48	0
7	c	1220	0	1231	0	0
8	Q	3373	0	3325	67	0
8	R	3373	0	3325	70	0
8	S	3373	0	3325	64	0
8	T	3373	0	3325	71	0
8	U	3373	0	3325	68	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	V	3373	0	3325	73	0
8	W	3373	0	3325	70	0
8	X	3373	0	3325	70	0
8	Y	3373	0	3325	73	0
8	Z	3373	0	3325	66	0
9	l	6	0	0	0	0
All	All	91535	0	91361	1146	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 1146 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:F:879:ARG:HD2	8:T:337:LEU:HD23	1.58	0.83
8:W:56:ASP:HB2	8:X:296:ARG:HD2	1.62	0.81
5:E:734:LEU:HB2	5:E:739:LEU:HD11	1.69	0.75
5:G:560:ARG:HB2	8:V:339:ARG:HE	1.52	0.74
5:E:696:TYR:O	5:E:700:GLU:HB2	1.89	0.71

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	J	506/1024 (49%)	464 (92%)	38 (8%)	4 (1%)	19 60
1	l	104/1024 (10%)	97 (93%)	5 (5%)	2 (2%)	8 38
2	e	360/375 (96%)	345 (96%)	15 (4%)	0	100 100
3	b	63/82 (77%)	61 (97%)	2 (3%)	0	100 100
3	d	57/82 (70%)	56 (98%)	1 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	i	63/82 (77%)	63 (100%)	0	0	100	100
3	k	63/82 (77%)	61 (97%)	2 (3%)	0	100	100
3	m	63/82 (77%)	63 (100%)	0	0	100	100
4	D	571/907 (63%)	552 (97%)	17 (3%)	2 (0%)	34	72
4	F	591/907 (65%)	561 (95%)	30 (5%)	0	100	100
4	H	584/907 (64%)	557 (95%)	23 (4%)	4 (1%)	22	63
4	a	112/907 (12%)	109 (97%)	3 (3%)	0	100	100
4	h	97/907 (11%)	94 (97%)	3 (3%)	0	100	100
4	j	105/907 (12%)	99 (94%)	6 (6%)	0	100	100
5	C	606/902 (67%)	575 (95%)	28 (5%)	3 (0%)	29	69
5	E	626/902 (69%)	587 (94%)	38 (6%)	1 (0%)	47	81
5	G	628/902 (70%)	596 (95%)	29 (5%)	3 (0%)	29	69
6	I	511/667 (77%)	484 (95%)	25 (5%)	2 (0%)	34	72
6	K	548/667 (82%)	530 (97%)	16 (3%)	2 (0%)	34	72
7	L	540/1819 (30%)	495 (92%)	38 (7%)	7 (1%)	12	48
7	c	148/1819 (8%)	142 (96%)	6 (4%)	0	100	100
8	Q	408/451 (90%)	386 (95%)	22 (5%)	0	100	100
8	R	408/451 (90%)	386 (95%)	22 (5%)	0	100	100
8	S	408/451 (90%)	386 (95%)	22 (5%)	0	100	100
8	T	408/451 (90%)	386 (95%)	22 (5%)	0	100	100
8	U	408/451 (90%)	386 (95%)	22 (5%)	0	100	100
8	V	408/451 (90%)	386 (95%)	22 (5%)	0	100	100
8	W	408/451 (90%)	386 (95%)	22 (5%)	0	100	100
8	X	408/451 (90%)	386 (95%)	22 (5%)	0	100	100
8	Y	408/451 (90%)	386 (95%)	22 (5%)	0	100	100
8	Z	408/451 (90%)	386 (95%)	22 (5%)	0	100	100
All	All	11026/20463 (54%)	10451 (95%)	545 (5%)	30 (0%)	44	77

5 of 30 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	I	121	PRO
5	C	465	HIS

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
5	G	581	ASP
4	H	454	VAL
4	H	632	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	J	498/933 (53%)	493 (99%)	5 (1%)	76 86
1	l	84/933 (9%)	82 (98%)	2 (2%)	49 69
2	e	310/318 (98%)	306 (99%)	4 (1%)	69 81
3	b	53/62 (86%)	52 (98%)	1 (2%)	57 75
3	d	53/62 (86%)	53 (100%)	0	100 100
3	i	53/62 (86%)	52 (98%)	1 (2%)	57 75
3	k	53/62 (86%)	51 (96%)	2 (4%)	33 57
3	m	53/62 (86%)	53 (100%)	0	100 100
4	D	525/798 (66%)	520 (99%)	5 (1%)	76 86
4	F	542/798 (68%)	538 (99%)	4 (1%)	84 90
4	H	539/798 (68%)	537 (100%)	2 (0%)	91 94
4	a	101/798 (13%)	99 (98%)	2 (2%)	55 74
4	h	88/798 (11%)	88 (100%)	0	100 100
4	j	88/798 (11%)	88 (100%)	0	100 100
5	C	556/791 (70%)	553 (100%)	3 (0%)	88 93
5	E	574/791 (73%)	573 (100%)	1 (0%)	93 96
5	G	572/791 (72%)	568 (99%)	4 (1%)	84 90
6	I	472/594 (80%)	467 (99%)	5 (1%)	73 84
6	K	509/594 (86%)	505 (99%)	4 (1%)	81 89
7	L	501/1546 (32%)	499 (100%)	2 (0%)	91 94
7	c	135/1546 (9%)	134 (99%)	1 (1%)	84 90

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
8	Q	376/400 (94%)	329 (88%)	47 (12%)	4	19
8	R	376/400 (94%)	329 (88%)	47 (12%)	4	19
8	S	376/400 (94%)	329 (88%)	47 (12%)	4	19
8	T	376/400 (94%)	329 (88%)	47 (12%)	4	19
8	U	376/400 (94%)	329 (88%)	47 (12%)	4	19
8	V	376/400 (94%)	329 (88%)	47 (12%)	4	19
8	W	376/400 (94%)	329 (88%)	47 (12%)	4	19
8	X	376/400 (94%)	329 (88%)	47 (12%)	4	19
8	Y	376/400 (94%)	329 (88%)	47 (12%)	4	19
8	Z	376/400 (94%)	329 (88%)	47 (12%)	4	19
All	All	10119/17935 (56%)	9601 (95%)	518 (5%)	27	48

5 of 518 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
8	Y	373	VAL
8	Z	112	GLU
8	Y	369	SER
8	T	112	GLU
8	T	50	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 144 such sidechains are listed below:

Mol	Chain	Res	Type
8	V	322	ASN
4	j	78	HIS
8	W	299	GLN
8	Y	299	GLN
4	H	703	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.



## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-14014. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)

#### 6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 134



Y Index: 84



Z Index: 72


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal surface views [i](#)

### 6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.0299. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 1956  $\text{nm}^3$ ; this corresponds to an approximate mass of 1767 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)



\*Reported resolution corresponds to spatial frequency of  $0.123 \text{\AA}^{-1}$

## 8 Fourier-Shell correlation


This section was not generated. No FSC curve or half-maps provided.



## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-14014 and PDB model 7QJ9. Per-residue inclusion information can be found in section 3 on page 7.

### 9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.0299 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Atom inclusion [i](#)



At the recommended contour level, 93% of all backbone atoms, 83% of all non-hydrogen atoms, are inside the map.