



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 2, 2023 – 11:04 AM EDT

PDB ID : 3QNC
Title : Crystal Structure of a Rationally Designed OXA-10 Variant Showing Carbapenemase Activity, OXA-10loop48
Authors : De Luca, F.; Benvenuti, M.; Carboni, F.; Pozzi, C.; Rossolini, G.M.; Mangani, S.; Docquier, J.D.
Deposited on : 2011-02-08
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	244	
1	B	244	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CO2	A	270	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4514 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Oxacillinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	242	1941	1246	323	366	6	0	3	0
1	B	243	1950	1249	326	369	6	0	3	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	208	TYR	-	insertion	UNP Q7BNC2
A	209	SER	-	insertion	UNP Q7BNC2
A	210	THR	-	insertion	UNP Q7BNC2
A	211	ARG	-	insertion	UNP Q7BNC2
A	212	ILE	-	insertion	UNP Q7BNC2
A	213	GLU	-	insertion	UNP Q7BNC2
A	217	PRO	-	insertion	UNP Q7BNC2
A	218	LYS	-	insertion	UNP Q7BNC2
A	219	ILE	-	insertion	UNP Q7BNC2
A	220	GLY	-	insertion	UNP Q7BNC2
B	208	TYR	-	insertion	UNP Q7BNC2
B	209	SER	-	insertion	UNP Q7BNC2
B	210	THR	-	insertion	UNP Q7BNC2
B	211	ARG	-	insertion	UNP Q7BNC2
B	212	ILE	-	insertion	UNP Q7BNC2
B	213	GLU	-	insertion	UNP Q7BNC2
B	217	PRO	-	insertion	UNP Q7BNC2
B	218	LYS	-	insertion	UNP Q7BNC2
B	219	ILE	-	insertion	UNP Q7BNC2
B	220	GLY	-	insertion	UNP Q7BNC2

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0

- Molecule 4 is CARBON DIOXIDE (three-letter code: CO2) (formula: CO₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 3 1 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	280	Total O 280 280	0	0
5	B	263	Total O 263 263	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Oxacillinase



- Molecule 1: Oxacillinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.21Å 97.01Å 125.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.83 – 1.60 24.83 – 1.60	Depositor EDS
% Data completeness (in resolution range)	98.1 (24.83-1.60) 98.1 (24.83-1.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.48 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.174 , 0.201 0.173 , 0.200	Depositor DCC
R_{free} test set	3983 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	16.9	Xtrriage
Anisotropy	0.069	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 53.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4514	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: KCX, EDO, CO2, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.59	0/1978	0.67	0/2671
1	B	0.52	0/1987	0.61	0/2682
All	All	0.56	0/3965	0.64	0/5353

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1941	0	1950	11	0
1	B	1950	0	1955	9	0
2	A	30	0	0	1	0
2	B	15	0	0	0	0
3	A	24	0	36	1	0
3	B	8	0	12	3	0
4	A	3	0	0	3	0
5	A	280	0	0	4	0
5	B	263	0	0	1	0
All	All	4514	0	3953	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:58:ARG:HH11	3:B:5:EDO:H22	1.00	1.15
1:B:58:ARG:NH1	3:B:5:EDO:H22	1.71	1.05
1:B:58:ARG:HH11	3:B:5:EDO:C2	1.90	0.81
1:A:44:CYS:SG	1:A:167[B]:VAL:HG11	2.42	0.59
1:A:91:LYS:HE2	5:A:495:HOH:O	2.10	0.52
1:A:61:LYS:NZ	4:A:270:CO2:O1	2.40	0.51
1:A:129:GLU:HG3	1:A:150:ILE:HD13	1.94	0.49
1:A:63:TYR:HH	4:A:270:CO2:C	2.28	0.46
1:A:49:LYS:HG2	2:A:5:SO4:O1	2.15	0.45
3:A:3:EDO:H22	5:A:538:HOH:O	2.16	0.44
1:B:142:GLY:H	1:B:165:ASN:ND2	2.16	0.43
1:A:167[B]:VAL:HG13	1:A:234:PHE:CB	2.49	0.42
1:A:240:ASP:OD1	4:A:270:CO2:O2	2.38	0.42
1:B:30:LYS:NZ	5:B:441:HOH:O	2.53	0.42
1:B:142:GLY:H	1:B:165:ASN:HD21	1.68	0.41
1:B:189:LYS:O	1:B:203:HIS:HD2	2.03	0.41
1:A:138:LYS:HE3	5:A:283:HOH:O	2.20	0.41
1:B:117:VAL:N	1:B:118:PRO:HD2	2.36	0.41
1:B:220:GLY:O	1:B:238:ASN:HA	2.21	0.40
1:A:29:ASN:CG	5:A:507:HOH:O	2.59	0.40
1:A:220:GLY:O	1:A:238:ASN:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	242/244 (99%)	235 (97%)	7 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	243/244 (100%)	239 (98%)	4 (2%)	0	100	100
All	All	485/488 (99%)	474 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	211/209 (101%)	209 (99%)	2 (1%)	78	65
1	B	212/209 (101%)	210 (99%)	2 (1%)	78	65
All	All	423/418 (101%)	419 (99%)	4 (1%)	88	65

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115[A]	SER
1	A	115[B]	SER
1	B	115[A]	SER
1	B	115[B]	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	165	ASN
1	B	101	GLN
1	B	165	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KCX	B	70	1	9,11,12	1.01	0	5,12,14	1.41	1 (20%)
1	KCX	A	70	1	9,11,12	0.83	0	5,12,14	1.50	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	B	70	1	-	0/9/10/12	-
1	KCX	A	70	1	-	0/9/10/12	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	70	KCX	CE-NZ-CX	2.60	126.05	121.89
1	A	70	KCX	OQ1-CX-NZ	-2.48	121.11	124.96
1	A	70	KCX	CE-NZ-CX	2.10	125.25	121.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	7	-	4,4,4	0.13	0	6,6,6	0.11	0
2	SO4	A	9	-	4,4,4	0.14	0	6,6,6	0.05	0
3	EDO	A	269	-	3,3,3	0.48	0	2,2,2	0.49	0
2	SO4	A	5	-	4,4,4	0.14	0	6,6,6	0.11	0
3	EDO	A	2	-	3,3,3	0.43	0	2,2,2	0.43	0
2	SO4	B	6	-	4,4,4	0.14	0	6,6,6	0.23	0
3	EDO	A	3	-	3,3,3	0.34	0	2,2,2	0.84	0
3	EDO	A	267	-	3,3,3	0.44	0	2,2,2	0.34	0
3	EDO	A	6	-	3,3,3	0.52	0	2,2,2	0.25	0
4	CO2	A	270	-	2,2,2	0.79	0	1,1,1	0.35	0
2	SO4	A	8	-	4,4,4	0.28	0	6,6,6	0.37	0
2	SO4	A	4	-	4,4,4	0.14	0	6,6,6	0.06	0
2	SO4	B	2	-	4,4,4	0.18	0	6,6,6	0.25	0
3	EDO	A	268	-	3,3,3	0.39	0	2,2,2	0.46	0
2	SO4	A	1	-	4,4,4	0.22	0	6,6,6	0.42	0
2	SO4	B	3	-	4,4,4	0.14	0	6,6,6	0.17	0
3	EDO	B	5	-	3,3,3	0.54	0	2,2,2	0.48	0
3	EDO	B	4	-	3,3,3	0.34	0	2,2,2	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	269	-	-	1/1/1/1	-
3	EDO	A	6	-	-	1/1/1/1	-
3	EDO	A	2	-	-	1/1/1/1	-
3	EDO	A	3	-	-	1/1/1/1	-
3	EDO	A	267	-	-	1/1/1/1	-
3	EDO	A	268	-	-	0/1/1/1	-
3	EDO	B	5	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	4	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	2	EDO	O1-C1-C2-O2
3	B	4	EDO	O1-C1-C2-O2
3	A	3	EDO	O1-C1-C2-O2
3	A	269	EDO	O1-C1-C2-O2
3	A	267	EDO	O1-C1-C2-O2
3	A	6	EDO	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	5	SO4	1	0
3	A	3	EDO	1	0
4	A	270	CO2	3	0
3	B	5	EDO	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	241/244 (98%)	-0.03	9 (3%) 41 39	8, 15, 33, 41	0
1	B	242/244 (99%)	0.02	10 (4%) 37 34	10, 20, 34, 41	0
All	All	483/488 (98%)	-0.00	19 (3%) 39 36	8, 18, 34, 41	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	26	THR	6.8
1	B	265	GLY	5.5
1	B	212	ILE	5.4
1	B	100	LYS	4.0
1	B	96	PRO	3.6
1	B	211	ARG	3.4
1	A	29	ASN	3.2
1	B	95	LYS	3.1
1	A	30	LYS	2.9
1	B	26	THR	2.9
1	A	49	LYS	2.7
1	A	243	ASN	2.7
1	B	93	ASP	2.6
1	A	263	ILE	2.4
1	A	199	GLU	2.2
1	B	27	SER	2.2
1	A	244	GLU	2.1
1	B	92	TRP	2.1
1	A	198	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	B	70	12/13	0.96	0.09	14,15,21,23	0
1	KCX	A	70	12/13	0.98	0.10	7,8,11,15	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	9	5/5	0.69	0.27	99,99,99,99	0
2	SO4	A	7	5/5	0.72	0.24	80,80,80,80	0
3	EDO	A	2	4/4	0.75	0.23	31,34,35,36	0
3	EDO	A	269	4/4	0.77	0.14	39,41,41,42	0
3	EDO	B	4	4/4	0.84	0.17	42,42,43,44	0
2	SO4	A	4	5/5	0.85	0.21	83,83,83,83	0
3	EDO	A	268	4/4	0.85	0.22	40,41,41,42	0
3	EDO	B	5	4/4	0.85	0.14	28,28,29,31	0
3	EDO	A	3	4/4	0.86	0.18	32,39,40,42	0
3	EDO	A	267	4/4	0.88	0.12	33,34,34,34	0
3	EDO	A	6	4/4	0.88	0.23	23,29,31,32	0
2	SO4	A	5	5/5	0.93	0.30	63,63,64,64	0
2	SO4	B	6	5/5	0.93	0.19	61,62,62,63	0
4	CO2	A	270	3/3	0.94	0.22	33,33,34,34	0
2	SO4	B	3	5/5	0.95	0.12	43,44,45,46	0
2	SO4	A	8	5/5	0.97	0.11	29,31,34,35	0
2	SO4	B	2	5/5	0.98	0.09	28,30,30,32	0
2	SO4	A	1	5/5	0.98	0.09	20,23,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.