



wwPDB X-ray Structure Validation Summary Report ⓘ

May 13, 2020 – 01:52 am BST

PDB ID : 1SS4
Title : Crystal Structure of the Glyoxalase Family Protein APC24694 from *Bacillus cereus*
Authors : Kim, Y.; Li, H.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2004-03-23
Resolution : 1.84 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4003 (1.86-1.82)
Clashscore	141614	4233 (1.86-1.82)
Ramachandran outliers	138981	4185 (1.86-1.82)
Sidechain outliers	138945	4186 (1.86-1.82)
RSRZ outliers	127900	3957 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	153	 8% (Poor fit) 80% (0-1 outliers) 16% (2-3 outliers) 5% (4+ outliers) 5% (Not modelled)
1	B	153	 5% (Poor fit) 81% (0-1 outliers) 14% (2-3 outliers) 5% (4+ outliers) 5% (Not modelled)

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 2596 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glyoxalase family protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	149	1148	721	198	221	2	6	0	0	0
1	B	145	1121	704	193	217	2	5	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	CLONING ARTIFACT	UNP Q81F54
A	-1	ASN	-	CLONING ARTIFACT	UNP Q81F54
A	0	ALA	-	CLONING ARTIFACT	UNP Q81F54
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q81F54
A	9	MSE	MET	MODIFIED RESIDUE	UNP Q81F54
A	59	MSE	MET	MODIFIED RESIDUE	UNP Q81F54
A	60	MSE	MET	MODIFIED RESIDUE	UNP Q81F54
A	97	MSE	MET	MODIFIED RESIDUE	UNP Q81F54
A	106	MSE	MET	MODIFIED RESIDUE	UNP Q81F54
B	-2	SER	-	CLONING ARTIFACT	UNP Q81F54
B	-1	ASN	-	CLONING ARTIFACT	UNP Q81F54
B	0	ALA	-	CLONING ARTIFACT	UNP Q81F54
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q81F54
B	9	MSE	MET	MODIFIED RESIDUE	UNP Q81F54
B	59	MSE	MET	MODIFIED RESIDUE	UNP Q81F54
B	60	MSE	MET	MODIFIED RESIDUE	UNP Q81F54
B	97	MSE	MET	MODIFIED RESIDUE	UNP Q81F54
B	106	MSE	MET	MODIFIED RESIDUE	UNP Q81F54

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Na 1 1	0	0

- Molecule 4 is CITRIC ACID (three-letter code: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	6	7		

- Molecule 5 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			3	1	2		
5	B	1	Total	C	O	0	0
			3	1	2		
5	B	1	Total	C	O	0	0
			3	1	2		
5	B	1	Total	C	O	0	0
			3	1	2		

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Mg	0	0
			1	1		

- Molecule 7 is GLUTATHIONE (three-letter code: GSH) (formula: C₁₀H₁₇N₃O₆S).

4 Data and refinement statistics

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	60.24Å 66.09Å 102.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.57 – 1.84 38.97 – 1.84	Depositor EDS
% Data completeness (in resolution range)	90.4 (33.57-1.84) 90.4 (38.97-1.84)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.05 (at 1.84Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.202 , 0.238 0.198 , 0.234	Depositor DCC
R_{free} test set	3233 reflections (9.19%)	wwPDB-VP
Wilson B-factor (Å ²)	21.8	Xtrriage
Anisotropy	0.801	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 56.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2596	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, NA, FMT, ACT, CIT, GSH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/1158	0.72	0/1556
1	B	0.47	0/1132	0.70	0/1524
All	All	0.46	0/2290	0.71	0/3080

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1148	0	1151	28	0
1	B	1121	0	1116	21	0
2	A	4	0	3	0	0
3	A	1	0	0	0	0
4	A	13	0	5	0	0
5	A	3	0	1	0	0
5	B	9	0	3	1	0
6	B	1	0	0	0	0
7	B	20	0	15	2	0
8	A	132	0	0	5	2
8	B	144	0	0	6	0
All	All	2596	0	2294	44	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 44 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:96:VAL:HG12	1:B:12:VAL:HG22	1.62	0.82
1:A:117:LEU:HA	1:A:133:ILE:HD13	1.67	0.77
1:B:111:THR:HG21	8:B:548:HOH:O	1.87	0.73
1:A:116:GLU:HG2	1:A:134:ARG:HB2	1.78	0.67
1:A:42:GLU:OE1	1:A:46:ARG:NH1	2.30	0.65

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:560:HOH:O	8:A:560:HOH:O[3_555]	2.01	0.19
8:A:533:HOH:O	8:A:533:HOH:O[3_655]	2.03	0.17

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	147/153 (96%)	144 (98%)	3 (2%)	0	100	100
1	B	143/153 (94%)	141 (99%)	2 (1%)	0	100	100
All	All	290/306 (95%)	285 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/122 (102%)	121 (98%)	3 (2%)	49	32
1	B	122/122 (100%)	120 (98%)	2 (2%)	62	49
All	All	246/244 (101%)	241 (98%)	5 (2%)	55	40

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	ARG
1	A	95	ARG
1	A	146	GLU
1	B	42	GLU
1	B	95	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	126	ASN
1	B	53	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ACT	A	411	-	1,3,3	2.55	1 (100%)	0,3,3	0.00	-
5	FMT	B	404	-	0,2,2	0.00	-	0,1,1	0.00	-
4	CIT	A	431	-	3,12,12	2.08	1 (33%)	3,17,17	0.48	0
7	GSH	B	432	-	12,19,19	1.76	5 (41%)	15,24,24	1.38	3 (20%)
5	FMT	B	402	-	0,2,2	0.00	-	0,1,1	0.00	-
5	FMT	B	401	-	0,2,2	0.00	-	0,1,1	0.00	-
5	FMT	A	403	-	0,2,2	0.00	-	0,1,1	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CIT	A	431	-	-	3/6/16/16	-
7	GSH	B	432	-	-	2/18/24/24	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	431	CIT	C4-C3	3.44	1.59	1.54
2	A	411	ACT	CH3-C	2.55	1.52	1.48
7	B	432	GSH	CA2-C2	2.47	1.59	1.52
7	B	432	GSH	CA3-N3	2.39	1.50	1.46
7	B	432	GSH	CA2-N2	2.16	1.50	1.45

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	432	GSH	CG1-CB1-CA1	2.38	119.40	113.84
7	B	432	GSH	CB2-CA2-N2	-2.15	108.22	111.28
7	B	432	GSH	CA3-N3-C2	-2.08	119.35	122.34

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	431	CIT	C2-C3-C4-C5
4	A	431	CIT	O7-C3-C4-C5
4	A	431	CIT	C6-C3-C4-C5
7	B	432	GSH	OE1-CD1-CG1-CB1
7	B	432	GSH	N2-CD1-CG1-CB1

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	432	GSH	2	0
5	B	402	FMT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	143/153 (93%)	0.37	12 (8%) 11 10	17, 30, 51, 59	0
1	B	140/153 (91%)	0.22	8 (5%) 23 21	18, 27, 49, 61	0
All	All	283/306 (92%)	0.30	20 (7%) 16 14	17, 29, 50, 61	0

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ALA	4.7
1	A	0	ALA	4.2
1	A	126	ASN	3.9
1	A	128	TYR	3.7
1	B	40	GLU	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	CIT	A	431	13/13	0.63	0.20	57,58,59,59	0
7	GSH	B	432	20/20	0.73	0.24	52,58,61,62	0
2	ACT	A	411	4/4	0.86	0.20	50,52,54,54	0
5	FMT	B	404	3/3	0.90	0.19	50,50,51,54	0
5	FMT	B	402	3/3	0.92	0.14	32,32,33,33	0
6	MG	B	302	1/1	0.94	0.30	14,14,14,14	1
5	FMT	A	403	3/3	0.94	0.24	40,40,41,41	0
5	FMT	B	401	3/3	0.98	0.09	23,23,23,28	0
3	NA	A	301	1/1	1.00	0.15	11,11,11,11	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.