



# Full wwPDB X-ray Structure Validation Report ⓘ

May 28, 2020 – 09:45 pm BST

PDB ID : 6TWB  
Title : Crystal Structure of the Catalytic Domain of Coagulation Factor XIa in Complex with Double Bridged Peptide F19  
Authors : Kong, X.D.; Pojer, F.; Heinis, C.  
Deposited on : 2020-01-13  
Resolution : 2.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2307 (2.94-2.90)
Clashscore	141614	2531 (2.94-2.90)
Ramachandran outliers	138981	2462 (2.94-2.90)
Sidechain outliers	138945	2464 (2.94-2.90)
RSRZ outliers	127900	2248 (2.94-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	262	 2% 65% 25% 9%
1	H	262	 97%
2	B	11	 18% 45% 18% 18%

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 2048 atoms, of which 12 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Coagulation factor XI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	238	1880	1188	333	348	11	0	0	0
1	H	8	62	39	10	12	1	0	0	0

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	initiating methionine	UNP P03951
A	-1	ASP	-	expression tag	UNP P03951
A	0	ASP	-	expression tag	UNP P03951
A	1	ASP	-	expression tag	UNP P03951
A	2	ASP	-	expression tag	UNP P03951
A	113	GLY	ASN	conflict	UNP P03951
A	115	GLY	THR	conflict	UNP P03951
A	246	HIS	-	expression tag	UNP P03951
A	247	HIS	-	expression tag	UNP P03951
A	248	HIS	-	expression tag	UNP P03951
A	249	HIS	-	expression tag	UNP P03951
A	250	HIS	-	expression tag	UNP P03951
A	251	HIS	-	expression tag	UNP P03951
H	352	MET	-	initiating methionine	UNP P03951
H	353	ASP	-	expression tag	UNP P03951
H	354	ASP	-	expression tag	UNP P03951
H	355	ASP	-	expression tag	UNP P03951
H	356	ASP	-	expression tag	UNP P03951
H	473	GLY	ASN	conflict	UNP P03951
H	475	GLY	THR	conflict	UNP P03951
H	608	HIS	-	expression tag	UNP P03951
H	609	HIS	-	expression tag	UNP P03951
H	610	HIS	-	expression tag	UNP P03951
H	611	HIS	-	expression tag	UNP P03951
H	612	HIS	-	expression tag	UNP P03951

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	613	HIS	-	expression tag	UNP P03951

- Molecule 2 is a protein called Double Bridged Peptide F19.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	11	81	49	14	13	5	0	0	1

- Molecule 3 is AMMONIUM ION (three-letter code: NH4) (formula: H<sub>4</sub>N).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	H	N		
3	A	1	5	4	1	0	0
3	A	1	5	4	1	0	0
3	B	1	5	4	1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	9	9	9	0	0
4	B	1	1	1	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.52Å 77.52Å 116.51Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.00 – 2.91 44.00 – 2.91	Depositor EDS
% Data completeness (in resolution range)	99.7 (44.00-2.91) 99.9 (44.00-2.91)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.32 (at 2.90Å)	Xtriage
Refinement program	PHENIX dev-3374	Depositor
R, $R_{free}$	0.179 , 0.241 0.179 , 0.240	Depositor DCC
$R_{free}$ test set	429 reflections (4.59%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	51.4	Xtriage
Anisotropy	0.805	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 48.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.034 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2048	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.59% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NH4, CSA, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/1925	0.60	0/2608
1	H	0.60	0/62	0.64	0/82
2	B	3.93	12/60 (20.0%)	1.75	2/78 (2.6%)
All	All	0.82	12/2047 (0.6%)	0.66	2/2768 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	1	0

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	8	ARG	CZ-NH1	12.67	1.49	1.33
2	B	8	ARG	NE-CZ	11.18	1.47	1.33
2	B	10	PRO	N-CD	10.93	1.63	1.47
2	B	9	CYS	C-N	8.72	1.50	1.34
2	B	8	ARG	CZ-NH2	-7.46	1.23	1.33
2	B	10	PRO	N-CA	-7.34	1.34	1.47
2	B	4	ILE	C-N	7.07	1.50	1.34
2	B	3	ASN	C-N	6.75	1.49	1.34
2	B	8	ARG	C-N	6.41	1.48	1.34
2	B	7	CYS	C-N	5.96	1.47	1.34
2	B	2	VAL	C-N	5.49	1.46	1.34
2	B	3	ASN	CG-ND2	5.49	1.46	1.32

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	B	8	ARG	CD-NE-CZ	-6.47	114.54	123.60
2	B	5	MET	CG-SD-CE	5.97	109.75	100.20

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	10	PRO	CA

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1880	0	1842	47	0
1	H	62	0	67	4	0
2	B	81	0	77	5	0
3	A	2	8	0	0	0
3	B	1	4	0	0	0
4	A	9	0	0	1	0
4	B	1	0	0	0	0
All	All	2036	12	1986	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance ( $\text{\AA}$ )	Clash overlap ( $\text{\AA}$ )
1:A:32:THR:HG23	1:A:67:TYR:HB2	1.65	0.79
1:A:102:ASP:OD2	1:A:214:SER:HB2	1.93	0.69
1:A:213:THR:HA	1:A:228:TYR:CD2	2.36	0.60
1:A:98:GLU:OE1	2:B:3:ASN:ND2	2.34	0.58
1:A:35:THR:O	1:A:38:GLN:HA	2.04	0.58
1:A:35:THR:OG1	1:A:65:ILE:HD12	2.06	0.55
1:A:57:HIS:NE2	1:A:195:SER:HB2	2.23	0.54
1:A:167:GLU:HG2	1:A:170:ARG:HH21	1.75	0.52

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:LYS:HG3	1:A:78:GLU:H	1.75	0.51
1:A:51:TRP:CZ3	1:A:107:LYS:HB2	2.45	0.51
1:A:148:ARG:HH22	1:A:192:LYS:HE3	1.76	0.50
1:A:73:GLN:HG2	1:A:141:TRP:CD1	2.47	0.50
1:H:362:CYS:O	1:H:365:LYS:HE2	2.11	0.50
1:A:121:ILE:HD13	1:A:209:LEU:HB2	1.94	0.49
2:B:4:ILE:O	2:B:4:ILE:HD12	2.12	0.49
1:A:65(B):ARG:HG2	1:A:82:PHE:CE1	2.48	0.49
1:A:56:ALA:HB1	1:A:90:ILE:HG23	1.93	0.49
1:A:214:SER:O	2:B:6:CSA:H12	2.12	0.49
1:A:132:ILE:HG23	1:A:132:ILE:O	2.13	0.48
1:A:84:GLY:C	1:A:109:GLU:HG3	2.33	0.48
1:A:77:LYS:HA	1:A:77:LYS:HD2	1.73	0.47
1:A:149:ASP:OD2	1:A:150:LYS:N	2.45	0.47
1:A:49:ASN:OD1	1:H:368:PRO:HA	2.15	0.47
1:A:148:ARG:HA	1:A:148:ARG:NE	2.28	0.47
1:A:187:GLY:H	1:A:220:ALA:HB1	1.78	0.47
1:A:166:GLU:N	1:A:166:GLU:OE2	2.44	0.46
2:B:9:CYS:HB3	2:B:10:PRO:HD2	1.98	0.46
1:A:52:ILE:HG13	1:A:108:LEU:HD21	1.98	0.45
1:A:192:LYS:NZ	2:B:10:PRO:O	2.49	0.45
1:A:58(A):PHE:CZ	1:A:106:LEU:HD21	2.51	0.45
1:A:170:ARG:NE	1:A:223:GLU:OE1	2.47	0.45
1:A:49:ASN:ND2	1:H:368:PRO:HB3	2.32	0.45
1:A:83:PHE:CD1	1:A:110:THR:HG23	2.52	0.44
1:A:170:ARG:NH1	1:A:223:GLU:OE1	2.50	0.44
1:A:145:ARG:HD3	1:A:152:GLN:OE1	2.18	0.44
1:A:135:ASP:O	1:A:201:CYS:HA	2.17	0.44
1:A:83:PHE:HD1	1:A:110:THR:HG23	1.83	0.44
1:A:156:GLN:HG3	4:A:405:HOH:O	2.18	0.43
1:A:45:SER:OG	1:A:198:PRO:HB3	2.19	0.43
1:A:33:LEU:HD13	1:A:66:VAL:HG22	2.01	0.43
1:A:214:SER:OG	1:A:215:TRP:HD1	2.02	0.42
1:A:29:TRP:CE2	1:A:121:ILE:HD12	2.55	0.42
1:A:148:ARG:HH22	1:A:192:LYS:CE	2.32	0.42
1:A:50:GLN:HG3	1:A:51:TRP:CD1	2.55	0.42
1:A:16:ILE:N	1:A:194:ASP:OD2	2.53	0.41
1:A:182:CYS:HA	1:A:226:GLY:O	2.20	0.41
1:A:121:ILE:HG12	1:A:122:CYS:N	2.35	0.41
1:A:17:VAL:O	1:A:188:LYS:HA	2.21	0.41
1:A:57:HIS:CE1	1:A:195:SER:HB2	2.56	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:367:LYS:HA	1:H:368:PRO:HD3	1.70	0.41
1:A:162:LEU:HD21	1:A:199:LEU:HD21	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	236/262 (90%)	224 (95%)	12 (5%)	0	100	100
1	H	6/262 (2%)	6 (100%)	0	0	100	100
2	B	8/11 (73%)	6 (75%)	1 (12%)	1 (12%)	0	0
All	All	250/535 (47%)	236 (94%)	13 (5%)	1 (0%)	34	65

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	10	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/226 (89%)	195 (96%)	7 (4%)	36	68
1	H	8/226 (4%)	8 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	8/8 (100%)	7 (88%)	1 (12%)	4	13
All	All	218/460 (47%)	210 (96%)	8 (4%)	34	66

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	81	SER
1	A	126	LYS
1	A	148	ARG
1	A	170	ARG
1	A	172	ARG
1	A	184(B)	ARG
1	A	214	SER
2	B	4	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	CSA	B	1	2	8,8,10	1.14	1 (12%)	5,8,12	2.05	1 (20%)
2	CSA	B	6	2	8,9,10	1.80	1 (12%)	4,10,12	1.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSA	B	1	2	-	2/5/6/10	-
2	CSA	B	6	2	-	2/6/8/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	6	CSA	C3-C2	4.28	1.58	1.51
2	B	1	CSA	C3-C2	2.07	1.54	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	CSA	CB-SG-C3	4.00	108.40	101.71

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	CSA	C1-C2-C3-SG
2	B	1	CSA	CA-CB-SG-C3
2	B	6	CSA	CA-CB-SG-C3
2	B	6	CSA	C1-C2-C3-SG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	6	CSA	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are modelled with single atom - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	238/262 (90%)	-0.27	5 (2%) 63 62	33, 51, 85, 109	0
1	H	8/262 (3%)	-0.57	0 100 100	63, 72, 89, 92	0
2	B	8/11 (72%)	-0.47	0 100 100	45, 51, 57, 62	0
All	All	254/535 (47%)	-0.28	5 (1%) 65 64	33, 51, 88, 109	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	130	ASN	4.5
1	A	61	GLU	3.4
1	A	129	ARG	2.6
1	A	131	VAL	2.4
1	A	132	ILE	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CSA	B	1	9/11	0.96	0.12	57,63,79,82	0
2	CSA	B	6	10/11	0.98	0.12	39,51,55,57	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NH4	A	302	1/1	0.75	0.17	45,54,54,54	0
3	NH4	A	301	1/1	0.84	0.51	53,64,64,64	0
3	NH4	B	101	1/1	0.89	0.36	55,66,66,66	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.