



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2023 – 10:08 PM EDT

PDB ID : 1V82  
Title : Crystal structure of human GlcAT-P apo form  
Authors : Kakuda, S.; Shiba, T.; Ishiguro, M.; Tagawa, H.; Oka, S.; Kajihara, Y.;  
Kawasaki, T.; Wakatsuki, S.; Kato, R.  
Deposited on : 2003-12-27  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	253	
1	B	253	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4343 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	247	Total 2005	C 1277	N 373	O 350	S 5	0	0	0
1	B	241	Total 1947	C 1241	N 359	O 342	S 5	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	82	ALA	-	cloning artifact	UNP Q9P2W7
B	82	ALA	-	cloning artifact	UNP Q9P2W7

- Molecule 2 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: C<sub>4</sub>H<sub>6</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 10	C 4	O 6	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	159	Total 159	O 159	0	0
3	B	222	Total 222	O 222	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 1



- Molecule 1: Galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.11Å 85.97Å 122.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 1.85 49.86 – 1.85	Depositor EDS
% Data completeness (in resolution range)	96.3 (40.00-1.85) 96.5 (49.86-1.85)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.93 (at 1.86Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.198 , 0.228 0.191 , 0.190	Depositor DCC
$R_{free}$ test set	2774 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.8	Xtrriage
Anisotropy	0.925	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 51.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4343	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.72% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.31	0/2054	0.58	0/2790
1	B	0.31	0/1995	0.57	0/2713
All	All	0.31	0/4049	0.58	0/5503

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2005	0	2039	49	0
1	B	1947	0	1976	45	0
2	A	10	0	4	0	0
3	A	159	0	0	2	0
3	B	222	0	0	2	0
All	All	4343	0	4019	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:171:ASN:HD21	1:A:273:PHE:H	1.25	0.83
1:B:171:ASN:HD21	1:B:273:PHE:H	1.25	0.80
1:A:315:GLU:HG2	1:B:315:GLU:HG3	1.65	0.79
1:B:261:ASN:HD22	1:B:263:ARG:H	1.28	0.78
1:B:274:LYS:HB3	1:B:278:VAL:HG21	1.68	0.75
1:A:163:ILE:HD11	1:A:276:ARG:HG2	1.73	0.69
1:B:182:PRO:HD3	3:B:412:HOH:O	1.92	0.68
1:B:231:ARG:HE	1:B:240:ARG:HE	1.39	0.68
1:A:111:HIS:HE1	1:B:107:ASN:O	1.77	0.68
1:B:261:ASN:ND2	1:B:263:ARG:H	1.93	0.65
1:A:240:ARG:HG2	1:A:240:ARG:HH21	1.60	0.65
1:A:231:ARG:NH2	1:A:240:ARG:CB	2.60	0.65
1:B:212:ARG:HD2	3:B:536:HOH:O	1.98	0.64
1:B:182:PRO:HG2	1:B:185:SER:HB3	1.80	0.64
1:A:263:ARG:HG2	1:A:267:GLN:HE21	1.63	0.64
1:B:187:GLN:H	1:B:187:GLN:NE2	1.96	0.64
1:A:292:VAL:CG2	1:A:296:ASP:HB2	2.28	0.63
1:B:150:ARG:HD3	1:B:150:ARG:N	2.14	0.62
1:B:246:ASP:O	1:B:249:ARG:HG2	1.99	0.62
1:A:233:ASN:C	1:A:233:ASN:HD22	2.03	0.61
1:A:218:PRO:HB2	1:A:307:ILE:HD13	1.83	0.60
1:A:274:LYS:HE2	1:A:276:ARG:O	2.01	0.60
1:A:133:LEU:O	1:A:137:THR:HG23	2.02	0.60
1:A:165:ARG:HB2	1:A:165:ARG:HH21	1.65	0.60
1:A:307:ILE:HG13	1:B:334:ILE:HG22	1.83	0.60
1:A:312:THR:C	1:A:313:ARG:HD2	2.23	0.59
1:A:292:VAL:HG22	1:A:293:THR:H	1.68	0.58
1:A:116:HIS:HE1	1:A:142:THR:OG1	1.86	0.58
1:A:292:VAL:HG22	1:A:296:ASP:HB2	1.85	0.57
1:A:240:ARG:HG2	1:A:240:ARG:NH2	2.20	0.56
1:B:261:ASN:HD22	1:B:263:ARG:N	2.03	0.55
1:B:149:PRO:HD2	1:B:168:MET:SD	2.48	0.54
1:A:312:THR:O	1:A:313:ARG:HD2	2.08	0.53
1:A:292:VAL:HG22	1:A:293:THR:N	2.22	0.53
1:A:231:ARG:NH2	1:A:240:ARG:HB2	2.24	0.53
1:A:155:ARG:N	1:A:155:ARG:HD3	2.24	0.52
1:A:162:ARG:HB3	3:A:646:HOH:O	2.09	0.51
1:B:87:HIS:ND1	1:B:116:HIS:HD2	2.08	0.51
1:A:263:ARG:HG2	1:A:267:GLN:NE2	2.26	0.51
1:B:231:ARG:HH21	1:B:240:ARG:HH21	1.58	0.51
1:B:163:ILE:HD12	1:B:163:ILE:O	2.11	0.51
1:B:146:VAL:HB	1:B:172:LEU:HD22	1.94	0.50

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:311:HIS:ND1	1:A:313:ARG:HD3	2.26	0.50
1:A:329:ASP:HB3	1:A:332:VAL:HG23	1.93	0.50
1:A:87:HIS:ND1	1:A:116:HIS:HD2	2.09	0.50
1:B:150:ARG:HD3	1:B:150:ARG:H	1.76	0.50
1:B:167:THR:HG21	1:B:275:LEU:HB3	1.93	0.50
1:B:222:VAL:HG23	1:B:228:GLU:HG3	1.93	0.49
1:B:163:ILE:HG22	1:B:275:LEU:O	2.12	0.49
1:B:275:LEU:H	1:B:275:LEU:HD23	1.76	0.49
1:B:129:LEU:C	1:B:129:LEU:HD23	2.32	0.49
1:B:116:HIS:HE1	1:B:142:THR:OG1	1.95	0.49
1:B:275:LEU:HD23	1:B:275:LEU:N	2.28	0.48
1:A:163:ILE:HD13	1:A:276:ARG:HA	1.94	0.48
1:B:133:LEU:O	1:B:137:THR:HG23	2.13	0.48
1:B:198:ASN:HB2	1:B:200:TYR:CZ	2.47	0.48
1:B:231:ARG:NE	1:B:240:ARG:HE	2.10	0.48
1:A:110:LEU:HA	1:A:139:LEU:HD21	1.96	0.48
1:A:125:ARG:HD3	3:A:536:HOH:O	2.13	0.47
1:A:231:ARG:CZ	1:A:240:ARG:HB2	2.44	0.47
1:B:175:ARG:O	1:B:179:GLU:HG3	2.13	0.47
1:B:292:VAL:HG22	1:B:293:THR:N	2.29	0.47
1:A:163:ILE:CD1	1:A:276:ARG:HG2	2.40	0.47
1:A:129:LEU:HD23	1:A:129:LEU:C	2.35	0.47
1:A:185:SER:C	1:A:187:GLN:H	2.19	0.46
1:A:233:ASN:C	1:A:233:ASN:ND2	2.69	0.46
1:B:166:GLY:HA2	1:B:169:GLN:HE21	1.81	0.46
1:B:231:ARG:HG3	1:B:242:LYS:HZ2	1.81	0.45
1:B:169:GLN:NE2	1:B:169:GLN:H	2.15	0.45
1:A:116:HIS:CE1	1:A:142:THR:OG1	2.69	0.45
1:A:97:VAL:O	1:A:101:GLU:HG2	2.17	0.44
1:A:146:VAL:HB	1:A:172:LEU:HD22	2.00	0.44
1:A:154:LEU:C	1:A:155:ARG:HD3	2.38	0.43
1:A:163:ILE:CD1	1:A:276:ARG:HA	2.48	0.43
1:A:212:ARG:HG3	1:A:212:ARG:HH11	1.84	0.43
1:A:194:ALA:HB1	1:A:200:TYR:CZ	2.53	0.43
1:B:187:GLN:H	1:B:187:GLN:CD	2.21	0.43
1:B:274:LYS:HB3	1:B:278:VAL:CG2	2.42	0.43
1:B:197:ASP:OD2	1:B:313:ARG:NH1	2.51	0.43
1:A:149:PRO:HD2	1:A:152:TYR:CD1	2.55	0.42
1:B:231:ARG:HE	1:B:240:ARG:NE	2.10	0.42
1:A:182:PRO:HG2	1:A:185:SER:HA	2.00	0.42
1:B:246:ASP:N	1:B:247:PRO:HD3	2.35	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:231:ARG:HG3	1:B:242:LYS:NZ	2.35	0.41
1:A:279:LYS:HE3	1:A:282:TYR:CZ	2.55	0.41
1:B:187:GLN:HA	1:B:188:PRO:HD2	1.94	0.41
1:B:329:ASP:HB3	1:B:332:VAL:HG23	2.01	0.41
1:A:198:ASN:HB2	1:A:200:TYR:CZ	2.56	0.41
1:A:185:SER:O	1:A:186:SER:HB2	2.21	0.41
1:B:239:VAL:O	1:B:240:ARG:HG3	2.21	0.41
1:A:133:LEU:HD23	1:A:133:LEU:C	2.42	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	243/253 (96%)	236 (97%)	6 (2%)	1 (0%)	34	19
1	B	237/253 (94%)	222 (94%)	13 (6%)	2 (1%)	19	7
All	All	480/506 (95%)	458 (95%)	19 (4%)	3 (1%)	25	12

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	276	ARG
1	A	240	ARG
1	B	277	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/222 (98%)	212 (97%)	6 (3%)	43	27
1	B	212/222 (96%)	204 (96%)	8 (4%)	33	16
All	All	430/444 (97%)	416 (97%)	14 (3%)	38	21

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	95	ARG
1	A	155	ARG
1	A	165	ARG
1	A	197	ASP
1	A	233	ASN
1	A	240	ARG
1	B	95	ARG
1	B	109	LEU
1	B	118	LEU
1	B	150	ARG
1	B	187	GLN
1	B	246	ASP
1	B	261	ASN
1	B	275	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	107	ASN
1	A	111	HIS
1	A	116	HIS
1	A	140	ASN
1	A	151	ASN
1	A	171	ASN
1	A	187	GLN
1	A	198	ASN
1	A	233	ASN
1	A	261	ASN
1	A	267	GLN
1	A	283	GLN
1	B	107	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	114	ASN
1	B	116	HIS
1	B	169	GLN
1	B	171	ASN
1	B	187	GLN
1	B	198	ASN
1	B	261	ASN
1	B	267	GLN
1	B	283	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	TLA	A	500	-	9,9,9	1.02	0	12,12,12	1.05	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TLA	A	500	-	-	0/12/12/12	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	A	500	TLA	O41-C4-C3	2.34	119.60	113.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	247/253 (97%)	0.28	20 (8%) <b>12</b> <b>12</b>	12, 20, 41, 59	0
1	B	241/253 (95%)	0.42	22 (9%) <b>9</b> <b>8</b>	12, 20, 45, 63	0
All	All	488/506 (96%)	0.35	42 (8%) <b>10</b> <b>10</b>	12, 20, 44, 63	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	186	SER	11.7
1	B	278	VAL	8.2
1	A	186	SER	7.8
1	B	277	GLY	7.4
1	B	276	ARG	7.4
1	A	185	SER	7.1
1	B	150	ARG	6.7
1	B	275	LEU	6.5
1	B	185	SER	6.0
1	B	184	ASN	5.9
1	B	82	ALA	5.9
1	A	156	GLY	5.7
1	A	155	ARG	5.5
1	A	187	GLN	5.5
1	B	163	ILE	5.5
1	A	324	LYS	4.7
1	B	248	HIS	4.2
1	A	162	ARG	4.0
1	A	325	LYS	4.0
1	B	240	ARG	3.9
1	B	183	ARG	3.6
1	A	184	ASN	3.6
1	B	279	LYS	3.4
1	B	282	TYR	3.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	182	PRO	3.2
1	A	326	GLY	3.1
1	B	324	LYS	2.9
1	A	240	ARG	2.9
1	A	188	PRO	2.8
1	B	125	ARG	2.7
1	A	212	ARG	2.6
1	A	320	VAL	2.5
1	B	280	GLY	2.5
1	B	274	LYS	2.4
1	A	125	ARG	2.3
1	A	263	ARG	2.3
1	A	331	SER	2.2
1	A	327	PHE	2.2
1	A	278	VAL	2.1
1	B	164	PRO	2.1
1	A	183	ARG	2.1
1	B	281	GLY	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	TLA	A	500	10/10	0.95	0.08	19,23,27,28	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.