



Full wwPDB NMR Structure Validation Report ⓘ

Feb 22, 2022 – 09:37 AM EST

PDB ID : 1VA2
Title : Solution Structure of Transcription Factor Sp1 DNA Binding Domain (Zinc Finger 2)
Authors : Oka, S.; Shiraishi, Y.; Yoshida, T.; Ohkubo, T.; Sugiura, Y.; Kobayashi, Y.
Deposited on : 2004-02-07

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.26
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.26


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	31	

2 Ensemble composition and analysis

This entry contains 31 models. Model 11 is the overall representative, medoid model (most similar to other models). The authors have identified model 31 as representative, based on the following criterion: *minimized average structure*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:566-A:591 (26)	0.32	11

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	8, 9, 10, 11, 13, 15, 17, 18, 19, 20, 21, 22, 23, 25, 26, 27, 29, 30
2	1, 2, 3, 4, 6, 16, 24
3	5, 7, 12, 14, 31
Single-model clusters	28

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 530 atoms, of which 260 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Transcription factor Sp1.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	31	529	165	260	55	46	3	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
			Total	Zn
2	A	1	1	1

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Transcription factor Sp1



4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Transcription factor Sp1




4.2.2 Score per residue for model 2

- Molecule 1: Transcription factor Sp1



4.2.3 Score per residue for model 3

- Molecule 1: Transcription factor Sp1




4.2.4 Score per residue for model 4

- Molecule 1: Transcription factor Sp1



4.2.5 Score per residue for model 5

- Molecule 1: Transcription factor Sp1




4.2.6 Score per residue for model 6

- Molecule 1: Transcription factor Sp1




4.2.7 Score per residue for model 7

- Molecule 1: Transcription factor Sp1




4.2.8 Score per residue for model 8

- Molecule 1: Transcription factor Sp1



4.2.9 Score per residue for model 9

- Molecule 1: Transcription factor Sp1




4.2.10 Score per residue for model 10

- Molecule 1: Transcription factor Sp1




4.2.11 Score per residue for model 11 (medoid)

- Molecule 1: Transcription factor Sp1




4.2.12 Score per residue for model 12

- Molecule 1: Transcription factor Sp1




4.2.13 Score per residue for model 13

- Molecule 1: Transcription factor Sp1




4.2.14 Score per residue for model 14

- Molecule 1: Transcription factor Sp1




4.2.15 Score per residue for model 15

- Molecule 1: Transcription factor Sp1




4.2.16 Score per residue for model 16

- Molecule 1: Transcription factor Sp1




4.2.17 Score per residue for model 17

- Molecule 1: Transcription factor Sp1



4.2.18 Score per residue for model 18

- Molecule 1: Transcription factor Sp1




4.2.19 Score per residue for model 19

- Molecule 1: Transcription factor Sp1




4.2.20 Score per residue for model 20

- Molecule 1: Transcription factor Sp1




4.2.21 Score per residue for model 21

- Molecule 1: Transcription factor Sp1




4.2.22 Score per residue for model 22

- Molecule 1: Transcription factor Sp1




4.2.23 Score per residue for model 23

- Molecule 1: Transcription factor Sp1




4.2.24 Score per residue for model 24

- Molecule 1: Transcription factor Sp1




4.2.25 Score per residue for model 25

- Molecule 1: Transcription factor Sp1




4.2.26 Score per residue for model 26

- Molecule 1: Transcription factor Sp1




4.2.27 Score per residue for model 27

- Molecule 1: Transcription factor Sp1




4.2.28 Score per residue for model 28

- Molecule 1: Transcription factor Sp1




4.2.29 Score per residue for model 29

- Molecule 1: Transcription factor Sp1



4.2.30 Score per residue for model 30

- Molecule 1: Transcription factor Sp1



4.2.31 Score per residue for model 31

- Molecule 1: Transcription factor Sp1



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics, simulated annealing*.

Of the 100 calculated structures, 31 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure solution	1.1
CNS	refinement	1.1

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	228	216	216	7±2
All	All	7099	6696	6696	208

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:591:HIS:CG	1:A:591:HIS:O	0.69	2.45	30	1
1:A:584:LEU:HD12	1:A:585:GLN:NE2	0.67	2.04	19	2
1:A:569:CYS:HB2	1:A:584:LEU:HD23	0.63	1.69	28	2
1:A:585:GLN:HG2	1:A:586:ARG:N	0.62	2.07	25	1
1:A:581:SER:HA	1:A:584:LEU:HD12	0.61	1.71	25	5
1:A:567:PHE:CD1	1:A:584:LEU:HD22	0.61	2.30	14	1
1:A:584:LEU:HD12	1:A:585:GLN:HE22	0.60	1.56	19	2
1:A:567:PHE:HB3	1:A:584:LEU:HD11	0.60	1.73	5	4
1:A:569:CYS:HA	1:A:584:LEU:HD23	0.59	1.73	16	3
1:A:567:PHE:CG	1:A:584:LEU:HD22	0.57	2.34	14	1
1:A:570:THR:HG22	1:A:576:LYS:H	0.57	1.58	28	1
1:A:583:GLU:O	1:A:587:HIS:HB3	0.56	2.00	4	25

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:578:PHE:HB2	1:A:584:LEU:HD11	0.56	1.77	3	1
1:A:571:TRP:CZ2	1:A:588:LYS:HB3	0.55	2.36	29	1
1:A:569:CYS:O	1:A:569:CYS:SG	0.55	2.64	28	1
1:A:587:HIS:CE1	1:A:591:HIS:CD2	0.53	2.97	30	2
1:A:569:CYS:SG	1:A:571:TRP:CE2	0.53	3.02	31	1
1:A:586:ARG:O	1:A:589:ARG:HD2	0.52	2.04	24	1
1:A:578:PHE:CD1	1:A:584:LEU:HD21	0.52	2.39	2	1
1:A:571:TRP:CD2	1:A:591:HIS:CD2	0.51	2.99	16	1
1:A:571:TRP:CD1	1:A:591:HIS:CE1	0.51	2.99	10	3
1:A:569:CYS:N	1:A:584:LEU:HD22	0.49	2.22	8	3
1:A:584:LEU:O	1:A:588:LYS:HB3	0.49	2.07	29	12
1:A:581:SER:O	1:A:585:GLN:NE2	0.49	2.46	12	15
1:A:586:ARG:O	1:A:589:ARG:HB3	0.48	2.08	26	2
1:A:569:CYS:CB	1:A:584:LEU:HD23	0.48	2.38	28	1
1:A:584:LEU:HD12	1:A:584:LEU:H	0.48	1.69	16	2
1:A:578:PHE:CD2	1:A:583:GLU:HB3	0.48	2.43	27	7
1:A:586:ARG:O	1:A:589:ARG:HB2	0.48	2.08	23	5
1:A:568:MET:N	1:A:584:LEU:HD11	0.47	2.25	8	1
1:A:569:CYS:CA	1:A:584:LEU:HD22	0.47	2.40	8	1
1:A:581:SER:HA	1:A:584:LEU:HD21	0.46	1.87	14	1
1:A:581:SER:HA	1:A:584:LEU:HD23	0.46	1.85	24	2
1:A:567:PHE:CG	1:A:584:LEU:CD2	0.46	2.98	14	1
1:A:585:GLN:O	1:A:588:LYS:CG	0.46	2.63	23	1
1:A:567:PHE:CB	1:A:584:LEU:HD11	0.46	2.41	5	1
1:A:569:CYS:N	1:A:584:LEU:CD2	0.46	2.79	6	8
1:A:568:MET:C	1:A:584:LEU:HD21	0.46	2.30	21	1
1:A:571:TRP:NE1	1:A:591:HIS:CG	0.46	2.84	10	2
1:A:578:PHE:CD1	1:A:584:LEU:CD2	0.45	2.99	2	2
1:A:584:LEU:O	1:A:588:LYS:CG	0.45	2.64	3	10
1:A:571:TRP:CZ2	1:A:588:LYS:HB2	0.45	2.46	27	5
1:A:586:ARG:O	1:A:589:ARG:CD	0.45	2.65	26	1
1:A:587:HIS:CD2	1:A:588:LYS:N	0.44	2.85	19	3
1:A:584:LEU:O	1:A:588:LYS:HG2	0.44	2.12	22	3
1:A:571:TRP:CD1	1:A:591:HIS:CG	0.44	3.06	26	1
1:A:584:LEU:O	1:A:588:LYS:HB2	0.44	2.12	2	5
1:A:588:LYS:HG3	1:A:589:ARG:N	0.44	2.28	20	3
1:A:571:TRP:NE1	1:A:574:CYS:HB2	0.43	2.28	28	1
1:A:569:CYS:SG	1:A:571:TRP:CH2	0.43	3.11	28	1
1:A:586:ARG:HG3	1:A:587:HIS:N	0.43	2.28	11	2
1:A:585:GLN:NE2	1:A:585:GLN:N	0.43	2.66	20	3
1:A:589:ARG:CD	1:A:590:THR:N	0.43	2.82	24	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:584:LEU:O	1:A:588:LYS:CB	0.43	2.67	12	6
1:A:571:TRP:N	1:A:571:TRP:HE3	0.42	2.12	27	1
1:A:569:CYS:SG	1:A:571:TRP:NE1	0.42	2.92	31	1
1:A:571:TRP:CD1	1:A:591:HIS:NE2	0.42	2.87	29	1
1:A:585:GLN:O	1:A:588:LYS:HG2	0.42	2.14	25	3
1:A:586:ARG:O	1:A:589:ARG:CB	0.42	2.68	14	2
1:A:569:CYS:SG	1:A:571:TRP:CE3	0.42	3.13	7	2
1:A:571:TRP:CH2	1:A:588:LYS:HB2	0.42	2.50	30	1
1:A:578:PHE:CE1	1:A:583:GLU:CG	0.42	3.03	31	1
1:A:577:ARG:O	1:A:578:PHE:CD1	0.41	2.73	9	2
1:A:587:HIS:O	1:A:590:THR:HG22	0.41	2.15	12	1
1:A:567:PHE:CE2	1:A:579:THR:O	0.41	2.74	28	1
1:A:570:THR:O	1:A:571:TRP:O	0.41	2.38	28	1
1:A:585:GLN:O	1:A:588:LYS:HG3	0.41	2.15	17	1
1:A:578:PHE:CD1	1:A:583:GLU:HG2	0.41	2.50	31	2
1:A:569:CYS:HB3	1:A:584:LEU:HD23	0.41	1.93	29	1
1:A:569:CYS:N	1:A:584:LEU:HD21	0.41	2.31	12	1
1:A:578:PHE:CD2	1:A:583:GLU:CG	0.41	3.04	27	1
1:A:578:PHE:CG	1:A:583:GLU:HG2	0.41	2.50	27	1
1:A:578:PHE:CE1	1:A:583:GLU:HG3	0.41	2.51	31	1
1:A:573:TYR:O	1:A:573:TYR:CD2	0.41	2.74	21	1
1:A:571:TRP:CZ2	1:A:591:HIS:HB3	0.40	2.52	21	1
1:A:567:PHE:HB3	1:A:584:LEU:HD13	0.40	1.92	12	1
1:A:571:TRP:CZ3	1:A:588:LYS:HE2	0.40	2.52	5	1
1:A:567:PHE:CE1	1:A:581:SER:OG	0.40	2.73	16	1
1:A:580:ARG:N	1:A:580:ARG:HD2	0.40	2.31	4	1
1:A:571:TRP:CG	1:A:591:HIS:CD2	0.40	3.10	24	1
1:A:588:LYS:O	1:A:591:HIS:N	0.40	2.53	26	1
1:A:569:CYS:HB2	1:A:571:TRP:CE2	0.40	2.51	29	1
1:A:578:PHE:CZ	1:A:587:HIS:CD2	0.40	3.09	18	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	26/31 (84%)	23±1 (89±5%)	2±1 (8±5%)	1±1 (3±3%)	7	40
All	All	806/961 (84%)	716 (89%)	66 (8%)	24 (3%)	7	40

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	571	TRP	16
1	A	573	TYR	5
1	A	575	GLY	1
1	A	569	CYS	1
1	A	591	HIS	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	25/29 (86%)	17±2 (67±9%)	8±2 (33±9%)	1	12
All	All	775/899 (86%)	520 (67%)	255 (33%)	1	12

All 22 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	587	HIS	31
1	A	589	ARG	24
1	A	591	HIS	22
1	A	585	GLN	17
1	A	571	TRP	17
1	A	569	CYS	16
1	A	588	LYS	15
1	A	582	ASP	14
1	A	586	ARG	14
1	A	581	SER	12
1	A	590	THR	12
1	A	568	MET	11
1	A	573	TYR	11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	584	LEU	10
1	A	576	LYS	8
1	A	572	SER	8
1	A	577	ARG	4
1	A	580	ARG	3
1	A	567	PHE	2
1	A	570	THR	2
1	A	579	THR	1
1	A	574	CYS	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided