



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 23, 2021 – 02:12 PM EST

PDB ID : 1VTO
Title : 1.9 Å RESOLUTION REFINED STRUCTURE OF TBP RECOGNIZING
THE MINOR GROOVE OF TATAAAAG
Authors : Kim, J.L.; Burley, S.K.
Deposited on : 1996-09-06
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.16


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	C	14	57% (green), 14% (yellow), 29% (orange)
1	E	14	29% (green), 36% (yellow), 36% (orange)
2	D	14	14% (green), 64% (yellow), 14% (orange), 7% (red)
2	F	14	14% (green), 36% (yellow), 43% (orange), 7% (red)
3	A	190	78% (green), 19% (yellow), .. (grey)
3	B	190	84% (green), 14% (yellow), .. (grey)

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6447 atoms, of which 1830 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*GP*CP*TP*AP*TP*AP*AP*AP*AP*GP*GP*GP*CP*A)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P			
1	C	14	Total 318	C 137	H 31	N 60	O 77	P 13	0	0	0
1	E	14	Total 321	C 138	H 32	N 60	O 78	P 13	0	0	0

- Molecule 2 is a DNA chain called DNA (5'-D(*TP*GP*CP*CP*CP*TP*TP*TP*TP*AP*TP*AP*GP*C)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P			
2	D	14	Total 305	C 136	H 26	N 44	O 86	P 13	0	0	0
2	F	14	Total 305	C 136	H 26	N 44	O 86	P 13	0	0	0

- Molecule 3 is a protein called TATA BINDING PROTEIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
3	A	187	Total 1806	C 960	H 333	N 250	O 255	S 8	0	0	0
3	B	188	Total 1831	C 970	H 342	N 255	O 256	S 8	0	1	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	H	O		
4	C	38	Total 114	H 76	O 38	0	0
4	D	39	Total 117	H 78	O 39	0	0
4	E	37	Total 111	H 74	O 37	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	F	50	Total 150	H 100	O 50	0	0
4	A	172	Total 516	H 344	O 172	0	0
4	B	185	Total 553	H 368	O 185	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: DNA (5'-D(*GP*CP*TP*AP*TP*AP*AP*AP*AP*GP*GP*GP*CP*A)-3')

Chain C: 



- Molecule 1: DNA (5'-D(*GP*CP*TP*AP*TP*AP*AP*AP*AP*GP*GP*GP*CP*A)-3')

Chain E: 



- Molecule 2: DNA (5'-D(*TP*GP*CP*CP*CP*TP*TP*TP*TP*AP*TP*AP*GP*C)-3')

Chain D: 




- Molecule 2: DNA (5'-D(*TP*GP*CP*CP*CP*TP*TP*TP*TP*AP*TP*AP*GP*C)-3')

Chain F: 





- Molecule 3: TATA BINDING PROTEIN

Chain A: 



- Molecule 3: TATA BINDING PROTEIN

Chain B:  84% 14% ..



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	42.00Å 147.00Å 57.00Å 90.00° 90.50° 90.00°	Depositor
Resolution (Å)	6.00 – 1.90	Depositor
% Data completeness (in resolution range)	94.4 (6.00-1.90)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	5.20	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.194 , 0.264	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	6447	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	1.37	0/324	1.68	7/499 (1.4%)
1	E	1.43	0/326	1.82	13/502 (2.6%)
2	D	1.46	2/310 (0.6%)	2.02	15/476 (3.2%)
2	F	1.53	5/310 (1.6%)	2.42	22/476 (4.6%)
3	A	0.65	1/1503 (0.1%)	0.77	0/2024
3	B	0.68	1/1525 (0.1%)	0.79	0/2052
All	All	0.96	9/4298 (0.2%)	1.31	57/6029 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
2	D	0	1
2	F	0	1
All	All	0	3

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1122	CYS	CB-SG	-8.31	1.68	1.82
2	D	215	DT	C5-C7	6.84	1.54	1.50
2	F	322	DT	C5-C7	6.69	1.54	1.50
2	F	315	DT	C5-C7	5.93	1.53	1.50
2	D	221	DT	C5-C7	5.75	1.53	1.50

The worst 5 of 57 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	317	DC	O4'-C4'-C3'	-13.58	97.85	106.00
2	D	227	DG	O4'-C1'-N9	13.48	117.44	108.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	F	320	DT	C6-C5-C7	-10.79	116.43	122.90
2	F	317	DC	O4'-C1'-N1	10.71	115.49	108.00
2	F	316	DG	O4'-C1'-N9	8.91	114.24	108.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	207	DA	Sidechain
2	D	227	DG	Sidechain
2	F	318	DC	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	287	31	154	4	0
1	E	289	32	158	5	5
2	D	279	26	162	8	0
2	F	279	26	162	13	0
3	A	1473	333	1549	31	5
3	B	1489	342	1575	26	0
4	A	172	344	0	8	5
4	B	185	368	0	8	4
4	C	38	76	0	1	0
4	D	39	78	0	1	1
4	E	37	74	0	2	1
4	F	50	100	0	0	1
All	All	4617	1830	3760	81	11

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 81 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:156:LYS:HA	3:A:156:LYS:HE2	1.43	0.97
3:B:1041:LYS:HD3	3:B:1052:TYR:HE2	1.33	0.89
2:F:317:DC:H2'	2:F:318:DC:C6	2.09	0.87
4:C:924:HOH:O	3:A:58:ALA:HB3	1.75	0.87
3:A:198:ILE:HD11	3:B:1190:PRO:HG2	1.56	0.86

The worst 5 of 11 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:693:HOH:H1	4:B:613:HOH:H1[2_647]	1.18	0.42
4:A:693:HOH:O	4:B:613:HOH:H1[2_647]	1.19	0.41
1:E:311:DG:N7	3:A:65:ARG:HH22[2_657]	1.21	0.39
3:A:46:GLN:HE21	4:B:885:HOH:O[2_647]	1.31	0.29
4:A:693:HOH:O	4:B:613:HOH:O[2_647]	2.03	0.17

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	185/190 (97%)	177 (96%)	8 (4%)	0	100	100
3	B	187/190 (98%)	179 (96%)	8 (4%)	0	100	100
All	All	372/380 (98%)	356 (96%)	16 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	159/163 (98%)	154 (97%)	5 (3%)	40	32
3	B	162/163 (99%)	159 (98%)	3 (2%)	57	53
All	All	321/326 (98%)	313 (98%)	8 (2%)	47	41

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	147	LEU
3	B	1104	LEU
3	B	1038	LEU
3	A	133	LEU
3	A	179	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	49	ASN
3	B	1053	ASN
3	B	1137	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.