



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2023 – 11:45 PM EDT

PDB ID : 6XOH  
Title : Structure of SUMO1-ML00789344 adduct bound to SAE  
Authors : Sintchak, M.; Lane, W.; Bump, N.  
Deposited on : 2020-07-07  
Resolution : 2.23 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5912 (2.24-2.20)
Clashscore	141614	6646 (2.24-2.20)
Ramachandran outliers	138981	6543 (2.24-2.20)
Sidechain outliers	138945	6544 (2.24-2.20)
RSRZ outliers	127900	5797 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	346	
2	B	640	
3	C	101	

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6714 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SUMO-activating enzyme subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	311	2395	1521	408	454	12	0	0	0

- Molecule 2 is a protein called SUMO-activating enzyme subunit 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	490	3681	2355	624	683	19	0	0	0

- Molecule 3 is a protein called Small ubiquitin-related modifier 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	52	408	263	69	76	0	1	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total	O	S
			5	4	1
				0	0

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total	Zn	
			1	1	
				0	0

- Molecule 6 is {(1R,2S,4R)-4-[(5-{4-[(1R)-3,4-dihydro-1H-2-benzopyran-1-yl]thiophene-2-carbonyl}pyrimidin-4-yl)amino]-2-hydroxycyclopentyl)methyl sulfamate (three-letter code: VB7) (formula: C<sub>24</sub>H<sub>26</sub>N<sub>4</sub>O<sub>6</sub>S<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf		
6	C	1	Total	C	N	O	S
			36	24	4	6	2
				0	0		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	127	Total	O	
			127	127	
7	B	56	Total	O	
			56	56	
7	C	5	Total	O	
			5	5	
				0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SUMO-activating enzyme subunit 1






- Molecule 2: SUMO-activating enzyme subunit 2



ASP  
VAL  
ILE  
ALA  
LEU  
ASP

● Molecule 3: Small ubiquitin-related modifier 1



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.13Å 72.95Å 126.59Å 90.00° 92.93° 90.00°	Depositor
Resolution (Å)	43.31 – 2.23 43.31 – 2.23	Depositor EDS
% Data completeness (in resolution range)	96.6 (43.31-2.23) 96.7 (43.31-2.23)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.67 (at 2.22Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.194 , 0.244 0.194 , 0.245	Depositor DCC
$R_{free}$ test set	2554 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.4	Xtrriage
Anisotropy	0.051	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 44.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.025 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6714	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	66.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ZN, VB7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/2438	0.81	2/3296 (0.1%)
2	B	0.35	0/3745	0.71	1/5094 (0.0%)
3	C	0.38	0/409	0.67	0/542
All	All	0.38	0/6592	0.75	3/8932 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	35	ARG	CB-CG-CD	-5.87	96.33	111.60
1	A	42	VAL	CA-CB-CG2	5.70	119.44	110.90
2	B	390	THR	CA-CB-OG1	-5.05	98.40	109.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2395	0	2356	27	0
2	B	3681	0	3603	54	0
3	C	408	0	391	10	0
4	A	5	0	0	0	0
5	B	1	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	C	36	0	0	1	0
7	A	127	0	0	5	0
7	B	56	0	0	2	0
7	C	5	0	0	0	0
All	All	6714	0	6350	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:152:LYS:HE2	7:A:531:HOH:O	1.73	0.88
2:B:482:ILE:HG12	2:B:511:ILE:HD11	1.60	0.84
2:B:456:VAL:HG13	2:B:511:ILE:HG22	1.58	0.83
2:B:462:LEU:HD13	2:B:502:ASN:ND2	2.04	0.72
2:B:456:VAL:CG1	2:B:511:ILE:HG22	2.22	0.69
1:A:173:GLU:OE1	1:A:208:LYS:HE2	1.94	0.68
2:B:465:GLN:HG3	2:B:491:ILE:HG21	1.76	0.68
2:B:513:ASN:C	2:B:513:ASN:HD22	1.97	0.68
2:B:114:ASN:ND2	2:B:138:GLU:OE1	2.27	0.67
2:B:492:SER:HB3	2:B:498:THR:HG22	1.75	0.66
1:A:105:MET:HE3	1:A:105:MET:HA	1.78	0.66
2:B:245:LYS:HA	2:B:249:TYR:CE1	2.37	0.59
3:C:62:LEU:HD22	3:C:90:VAL:CG1	2.34	0.58
2:B:462:LEU:HD13	2:B:502:ASN:HD22	1.69	0.56
2:B:456:VAL:HG11	2:B:511:ILE:O	2.05	0.56
2:B:451:THR:HG22	2:B:547:VAL:HG21	1.88	0.55
2:B:455:ASN:HD21	2:B:458:LYS:HE3	1.71	0.55
1:A:10:GLY:HA3	1:A:32:LYS:HE2	1.89	0.55
2:B:521:ASP:OD1	2:B:523:LEU:HB3	2.07	0.53
2:B:476:VAL:O	2:B:478:PRO:HD3	2.08	0.53
2:B:178:THR:HG22	2:B:178:THR:O	2.09	0.53
3:C:62:LEU:HD22	3:C:90:VAL:HG12	1.91	0.52
2:B:170:PHE:HB2	2:B:175:ILE:HD11	1.90	0.52
2:B:476:VAL:C	2:B:478:PRO:HD3	2.30	0.52
2:B:516:ARG:NH2	7:B:803:HOH:O	2.43	0.51
3:C:36:PHE:HZ	3:C:50:SER:HB3	1.76	0.51
2:B:263:ILE:HG12	2:B:365:ILE:HG21	1.92	0.51
3:C:26:VAL:HG13	3:C:90:VAL:HG23	1.93	0.51
1:A:42:VAL:HG13	1:A:131:LEU:HD12	1.94	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:263:ILE:HG22	2:B:281:PRO:HG3	1.92	0.50
2:B:462:LEU:HB2	2:B:502:ASN:HB3	1.94	0.49
2:B:403:LEU:O	2:B:407:SER:HB3	2.12	0.49
2:B:456:VAL:HG23	2:B:534:HIS:ND1	2.27	0.49
1:A:325:HIS:HE1	7:A:616:HOH:O	1.95	0.49
2:B:257:LYS:HA	2:B:261:ASP:HB2	1.95	0.49
2:B:478:PRO:HA	2:B:520:ASP:O	2.12	0.49
2:B:456:VAL:HG12	2:B:507:SER:CA	2.43	0.48
1:A:164:TYR:HA	1:A:329:PHE:O	2.12	0.48
2:B:141:THR:HG22	2:B:394:ILE:HG12	1.96	0.48
2:B:284:TRP:O	2:B:288:GLN:HG2	2.12	0.48
2:B:333:LEU:O	2:B:333:LEU:HG	2.14	0.48
1:A:175:VAL:CG2	1:A:206:MET:CB	2.92	0.48
1:A:277:LEU:HD12	1:A:284:PRO:HA	1.96	0.47
2:B:468:ILE:HG12	2:B:544:PHE:CE2	2.50	0.47
1:A:159:PHE:CG	1:A:246:LEU:HD13	2.51	0.46
2:B:455:ASN:ND2	2:B:536:GLU:HG2	2.31	0.46
6:C:201:VB7:O10	6:C:201:VB7:N2	2.45	0.46
1:A:296:PHE:O	1:A:297:SER:HB2	2.15	0.46
2:B:513:ASN:C	2:B:513:ASN:ND2	2.68	0.46
2:B:123:ASN:ND2	2:B:160:GLU:HG2	2.31	0.46
2:B:432:CYS:SG	3:C:65:LEU:HD21	2.56	0.46
3:C:30:ASP:O	3:C:31:SER:HB2	2.15	0.45
1:A:10:GLY:CA	1:A:32:LYS:HE2	2.46	0.45
2:B:198:GLY:HA2	2:B:342:LEU:O	2.17	0.45
1:A:92:ALA:HB1	1:A:110:VAL:HG22	1.99	0.45
2:B:138:GLU:OE2	7:B:801:HOH:O	2.21	0.45
2:B:456:VAL:HG12	2:B:507:SER:HA	1.98	0.44
1:A:256:ASP:HB3	1:A:257:PRO:HD2	1.99	0.44
2:B:44:ILE:O	2:B:89:ILE:HA	2.17	0.44
2:B:457:HIS:CD2	2:B:507:SER:CB	3.01	0.44
1:A:18:GLN:NE2	7:A:508:HOH:O	2.50	0.44
1:A:105:MET:HA	1:A:105:MET:CE	2.47	0.44
2:B:49:LEU:HD13	2:B:50:ASP:OD1	2.17	0.44
1:A:282:ILE:HG21	1:A:286:LEU:HD22	2.00	0.44
2:B:513:ASN:HB2	2:B:534:HIS:HB3	2.00	0.43
2:B:425:ARG:O	2:B:426:LYS:HB2	2.19	0.43
2:B:34:LYS:O	2:B:38:LEU:HG	2.19	0.43
3:C:67:GLU:HG3	3:C:67:GLU:O	2.19	0.43
1:A:175:VAL:HG23	1:A:206:MET:CB	2.49	0.42
3:C:36:PHE:CZ	3:C:50:SER:HB3	2.54	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:46:LYS:HG2	3:C:47:LEU:N	2.34	0.42
1:A:12:SER:OG	1:A:15:GLU:HG3	2.19	0.42
1:A:159:PHE:HD2	7:A:619:HOH:O	2.03	0.42
1:A:177:GLU:OE1	1:A:177:GLU:N	2.53	0.41
2:B:457:HIS:CD2	2:B:507:SER:HB2	2.55	0.41
2:B:475:MET:HG2	2:B:521:ASP:HB2	2.02	0.41
1:A:147:HIS:HB2	7:A:566:HOH:O	2.20	0.41
1:A:53:LYS:HB3	1:A:53:LYS:HE3	1.65	0.41
2:B:282:LEU:HD23	2:B:282:LEU:HA	1.95	0.41
1:A:146:CYS:HB3	1:A:151:ILE:O	2.20	0.41
2:B:516:ARG:HG2	2:B:531:ASN:OD1	2.21	0.41
2:B:65:LYS:O	2:B:68:VAL:HG23	2.20	0.41
2:B:23:VAL:HG12	2:B:116:LEU:HD21	2.03	0.41
2:B:98:ASN:OD1	2:B:98:ASN:C	2.59	0.40
1:A:22:GLN:HB3	2:B:387:ILE:HG22	2.02	0.40
1:A:131:LEU:HG	1:A:134:CYS:SG	2.61	0.40
1:A:323:PRO:HG2	2:B:421:GLN:NE2	2.36	0.40
2:B:157:GLU:HG3	2:B:435:ASP:HB2	2.03	0.40
3:C:64:PHE:CE1	3:C:90:VAL:HG22	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/346 (89%)	298 (97%)	8 (3%)	1 (0%)	41	45
2	B	472/640 (74%)	437 (93%)	34 (7%)	1 (0%)	47	54
3	C	44/101 (44%)	39 (89%)	5 (11%)	0	100	100
All	All	823/1087 (76%)	774 (94%)	47 (6%)	2 (0%)	47	54

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	226	SER
2	B	513	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	256/296 (86%)	244 (95%)	12 (5%)	26	31
2	B	387/552 (70%)	369 (95%)	18 (5%)	26	31
3	C	41/91 (45%)	34 (83%)	7 (17%)	2	1
All	All	684/939 (73%)	647 (95%)	37 (5%)	22	25

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	LEU
1	A	84	THR
1	A	89	ARG
1	A	110	VAL
1	A	113	GLU
1	A	131	LEU
1	A	159	PHE
1	A	175	VAL
1	A	177	GLU
1	A	205	THR
1	A	228	LYS
1	A	280	LEU
2	B	34	LYS
2	B	49	LEU
2	B	97	MET
2	B	138	GLU
2	B	158	CYS
2	B	160	GLU
2	B	162	HIS
2	B	170	PHE
2	B	277	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	394	ILE
2	B	465	GLN
2	B	476	VAL
2	B	497	GLU
2	B	513	ASN
2	B	515	SER
2	B	517	LEU
2	B	520	ASP
2	B	538	LEU
3	C	22	ILE
3	C	25	LYS
3	C	46	LYS
3	C	47	LEU
3	C	61	SER
3	C	66	PHE
3	C	86	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
2	B	74	GLN
2	B	108	GLN
2	B	123	ASN
2	B	311	GLN
2	B	457	HIS
2	B	481	GLN
2	B	513	ASN
3	C	94	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	VB7	C	201	3	37,40,40	2.72	13 (35%)	45,58,58	3.09	17 (37%)
4	SO4	A	401	-	4,4,4	0.43	0	6,6,6	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	VB7	C	201	3	-	0/14/44/44	0/5/5/5

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	201	VB7	O35-S33	8.48	1.49	1.42
6	C	201	VB7	O36-S33	7.70	1.48	1.42
6	C	201	VB7	O31-S33	-5.46	1.50	1.57
6	C	201	VB7	S33-N34	4.78	1.63	1.58
6	C	201	VB7	C13-C16	3.45	1.56	1.52
6	C	201	VB7	C11-S15	-3.36	1.68	1.72
6	C	201	VB7	C14-C13	-3.23	1.35	1.37
6	C	201	VB7	C29-C1	3.07	1.59	1.53
6	C	201	VB7	O21-C20	2.73	1.50	1.44
6	C	201	VB7	C11-C9	2.70	1.56	1.50
6	C	201	VB7	C19-C18	2.69	1.55	1.51
6	C	201	VB7	O32-C28	2.05	1.47	1.43

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	201	VB7	C17-C16	2.01	1.55	1.51

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	201	VB7	O36-S33-O35	-8.88	111.56	119.97
6	C	201	VB7	C7-C8-C3	7.39	121.78	114.55
6	C	201	VB7	C13-C14-S15	-6.40	107.81	112.25
6	C	201	VB7	O36-S33-N34	5.53	117.48	109.14
6	C	201	VB7	C30-O31-S33	5.11	123.61	117.21
6	C	201	VB7	C20-O21-C16	4.93	117.84	111.88
6	C	201	VB7	N6-C5-N4	-4.92	122.29	127.70
6	C	201	VB7	N2-C3-N4	4.26	124.18	117.87
6	C	201	VB7	C8-C3-N4	-4.20	116.17	121.90
6	C	201	VB7	C13-C16-C17	-3.82	108.47	114.32
6	C	201	VB7	C7-N6-C5	3.46	120.22	115.80
6	C	201	VB7	O21-C16-C13	3.45	117.41	108.62
6	C	201	VB7	O35-S33-N34	-3.25	104.22	109.14
6	C	201	VB7	C8-C3-N2	-2.75	119.61	121.72
6	C	201	VB7	C8-C7-N6	-2.70	117.78	123.81
6	C	201	VB7	C19-C18-C17	2.50	124.87	121.13
6	C	201	VB7	C3-C8-C9	-2.26	117.99	124.67

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	201	VB7	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	311/346 (89%)	-0.23	7 (2%) 60 58	29, 47, 77, 108	0
2	B	490/640 (76%)	0.53	61 (12%) 4 3	29, 69, 124, 158	0
3	C	52/101 (51%)	1.61	18 (34%) 0 0	39, 102, 141, 149	0
All	All	853/1087 (78%)	0.32	86 (10%) 7 5	29, 59, 123, 158	0

All (86) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	500	ALA	7.4
2	B	506	LEU	6.0
2	B	458	LYS	6.0
3	C	22	ILE	5.7
3	C	34	ILE	5.6
3	C	36	PHE	5.5
3	C	24	LEU	5.5
3	C	47	LEU	5.5
2	B	482	ILE	4.7
2	B	454	LEU	4.4
2	B	333	LEU	4.4
2	B	459	VAL	4.3
2	B	538	LEU	4.2
3	C	51	TYR	4.1
2	B	206	VAL	3.9
2	B	479	ASP	3.9
2	B	334	ALA	3.9
2	B	488	THR	3.6
3	C	32	SER	3.6
2	B	481	GLN	3.6
2	B	243	TRP	3.6
3	C	33	GLU	3.5
2	B	489	ILE	3.5

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	B	529	LEU	3.4
2	B	532	ILE	3.4
3	C	38	VAL	3.4
1	A	205	THR	3.4
3	C	52	CYS	3.4
2	B	329	LEU	3.3
2	B	517	LEU	3.3
2	B	393	VAL	3.2
2	B	490	LEU	3.1
2	B	452	VAL	3.1
3	C	44	LEU	3.1
1	A	302	VAL	3.0
1	A	306	VAL	3.0
2	B	249	TYR	3.0
2	B	331	VAL	2.9
2	B	397	LEU	2.8
2	B	537	ASP	2.8
2	B	533	LEU	2.8
2	B	539	GLY	2.8
2	B	534	HIS	2.8
2	B	464	LEU	2.7
2	B	395	ALA	2.7
2	B	394	ILE	2.7
1	A	305	VAL	2.7
3	C	86	ASP	2.7
2	B	366	PHE	2.7
2	B	263	ILE	2.7
2	B	530	ILE	2.7
2	B	192	LEU	2.7
2	B	195	GLN	2.6
2	B	480	VAL	2.6
2	B	461	VAL	2.6
1	A	304	ALA	2.6
2	B	197	PHE	2.6
2	B	392	ALA	2.5
3	C	37	LYS	2.5
1	A	307	GLY	2.5
2	B	508	GLU	2.5
2	B	140	GLY	2.5
3	C	23	LYS	2.4
2	B	398	ILE	2.4
2	B	247	THR	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
3	C	66	PHE	2.4
2	B	396	GLY	2.3
2	B	429	LEU	2.3
2	B	400	LEU	2.2
3	C	53	GLN	2.2
2	B	196	LEU	2.2
1	A	303	CYS	2.2
2	B	269	MET	2.2
2	B	336	LYS	2.1
2	B	399	VAL	2.1
2	B	512	ARG	2.1
2	B	350	SER	2.1
3	C	50	SER	2.1
2	B	484	ASP	2.1
2	B	211	ALA	2.1
2	B	493	SER	2.1
2	B	273	TRP	2.1
2	B	335	GLU	2.0
2	B	330	ARG	2.0
3	C	46	LYS	2.0
2	B	214	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	VB7	C	201	36/36	0.97	0.12	36,46,74,76	0
5	ZN	B	700	1/1	0.98	0.10	61,61,61,61	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	SO4	A	401	5/5	0.98	0.16	59,59,70,78	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.