



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 30, 2023 – 07:54 PM JST

PDB ID : 4YSX
Title : Crystal structure of Mitochondrial rhodoquinol-fumarate reductase from *Ascaris suum* with the specific inhibitor NN23
Authors : Harada, S.; Shiba, T.; Sato, D.; Yamamoto, A.; Nagahama, M.; Yone, A.; Inaoka, D.K.; Sakamoto, K.; Inoue, M.; Honma, T.; Kita, K.
Deposited on : 2015-03-17
Resolution : 2.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	645	 2% 79% 15% . .
1	E	645	 2% 81% 13% . .
2	B	282	 2% 77% 10% . 11%
2	F	282	 2% 79% 9% . 11%
3	C	188	 5% 73% 7% . 19%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	188	 <p>7% 75% 7% 18%</p>
4	D	156	 <p>4% 71% 10% 17%</p>
4	H	156	 <p>6% 69% 12% 17%</p>

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 19103 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Succinate dehydrogenase flavoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	616	4787	3004	855	900	28	0	0	0
1	E	616	4787	3004	855	900	28	0	0	0

- Molecule 2 is a protein called Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	250	1985	1263	338	361	23	0	0	0
2	F	252	1994	1268	340	363	23	0	0	0

- Molecule 3 is a protein called Cytochrome b-large subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	153	1217	813	204	194	6	0	0	0
3	G	155	1236	823	209	198	6	0	0	0

- Molecule 4 is a protein called Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	129	998	659	165	169	5	0	0	0
4	H	129	998	659	165	169	5	0	0	0

- Molecule 5 is MALONATE ION (three-letter code: MLI) (formula: C₃H₂O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	3	4		
5	E	1	Total	C	O	0	0
			7	3	4		

- Molecule 6 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
6	E	1	53	27	9	15	2	0	0

- Molecule 7 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Fe	S		
7	B	1	4	2	2	0	0
7	F	1	4	2	2	0	0

- Molecule 8 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	1	Total	Fe S	0	0
			8	4 4		
8	F	1	Total	Fe S	0	0
			8	4 4		

- Molecule 9 is FE3-S4 CLUSTER (three-letter code: F3S) (formula: Fe₃S₄).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	B	1	Total	Fe S	0	0
			7	3 4		
9	F	1	Total	Fe S	0	0
			7	3 4		

- Molecule 10 is N-(4-tert-butylbenzyl)-2-(trifluoromethyl)benzamide (three-letter code: E23) (formula: $C_{19}H_{20}F_3NO$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
10	B	1	Total	C	F	N	O	0	0
			24	19	3	1	1		
10	G	1	Total	C	F	N	O	0	0
			24	19	3	1	1		

- Molecule 11 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
11	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
11	G	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 12 is L-ALPHA-PHOSPHATIDYL-BETA-OLEOYL-GAMMA-PALMITOYL-PHOSPHATIDYLETHANOLAMINE (three-letter code: EPH) (formula: C₃₉H₆₈NO₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	D	1	Total	C	N	O	P	0	0
			44	34	1	8	1		
12	H	1	Total	C	N	O	P	0	0
			44	34	1	8	1		

- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	186	Total	O	0	0
			186	186		
13	B	104	Total	O	0	0
			104	104		
13	C	33	Total	O	0	0
			33	33		
13	D	27	Total	O	0	0
			27	27		
13	E	207	Total	O	0	0
			207	207		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	F	111	Total 111	O 111	0	0
13	G	36	Total 36	O 36	0	0
13	H	17	Total 17	O 17	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Succinate dehydrogenase flavoprotein




- Molecule 1: Succinate dehydrogenase flavoprotein




- Molecule 2: Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial

Chain B: 2% 77% 10% 11%




- Molecule 2: Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial

Chain F: 2% 79% 9% 11%




- Molecule 3: Cytochrome b-large subunit

Chain C: 5% 73% 7% 19%




- Molecule 3: Cytochrome b-large subunit

Chain G: 7% 75% 7% 18%




- Molecule 4: Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial

Chain D: 4% 71% 10% 17%



- Molecule 4: Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	124.20Å 127.91Å 220.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.25 20.00 – 2.25	Depositor EDS
% Data completeness (in resolution range)	96.3 (20.00-2.25) 96.4 (20.00-2.25)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.15 (at 2.26Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.181 , 0.221 0.185 , 0.223	Depositor DCC
R_{free} test set	8058 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	42.8	Xtrriage
Anisotropy	0.127	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 35.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.017 for k,h,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	19103	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FES, F3S, E23, SF4, EPH, MLI, HEM, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/4889	0.67	2/6605 (0.0%)
1	E	0.45	0/4889	0.70	7/6605 (0.1%)
2	B	0.46	0/2029	0.67	0/2739
2	F	0.46	1/2038 (0.0%)	0.63	0/2751
3	C	0.40	0/1255	0.59	0/1709
3	G	0.39	0/1275	0.60	0/1735
4	D	0.40	0/1030	0.58	0/1406
4	H	0.40	0/1030	0.60	0/1406
All	All	0.44	1/18435 (0.0%)	0.66	9/24956 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	94	CYS	C-N	-5.26	1.23	1.33

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	E	604	ARG	NE-CZ-NH2	-6.32	117.14	120.30
1	E	320	ARG	NE-CZ-NH2	6.09	123.34	120.30
1	E	241	ARG	NE-CZ-NH2	5.82	123.21	120.30
1	A	168	ARG	NE-CZ-NH2	-5.74	117.43	120.30
1	E	359	ARG	NE-CZ-NH1	5.74	123.17	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4787	0	4722	61	0
1	E	4787	0	4720	54	0
2	B	1985	0	2001	16	0
2	F	1994	0	2006	12	0
3	C	1217	0	1265	7	0
3	G	1236	0	1278	8	0
4	D	998	0	985	11	0
4	H	998	0	985	11	0
5	A	7	0	2	0	0
5	E	7	0	2	1	0
6	A	53	0	31	7	0
6	E	53	0	31	7	0
7	B	4	0	0	0	0
7	F	4	0	0	0	0
8	B	8	0	0	0	0
8	F	8	0	0	0	0
9	B	7	0	0	0	0
9	F	7	0	0	0	0
10	B	24	0	20	0	0
10	G	24	0	20	0	0
11	C	43	0	30	5	0
11	G	43	0	30	5	0
12	D	44	0	53	2	0
12	H	44	0	53	1	0
13	A	186	0	0	4	0
13	B	104	0	0	2	0
13	C	33	0	0	0	0
13	D	27	0	0	1	0
13	E	207	0	0	3	0
13	F	111	0	0	3	0
13	G	36	0	0	1	0
13	H	17	0	0	1	0
All	All	19103	0	18234	175	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 175 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:HIS:NE2	6:A:702:FAD:HM82	1.25	1.48
1:E:79:HIS:NE2	6:E:702:FAD:HM82	1.33	1.36
1:E:79:HIS:NE2	6:E:702:FAD:C8M	2.07	1.16
1:A:79:HIS:NE2	6:A:702:FAD:C8M	2.12	1.13
1:E:476:ASN:HD21	1:E:550:GLN:HE22	1.19	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	614/645 (95%)	594 (97%)	20 (3%)	0	100	100
1	E	614/645 (95%)	599 (98%)	15 (2%)	0	100	100
2	B	248/282 (88%)	239 (96%)	9 (4%)	0	100	100
2	F	250/282 (89%)	243 (97%)	7 (3%)	0	100	100
3	C	151/188 (80%)	147 (97%)	4 (3%)	0	100	100
3	G	153/188 (81%)	151 (99%)	2 (1%)	0	100	100
4	D	127/156 (81%)	125 (98%)	2 (2%)	0	100	100
4	H	127/156 (81%)	122 (96%)	5 (4%)	0	100	100
All	All	2284/2542 (90%)	2220 (97%)	64 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	502/527 (95%)	471 (94%)	31 (6%)	18	17
1	E	502/527 (95%)	470 (94%)	32 (6%)	17	16
2	B	220/242 (91%)	205 (93%)	15 (7%)	16	14
2	F	220/242 (91%)	208 (94%)	12 (6%)	21	21
3	C	127/158 (80%)	119 (94%)	8 (6%)	18	17
3	G	129/158 (82%)	123 (95%)	6 (5%)	26	29
4	D	98/119 (82%)	89 (91%)	9 (9%)	9	7
4	H	98/119 (82%)	87 (89%)	11 (11%)	6	4
All	All	1896/2092 (91%)	1772 (94%)	124 (6%)	17	16

5 of 124 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	79	ILE
3	G	160	LEU
1	E	205	LEU
3	G	136	ILE
4	H	100	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 50 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	252	ASN
1	E	497	GLN
4	H	140	ASN
1	E	355	GLN
1	E	451	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	FAD	E	702	-	53,58,58	1.19	5 (9%)	68,89,89	1.60	16 (23%)
11	HEM	C	201	4,3	41,50,50	1.37	6 (14%)	45,82,82	2.41	19 (42%)
12	EPH	D	201	-	43,43,48	1.13	2 (4%)	45,48,53	1.18	4 (8%)
8	SF4	F	302	2	0,12,12	-	-	-	-	-
7	FES	F	301	2	0,4,4	-	-	-	-	-
6	FAD	A	702	-	53,58,58	1.30	7 (13%)	68,89,89	1.53	15 (22%)
5	MLI	E	701	-	6,6,6	1.07	0	7,7,7	1.41	1 (14%)
10	E23	G	202	-	25,25,25	0.40	0	37,37,37	0.97	4 (10%)
7	FES	B	301	2	0,4,4	-	-	-	-	-
11	HEM	G	201	4,3	41,50,50	1.32	5 (12%)	45,82,82	2.57	20 (44%)
8	SF4	B	302	2	0,12,12	-	-	-	-	-
9	F3S	B	303	2	0,9,9	-	-	-	-	-
10	E23	B	304	-	25,25,25	0.48	0	37,37,37	1.09	3 (8%)
5	MLI	A	701	-	6,6,6	1.03	0	7,7,7	1.07	0
9	F3S	F	303	2	0,9,9	-	-	-	-	-
12	EPH	H	201	-	43,43,48	1.17	2 (4%)	45,48,53	1.18	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	FAD	E	702	-	-	6/30/50/50	0/6/6/6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	HEM	C	201	4,3	-	4/12/54/54	-
12	EPH	D	201	-	-	28/47/47/52	-
8	SF4	F	302	2	-	-	0/6/5/5
7	FES	F	301	2	-	-	0/1/1/1
6	FAD	A	702	-	-	6/30/50/50	0/6/6/6
5	MLI	E	701	-	-	2/4/4/4	-
10	E23	G	202	-	-	0/21/21/21	0/2/2/2
7	FES	B	301	2	-	-	0/1/1/1
11	HEM	G	201	4,3	-	4/12/54/54	-
8	SF4	B	302	2	-	-	0/6/5/5
9	F3S	B	303	2	-	-	0/3/3/3
10	E23	B	304	-	-	6/21/21/21	0/2/2/2
5	MLI	A	701	-	-	0/4/4/4	-
9	F3S	F	303	2	-	-	0/3/3/3
12	EPH	H	201	-	-	22/47/47/52	-

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	H	201	EPH	O2-C4	4.99	1.47	1.33
12	D	201	EPH	O2-C4	4.63	1.46	1.33
12	H	201	EPH	O1-C3	4.61	1.47	1.34
12	D	201	EPH	O1-C3	4.61	1.47	1.34
6	A	702	FAD	C9A-C5X	4.33	1.48	1.41

The worst 5 of 86 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	G	201	HEM	CAD-C3D-C4D	6.28	135.63	124.66
11	C	201	HEM	CAD-C3D-C4D	5.65	134.53	124.66
11	G	201	HEM	O2D-CGD-CBD	5.27	130.96	114.03
11	G	201	HEM	C1B-NB-C4B	5.27	110.51	105.07
11	G	201	HEM	CHD-C1D-ND	4.91	129.77	124.43

There are no chirality outliers.

5 of 78 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	702	FAD	N10-C1'-C2'-O2'
6	A	702	FAD	N10-C1'-C2'-C3'

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
6	E	702	FAD	N10-C1'-C2'-O2'
6	E	702	FAD	N10-C1'-C2'-C3'
11	C	201	HEM	C2D-C3D-CAD-CBD


There are no ring outliers.



7 monomers are involved in 27 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	E	702	FAD	7	0
11	C	201	HEM	5	0
12	D	201	EPH	2	0
6	A	702	FAD	7	0
5	E	701	MLI	1	0
11	G	201	HEM	5	0
12	H	201	EPH	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	616/645 (95%)	-0.26	11 (1%) 68 71	32, 46, 70, 100	0
1	E	616/645 (95%)	-0.32	11 (1%) 68 71	31, 43, 64, 123	0
2	B	250/282 (88%)	-0.29	6 (2%) 59 62	32, 41, 64, 87	0
2	F	252/282 (89%)	-0.29	6 (2%) 59 62	33, 41, 68, 98	0
3	C	153/188 (81%)	-0.09	9 (5%) 22 24	35, 50, 88, 127	0
3	G	155/188 (82%)	-0.04	14 (9%) 9 10	38, 51, 88, 129	0
4	D	129/156 (82%)	-0.18	7 (5%) 25 28	40, 51, 91, 116	0
4	H	129/156 (82%)	0.00	9 (6%) 16 17	42, 53, 110, 151	0
All	All	2300/2542 (90%)	-0.24	73 (3%) 47 50	31, 46, 76, 151	0

The worst 5 of 73 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	188	HIS	8.4
4	H	51	PRO	8.3
4	H	49	PHE	8.1
4	H	52	LEU	7.7
4	H	50	LYS	7.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
12	EPH	H	201	44/49	0.80	0.26	61,76,84,97	0
12	EPH	D	201	44/49	0.89	0.18	57,68,73,87	0
10	E23	G	202	24/24	0.95	0.09	36,42,60,61	0
11	HEM	G	201	43/43	0.97	0.12	43,45,54,59	0
10	E23	B	304	24/24	0.97	0.09	36,40,58,59	0
8	SF4	F	302	8/8	0.97	0.07	35,36,37,37	0
5	MLI	A	701	7/7	0.98	0.07	40,41,45,46	0
5	MLI	E	701	7/7	0.98	0.07	37,39,42,45	0
6	FAD	A	702	53/53	0.98	0.08	31,35,39,40	0
11	HEM	C	201	43/43	0.98	0.12	43,47,52,56	0
6	FAD	E	702	53/53	0.98	0.09	30,34,40,45	0
7	FES	F	301	4/4	0.98	0.07	32,32,33,34	0
8	SF4	B	302	8/8	0.98	0.07	34,35,36,36	0
7	FES	B	301	4/4	0.99	0.07	33,33,34,34	0
9	F3S	B	303	7/7	0.99	0.10	35,36,38,38	0
9	F3S	F	303	7/7	0.99	0.09	35,35,37,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around EPH H 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around EPH D 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around E23 G 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM G 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around E23 B 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD A 702:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM C 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.