



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 12, 2023 – 12:13 pm GMT

PDB ID : 3ZLP
Title : Crystal structure of Schistosoma mansoni Peroxiredoxin 1 C48P mutant form with four decamers in the asymmetric unit
Authors : Saccoccia, F.; Angelucci, F.; Ardini, M.; Boumis, G.; Brunori, M.; DiLeandro, L.; Ippoliti, R.; Miele, A.E.; Natoli, G.; Scotti, S.; Bellelli, A.
Deposited on : 2013-02-04
Resolution : 3.52 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1161 (3.60-3.44)
Ramachandran outliers	138981	1206 (3.60-3.44)
Sidechain outliers	138945	1207 (3.60-3.44)
RSRZ outliers	127900	1080 (3.60-3.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	186	 73% 14% 12%
1	B	186	 76% 13% 11%
1	C	186	 76% 12% 11%
1	D	186	 75% 13% 12%
1	E	186	 78% 7% 13%
1	F	186	 74% 14% 11%
1	G	186	 72% 16% 13%








Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	H	186	% 70% 17% • 11%
1	I	186	74% 13% 12%
1	J	186	76% 13% • 11%
1	K	186	70% 16% • 13%
1	L	186	% 69% 18% • 12%
1	M	186	75% 13% • 11%
1	N	186	72% 15% • 12%
1	O	186	% 76% 11% • 12%
1	P	186	73% 13% • 11%
1	Q	186	2% 66% 18% • 16%
1	R	186	77% 10% • 12%
1	S	186	73% 15% 12%
1	T	186	73% 16% 11%
1	U	186	74% 11% • 12%
1	V	186	72% 16% • 11%
1	W	186	74% 14% • 11%
1	X	186	74% 13% • 12%
1	Y	186	% 72% 15% • 12%
1	Z	186	74% 14% • 11%
1	a	186	% 69% 18% • 12%
1	b	186	75% 14% 11%
1	c	186	% 72% 13% •• 12%
1	d	186	% 74% 14% • 11%
1	e	186	70% 17% • 13%
1	f	186	69% 18% • 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	g	186	 73% 15% • 11%
1	h	186	%  72% 15% •• 12%
1	i	186	%  65% 19% • 15%
1	j	186	%  72% 15% • 11%
1	k	186	3%  72% 15% • 12%
1	l	186	 69% 19% • 11%
1	m	186	%  72% 13% 16%
1	n	186	%  74% 14% • 11%

2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 52692 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THIOREDOXIN PEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	163	1310	840	222	244	4	0	0	0
1	B	165	1337	860	226	246	5	0	2	0
1	C	165	1325	850	224	246	5	0	0	0
1	D	164	1331	854	227	246	4	0	2	0
1	E	162	1300	834	219	243	4	0	0	0
1	F	165	1325	850	224	246	5	0	0	0
1	G	162	1306	838	219	245	4	0	1	0
1	H	165	1325	850	224	246	5	0	0	0
1	I	163	1316	844	222	246	4	0	1	0
1	J	166	1332	854	225	248	5	0	0	0
1	K	161	1291	828	217	242	4	0	0	0
1	L	164	1327	854	223	245	5	0	2	0
1	M	165	1325	850	224	246	5	0	0	0
1	N	164	1331	854	227	246	4	0	2	0
1	O	163	1310	840	222	244	4	0	0	0
1	P	165	1325	850	224	246	5	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	Q	157	1267	810	214	239	4	0	1	0
1	R	164	1315	844	221	245	5	0	0	0
1	S	163	1316	844	222	246	4	0	1	0
1	T	166	1332	854	225	248	5	0	0	0
1	U	163	1310	840	222	244	4	0	0	0
1	V	165	1337	860	226	246	5	0	2	0
1	W	165	1325	850	224	246	5	0	0	0
1	X	164	1331	854	227	246	4	0	2	0
1	Y	163	1310	840	222	244	4	0	0	0
1	Z	165	1325	850	224	246	5	0	0	0
1	a	163	1316	844	222	246	4	0	1	0
1	b	165	1325	850	224	246	5	0	0	0
1	c	163	1316	844	222	246	4	0	1	0
1	d	166	1332	854	225	248	5	0	0	0
1	e	162	1300	834	219	243	4	0	0	0
1	f	165	1337	860	226	246	5	0	2	0
1	g	165	1325	850	224	246	5	0	0	0
1	h	163	1321	848	224	245	4	0	2	0
1	i	158	1271	812	217	238	4	0	0	0
1	j	165	1325	850	224	246	5	0	0	0
1	k	163	1316	844	222	246	4	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	l	165	1325	850	224	246	5	0	0	0
1	m	157	1267	810	214	239	4	0	1	0
1	n	166	1332	854	225	248	5	0	0	0

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	THR	-	expression tag	UNP O97161
A	48	PRO	CYS	engineered mutation	UNP O97161
B	0	THR	-	expression tag	UNP O97161
B	48	PRO	CYS	engineered mutation	UNP O97161
C	0	THR	-	expression tag	UNP O97161
C	48	PRO	CYS	engineered mutation	UNP O97161
D	0	THR	-	expression tag	UNP O97161
D	48	PRO	CYS	engineered mutation	UNP O97161
E	0	THR	-	expression tag	UNP O97161
E	48	PRO	CYS	engineered mutation	UNP O97161
F	0	THR	-	expression tag	UNP O97161
F	48	PRO	CYS	engineered mutation	UNP O97161
G	0	THR	-	expression tag	UNP O97161
G	48	PRO	CYS	engineered mutation	UNP O97161
H	0	THR	-	expression tag	UNP O97161
H	48	PRO	CYS	engineered mutation	UNP O97161
I	0	THR	-	expression tag	UNP O97161
I	48	PRO	CYS	engineered mutation	UNP O97161
J	0	THR	-	expression tag	UNP O97161
J	48	PRO	CYS	engineered mutation	UNP O97161
K	0	THR	-	expression tag	UNP O97161
K	48	PRO	CYS	engineered mutation	UNP O97161
L	0	THR	-	expression tag	UNP O97161
L	48	PRO	CYS	engineered mutation	UNP O97161
M	0	THR	-	expression tag	UNP O97161
M	48	PRO	CYS	engineered mutation	UNP O97161
N	0	THR	-	expression tag	UNP O97161
N	48	PRO	CYS	engineered mutation	UNP O97161
O	0	THR	-	expression tag	UNP O97161
O	48	PRO	CYS	engineered mutation	UNP O97161
P	0	THR	-	expression tag	UNP O97161
P	48	PRO	CYS	engineered mutation	UNP O97161

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
Q	0	THR	-	expression tag	UNP O97161
Q	48	PRO	CYS	engineered mutation	UNP O97161
R	0	THR	-	expression tag	UNP O97161
R	48	PRO	CYS	engineered mutation	UNP O97161
S	0	THR	-	expression tag	UNP O97161
S	48	PRO	CYS	engineered mutation	UNP O97161
T	0	THR	-	expression tag	UNP O97161
T	48	PRO	CYS	engineered mutation	UNP O97161
U	0	THR	-	expression tag	UNP O97161
U	48	PRO	CYS	engineered mutation	UNP O97161
V	0	THR	-	expression tag	UNP O97161
V	48	PRO	CYS	engineered mutation	UNP O97161
W	0	THR	-	expression tag	UNP O97161
W	48	PRO	CYS	engineered mutation	UNP O97161
X	0	THR	-	expression tag	UNP O97161
X	48	PRO	CYS	engineered mutation	UNP O97161
Y	0	THR	-	expression tag	UNP O97161
Y	48	PRO	CYS	engineered mutation	UNP O97161
Z	0	THR	-	expression tag	UNP O97161
Z	48	PRO	CYS	engineered mutation	UNP O97161
a	0	THR	-	expression tag	UNP O97161
a	48	PRO	CYS	engineered mutation	UNP O97161
b	0	THR	-	expression tag	UNP O97161
b	48	PRO	CYS	engineered mutation	UNP O97161
c	0	THR	-	expression tag	UNP O97161
c	48	PRO	CYS	engineered mutation	UNP O97161
d	0	THR	-	expression tag	UNP O97161
d	48	PRO	CYS	engineered mutation	UNP O97161
e	0	THR	-	expression tag	UNP O97161
e	48	PRO	CYS	engineered mutation	UNP O97161
f	0	THR	-	expression tag	UNP O97161
f	48	PRO	CYS	engineered mutation	UNP O97161
g	0	THR	-	expression tag	UNP O97161
g	48	PRO	CYS	engineered mutation	UNP O97161
h	0	THR	-	expression tag	UNP O97161
h	48	PRO	CYS	engineered mutation	UNP O97161
i	0	THR	-	expression tag	UNP O97161
i	48	PRO	CYS	engineered mutation	UNP O97161
j	0	THR	-	expression tag	UNP O97161
j	48	PRO	CYS	engineered mutation	UNP O97161
k	0	THR	-	expression tag	UNP O97161
k	48	PRO	CYS	engineered mutation	UNP O97161

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
l	0	THR	-	expression tag	UNP O97161
l	48	PRO	CYS	engineered mutation	UNP O97161
m	0	THR	-	expression tag	UNP O97161
m	48	PRO	CYS	engineered mutation	UNP O97161
n	0	THR	-	expression tag	UNP O97161
n	48	PRO	CYS	engineered mutation	UNP O97161


- Molecule 1: THIOREDOXIN PEROXIDASE

Chain E:  78% 7% 13%



- Molecule 1: THIOREDOXIN PEROXIDASE

Chain F:  74% 14% 11%




- Molecule 1: THIOREDOXIN PEROXIDASE

Chain G:  72% 16% 13%




- Molecule 1: THIOREDOXIN PEROXIDASE

Chain H:  70% 17% 11%




- Molecule 1: THIOREDOXIN PEROXIDASE

Chain I:  74% 13% 12%




- Molecule 1: THIOREDOXIN PEROXIDASE

Chain J:  76% 13% 11%




- Molecule 1: THIOREDOXIN PEROXIDASE

Chain K: 70% 16% 13%




- Molecule 1: THIOREDOXIN PEROXIDASE

Chain L: 69% 18% 12%




- Molecule 1: THIOREDOXIN PEROXIDASE

Chain M: 75% 13% 11%



- Molecule 1: THIOREDOXIN PEROXIDASE

Chain N: 72% 15% 12%



- Molecule 1: THIOREDOXIN PEROXIDASE


Chain O: 76% 11% 12%



GLY
GLN
HIS
GLY
ILE
LYS
VAL
GLN
LYS

• Molecule 1: THIOREDOXIN PEROXIDASE

Chain P: 73% 13% 11%



VAL
CYS
PRO
VAL
ASN
TRP
LYS
ARG
GLY
GLN
HIS
HIS
ILE
LYS
VAL
ASN
GLN
LYS

• Molecule 1: THIOREDOXIN PEROXIDASE


Chain Q: 2% 66% 18% 16%



D119, G120, M121, F123, I129, I134, L135, V145, R147, D150, K164, HIS, GLY, VAL, CYS, PRO, VAL, ASN, TRP, ARG, GLY, HIS, ILE, VAL, ASN, GLN, LYS

• Molecule 1: THIOREDOXIN PEROXIDASE

Chain R: 77% 10% 12%



ASN
GLN
LYS

• Molecule 1: THIOREDOXIN PEROXIDASE

Chain S: 73% 15% 12%




TRP
LYS
ARG
GLY
GLN
HIS
ILE
LYS
VAL
ASN
GLN
GLY
LYS

• Molecule 1: THIOREDOXIN PEROXIDASE


Chain T: 73% 16% 11%



VAL
CYS
PRO
VAL
ASN
TRP
LYS
ARG
GLY
GLN
HIS
ILE
LYS
VAL
ASN
GLN
LYS





- Molecule 1: THIOREDOXIN PEROXIDASE





- Molecule 1: THIOREDOXIN PEROXIDASE



- Molecule 1: THIOREDOXIN PEROXIDASE




- Molecule 1: THIOREDOXIN PEROXIDASE




- Molecule 1: THIOREDOXIN PEROXIDASE






● Molecule 1: THIOREDOXIN PEROXIDASE




● Molecule 1: THIOREDOXIN PEROXIDASE



● Molecule 1: THIOREDOXIN PEROXIDASE




● Molecule 1: THIOREDOXIN PEROXIDASE




● Molecule 1: THIOREDOXIN PEROXIDASE







● Molecule 1: THIOREDOXIN PEROXIDASE





● Molecule 1: THIOREDOXIN PEROXIDASE



● Molecule 1: THIOREDOXIN PEROXIDASE



● Molecule 1: THIOREDOXIN PEROXIDASE



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	95.18Å 190.30Å 212.00Å 90.00° 90.09° 90.00°	Depositor
Resolution (Å)	48.76 – 3.52 48.76 – 3.51	Depositor EDS
% Data completeness (in resolution range)	98.1 (48.76-3.52) 97.7 (48.76-3.51)	Depositor EDS
R_{merge}	0.28	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.03 (at 3.48Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.277 , 0.289 0.275 , 0.282	Depositor DCC
R_{free} test set	4589 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	95.0	Xtrriage
Anisotropy	0.536	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 66.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtrriage
Estimated twinning fraction	0.327 for h,-k,-l	Xtrriage
Reported twinning fraction	0.500 for -h,-k,l	Depositor
Outliers	2 of 91789 reflections (0.002%)	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	52692	wwPDB-VP
Average B, all atoms (Å ²)	111.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/1341	0.67	1/1813 (0.1%)
1	B	0.30	0/1374	0.63	1/1855 (0.1%)
1	C	0.30	0/1356	0.60	0/1833
1	D	0.30	0/1368	0.69	0/1849
1	E	0.30	0/1330	0.65	0/1798
1	F	0.27	0/1356	0.60	0/1833
1	G	0.29	0/1339	0.68	1/1810 (0.1%)
1	H	0.31	0/1356	0.69	0/1833
1	I	0.31	0/1350	0.69	0/1825
1	J	0.30	0/1363	0.64	0/1843
1	K	0.29	0/1321	0.66	0/1787
1	L	0.30	0/1363	0.68	3/1840 (0.2%)
1	M	0.30	0/1356	0.74	1/1833 (0.1%)
1	N	0.29	0/1368	0.72	1/1849 (0.1%)
1	O	0.29	0/1341	0.64	0/1813
1	P	0.28	0/1356	0.66	1/1833 (0.1%)
1	Q	0.29	0/1296	0.61	1/1747 (0.1%)
1	R	0.30	0/1345	0.63	0/1818
1	S	0.29	0/1350	0.74	1/1825 (0.1%)
1	T	0.28	0/1363	0.67	0/1843
1	U	0.31	0/1341	0.71	1/1813 (0.1%)
1	V	0.31	0/1374	0.74	2/1855 (0.1%)
1	W	0.29	0/1356	0.63	0/1833
1	X	0.29	0/1368	0.69	1/1849 (0.1%)
1	Y	0.31	0/1341	0.66	0/1813
1	Z	0.30	0/1356	0.69	1/1833 (0.1%)
1	a	0.33	0/1350	0.72	1/1825 (0.1%)
1	b	0.34	0/1356	0.74	1/1833 (0.1%)
1	c	0.29	0/1350	0.73	2/1825 (0.1%)
1	d	0.29	0/1363	0.67	2/1843 (0.1%)
1	e	0.30	0/1330	0.66	1/1798 (0.1%)
1	f	0.30	0/1374	0.70	3/1855 (0.2%)
1	g	0.31	0/1356	0.67	1/1833 (0.1%)
1	h	0.30	0/1357	0.72	0/1834

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	i	0.27	0/1298	0.55	0/1750
1	j	0.31	0/1356	0.68	1/1833 (0.1%)
1	k	0.31	0/1350	0.74	1/1825 (0.1%)
1	l	0.29	0/1356	0.68	0/1833
1	m	0.25	0/1296	0.53	0/1747
1	n	0.31	0/1363	0.66	0/1843
All	All	0.30	0/53983	0.67	29/72953 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	3
1	E	0	1
1	G	0	2
1	H	0	1
1	K	0	1
1	L	0	1
1	M	0	1
1	N	0	2
1	P	0	1
1	Q	0	2
1	R	0	1
1	S	0	2
1	T	0	1
1	U	0	1
1	V	0	4
1	W	0	1
1	X	0	1
1	Y	0	2
1	a	0	3
1	b	0	2
1	c	0	3
1	d	0	3
1	e	0	2
1	f	0	4
1	h	0	4
1	j	0	1
1	k	0	3

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
1	l	0	2
1	m	0	1
1	n	0	1
All	All	0	59

There are no bond length outliers.

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	N	48	PRO	C-N-CD	6.71	142.49	128.40
1	f	46	PHE	N-CA-C	-6.41	93.70	111.00
1	U	72	CYS	N-CA-C	6.09	127.46	111.00
1	P	133	GLY	N-CA-C	6.09	128.32	113.10
1	f	48	PRO	C-N-CD	-6.05	107.29	120.60

There are no chirality outliers.

5 of 59 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	21	GLU	Peptide
1	A	94	GLY	Peptide
1	B	105	LYS	Peptide
1	B	31	ARG	Peptide
1	B	43	ASP	Peptide

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	161/186 (87%)	132 (82%)	27 (17%)	2 (1%)	13	51
1	B	165/186 (89%)	145 (88%)	18 (11%)	2 (1%)	13	51
1	C	163/186 (88%)	141 (86%)	18 (11%)	4 (2%)	5	35
1	D	164/186 (88%)	148 (90%)	13 (8%)	3 (2%)	8	42
1	E	160/186 (86%)	134 (84%)	20 (12%)	6 (4%)	3	27
1	F	163/186 (88%)	140 (86%)	17 (10%)	6 (4%)	3	28
1	G	161/186 (87%)	135 (84%)	25 (16%)	1 (1%)	25	64
1	H	163/186 (88%)	131 (80%)	25 (15%)	7 (4%)	2	23
1	I	162/186 (87%)	131 (81%)	28 (17%)	3 (2%)	8	41
1	J	164/186 (88%)	144 (88%)	19 (12%)	1 (1%)	25	64
1	K	159/186 (86%)	133 (84%)	23 (14%)	3 (2%)	8	41
1	L	164/186 (88%)	140 (85%)	22 (13%)	2 (1%)	13	51
1	M	163/186 (88%)	137 (84%)	24 (15%)	2 (1%)	13	51
1	N	164/186 (88%)	143 (87%)	20 (12%)	1 (1%)	25	64
1	O	161/186 (87%)	139 (86%)	19 (12%)	3 (2%)	8	41
1	P	163/186 (88%)	134 (82%)	23 (14%)	6 (4%)	3	28
1	Q	154/186 (83%)	118 (77%)	29 (19%)	7 (4%)	2	22
1	R	162/186 (87%)	136 (84%)	25 (15%)	1 (1%)	25	64
1	S	162/186 (87%)	136 (84%)	23 (14%)	3 (2%)	8	41
1	T	164/186 (88%)	141 (86%)	20 (12%)	3 (2%)	8	42
1	U	161/186 (87%)	135 (84%)	21 (13%)	5 (3%)	4	32
1	V	165/186 (89%)	141 (86%)	21 (13%)	3 (2%)	8	42
1	W	163/186 (88%)	132 (81%)	28 (17%)	3 (2%)	8	42
1	X	164/186 (88%)	137 (84%)	23 (14%)	4 (2%)	6	36
1	Y	161/186 (87%)	133 (83%)	26 (16%)	2 (1%)	13	51
1	Z	163/186 (88%)	136 (83%)	25 (15%)	2 (1%)	13	51
1	a	162/186 (87%)	135 (83%)	24 (15%)	3 (2%)	8	41
1	b	163/186 (88%)	133 (82%)	28 (17%)	2 (1%)	13	51
1	c	162/186 (87%)	136 (84%)	23 (14%)	3 (2%)	8	41
1	d	164/186 (88%)	146 (89%)	17 (10%)	1 (1%)	25	64
1	e	160/186 (86%)	134 (84%)	21 (13%)	5 (3%)	4	32
1	f	165/186 (89%)	149 (90%)	14 (8%)	2 (1%)	13	51

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	g	163/186 (88%)	134 (82%)	25 (15%)	4 (2%)	5	35
1	h	163/186 (88%)	136 (83%)	25 (15%)	2 (1%)	13	51
1	i	154/186 (83%)	121 (79%)	31 (20%)	2 (1%)	12	49
1	j	163/186 (88%)	136 (83%)	24 (15%)	3 (2%)	8	42
1	k	162/186 (87%)	136 (84%)	23 (14%)	3 (2%)	8	41
1	l	163/186 (88%)	135 (83%)	23 (14%)	5 (3%)	4	32
1	m	154/186 (83%)	129 (84%)	24 (16%)	1 (1%)	25	64
1	n	164/186 (88%)	125 (76%)	32 (20%)	7 (4%)	2	23
All	All	6481/7440 (87%)	5437 (84%)	916 (14%)	128 (2%)	7	41

5 of 128 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	128	ILE
1	E	34	TYR
1	E	106	GLN
1	F	19	ASN
1	H	69	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/162 (88%)	119 (84%)	23 (16%)	2	14
1	B	146/162 (90%)	127 (87%)	19 (13%)	4	22
1	C	144/162 (89%)	124 (86%)	20 (14%)	3	21
1	D	145/162 (90%)	124 (86%)	21 (14%)	3	19
1	E	141/162 (87%)	127 (90%)	14 (10%)	8	34
1	F	144/162 (89%)	120 (83%)	24 (17%)	2	13
1	G	142/162 (88%)	117 (82%)	25 (18%)	2	11
1	H	144/162 (89%)	115 (80%)	29 (20%)	1	7

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	I	143/162 (88%)	121 (85%)	22 (15%)	2	17
1	J	145/162 (90%)	120 (83%)	25 (17%)	2	12
1	K	140/162 (86%)	113 (81%)	27 (19%)	1	8
1	L	145/162 (90%)	113 (78%)	32 (22%)	1	5
1	M	144/162 (89%)	120 (83%)	24 (17%)	2	13
1	N	145/162 (90%)	118 (81%)	27 (19%)	1	9
1	O	142/162 (88%)	121 (85%)	21 (15%)	3	18
1	P	144/162 (89%)	119 (83%)	25 (17%)	2	11
1	Q	137/162 (85%)	110 (80%)	27 (20%)	1	8
1	R	143/162 (88%)	125 (87%)	18 (13%)	4	23
1	S	143/162 (88%)	121 (85%)	22 (15%)	2	17
1	T	145/162 (90%)	119 (82%)	26 (18%)	2	10
1	U	142/162 (88%)	120 (84%)	22 (16%)	2	17
1	V	146/162 (90%)	121 (83%)	25 (17%)	2	12
1	W	144/162 (89%)	120 (83%)	24 (17%)	2	13
1	X	145/162 (90%)	124 (86%)	21 (14%)	3	19
1	Y	142/162 (88%)	117 (82%)	25 (18%)	2	11
1	Z	144/162 (89%)	119 (83%)	25 (17%)	2	11
1	a	143/162 (88%)	113 (79%)	30 (21%)	1	6
1	b	144/162 (89%)	124 (86%)	20 (14%)	3	21
1	c	143/162 (88%)	117 (82%)	26 (18%)	1	9
1	d	145/162 (90%)	119 (82%)	26 (18%)	2	10
1	e	141/162 (87%)	116 (82%)	25 (18%)	2	11
1	f	146/162 (90%)	115 (79%)	31 (21%)	1	6
1	g	144/162 (89%)	119 (83%)	25 (17%)	2	11
1	h	144/162 (89%)	117 (81%)	27 (19%)	1	8
1	i	137/162 (85%)	101 (74%)	36 (26%)	0	3
1	j	144/162 (89%)	115 (80%)	29 (20%)	1	7
1	k	143/162 (88%)	118 (82%)	25 (18%)	2	11
1	l	144/162 (89%)	112 (78%)	32 (22%)	1	5
1	m	137/162 (85%)	115 (84%)	22 (16%)	2	15

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	n	145/162 (90%)	123 (85%)	22 (15%)	3 17
All	All	5727/6480 (88%)	4738 (83%)	989 (17%)	2 11

5 of 989 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	T	130	ASP
1	j	154	ARG
1	Y	80	HIS
1	j	104	ARG
1	m	6	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 49 such sidechains are listed below:

Mol	Chain	Res	Type
1	a	68	GLN
1	g	95	HIS
1	b	121	ASN
1	f	165	HIS
1	i	58	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	163/186 (87%)	-0.27	1 (0%) 89 81	110, 111, 112, 112	0
1	B	165/186 (88%)	-0.20	1 (0%) 89 81	110, 111, 112, 112	0
1	C	165/186 (88%)	-0.09	1 (0%) 89 81	110, 111, 111, 112	0
1	D	164/186 (88%)	-0.13	0 100 100	109, 111, 111, 112	0
1	E	162/186 (87%)	-0.14	0 100 100	110, 111, 111, 112	0
1	F	165/186 (88%)	-0.22	1 (0%) 89 81	110, 111, 112, 112	0
1	G	162/186 (87%)	-0.21	0 100 100	110, 111, 112, 112	0
1	H	165/186 (88%)	-0.21	1 (0%) 89 81	110, 111, 112, 112	0
1	I	163/186 (87%)	-0.12	0 100 100	110, 111, 112, 112	0
1	J	166/186 (89%)	-0.17	0 100 100	110, 111, 112, 112	0
1	K	161/186 (86%)	-0.21	0 100 100	110, 111, 112, 112	0
1	L	164/186 (88%)	-0.20	2 (1%) 79 67	110, 111, 112, 112	0
1	M	165/186 (88%)	-0.22	0 100 100	110, 111, 112, 112	0
1	N	164/186 (88%)	-0.18	0 100 100	110, 111, 112, 112	0
1	O	163/186 (87%)	-0.08	1 (0%) 89 81	110, 111, 112, 112	0
1	P	165/186 (88%)	-0.19	0 100 100	110, 111, 112, 112	0
1	Q	157/186 (84%)	-0.12	3 (1%) 66 53	110, 111, 112, 112	0
1	R	164/186 (88%)	-0.24	0 100 100	110, 111, 112, 112	0
1	S	163/186 (87%)	-0.22	0 100 100	109, 111, 111, 112	0
1	T	166/186 (89%)	-0.13	0 100 100	109, 111, 111, 112	0
1	U	163/186 (87%)	-0.22	0 100 100	110, 111, 112, 112	0
1	V	165/186 (88%)	-0.18	0 100 100	110, 111, 112, 112	0
1	W	165/186 (88%)	-0.18	0 100 100	109, 111, 111, 112	0
1	X	164/186 (88%)	-0.22	0 100 100	109, 111, 111, 112	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	Y	163/186 (87%)	-0.23	1 (0%) 89 81	110, 111, 112, 112	0
1	Z	165/186 (88%)	-0.23	0 100 100	103, 111, 112, 112	0
1	a	163/186 (87%)	-0.21	1 (0%) 89 81	110, 111, 112, 112	0
1	b	165/186 (88%)	-0.19	0 100 100	110, 111, 112, 112	0
1	c	163/186 (87%)	-0.19	1 (0%) 89 81	110, 111, 112, 113	0
1	d	166/186 (89%)	-0.23	1 (0%) 89 81	110, 111, 112, 112	0
1	e	162/186 (87%)	-0.13	0 100 100	110, 111, 111, 112	0
1	f	165/186 (88%)	-0.14	0 100 100	110, 111, 111, 112	0
1	g	165/186 (88%)	-0.17	0 100 100	110, 111, 112, 112	0
1	h	163/186 (87%)	-0.16	1 (0%) 89 81	110, 111, 112, 113	0
1	i	158/186 (84%)	-0.15	2 (1%) 77 65	110, 111, 112, 112	0
1	j	165/186 (88%)	-0.07	2 (1%) 79 67	109, 111, 112, 112	0
1	k	163/186 (87%)	0.01	5 (3%) 49 36	110, 111, 112, 112	0
1	l	165/186 (88%)	-0.26	0 100 100	110, 111, 112, 112	0
1	m	157/186 (84%)	-0.10	2 (1%) 77 65	110, 111, 112, 112	0
1	n	166/186 (89%)	-0.25	1 (0%) 89 81	110, 111, 112, 112	0
All	All	6543/7440 (87%)	-0.18	28 (0%) 92 87	103, 111, 112, 113	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	h	14	GLY	5.8
1	a	45	THR	4.4
1	k	35	VAL	3.7
1	O	16	ALA	3.2
1	Q	99	PRO	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.